



Original article

Molecular characterization of goats from Saudi Arabia using microsatellite markers

Ahmed Hossam Mahmoud^{a,*}, Mohammad Abul Farah^a, Ahmed Rady^a, Khalid Mashai Alanazi^a, Osama Mohammed^b, Nabil Amor^b, Khaloud Mohammed Alarjani^c

^a Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia

^b KSU Mammals Research Chair, Department of Zoology, College of Science, King Saud University, Riyadh, Saudi Arabia

^c Department of Botany and Microbiology, College of Science, King Saud University, Riyadh, Saudi Arabia

ARTICLE INFO

Article history:

Received 23 November 2019

Revised 12 December 2019

Accepted 2 January 2020

Available online 9 January 2020

Keywords:

Saudi goats

Microsatellite marker

Population structure

Genetic variability

ABSTRACT

This study is undertaken for genetic characterization of local Saudi goat populations for maintaining genetic variability of livestock population. Genomic DNA was extracted from three goat populations namely Ardi, Hollandi and Shami and subjected for genotyping using eighteen recommended microsatellite markers. The average values obtained for number of private alleles, effective number of alleles and different alleles 2.071, 5.343 and 9.389, respectively. The expected and observed heterozygosity values were 0.757 and 0.913, respectively. F- values were 0.081, –0.115, –0.18 and Shannon's Information Index was found to be 1.751. All loci were highly differed from Hardy-Weinburg Equilibrium and an average migrant's value was 4.302. The maximum value was obtained between Hollandi and Ardi (8.474) and the value was extremely low between Shami and Ardi (2.518). Pairwise population of Fst and Nei's Genetic Distance showed very close relationship between Hollandi and Ardi goat population than between this population with the Shami. STRUCTURE software revealed 3 clusters at K = 3 in goat population. The present findings revealed genetic variations in Hollandi and Ardi populations. Molecular characterization of Saudi goats is important for maintaining and improving genetic resources.

© 2020 Published by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Domestication, a complex and gradual process, is a mutualism between human and associated animal resulting in around 40 domesticated species of animals to justify his needs. Goat is one of the earliest animals domesticated by human around 10,000 years ago (Groeneveld et al., 2010). Domestic goats (*Capra aegagrus hircus*) were widely distributed all over the world with more than 500 breeds for meat, milk, butter, cheese, skin and fiber productions and they are used as a brush control too as well as in maintaining rural populations and in various traditional, social feasts. Goat is the one of the important domestic ruminant among other domestic ruminants and can survive under wide range of climatic

conditions (Serrano et al., 2009). However, goats are very minimally considered by international community and several local breeds are facing possible extinction for different reasons. Thus, characterization of the genetic make-up of the endogenous breeds is an important issue for their optimum utilization (Galal, 2005) and avoiding the risk of loss of genetic variability among local goat breeds and their extinction through selection and inbreeding practices. Scientists recognized the need for the conservation of livestock resources. Therefore, studies on diversity and variability among local goat breeds were performed in many countries in Asia, Europe and Africa; with very few in the Middle Eastern countries even though, these countries hold around 54 goat breeds. Genetic diversity in livestock is a reservoir of traits that enables farmers to improve their stock and allows animals to adapt to changing conditions.

Molecular methods have been widely used to study the genetic variance among livestock. Polymerase chain reaction (PCR) technique is widely used to study the specific genetic material (DNA) in a controlled and logarithmic fashion. Numerous DNA markers have been used to study the genetic variance. Also, various types of genetic markers vary widely based on their content. In recent

* Corresponding author.

E-mail address: ahmahmoud@ksu.edu.sa (A. Hossam Mahmoud).

Peer review under responsibility of King Saud University.



years, microsatellites are widely used to study the genetic variations in animal population. These microsatellites regularly found throughout the genetic material, polymorphic in nature and are highly stable and very simple to analyze. Hence, application of genetic markers in farm animals could show a bottleneck method of analyzing the genetic makeup of an individual and has been reported earlier by [Blott et al. \(1998\)](#). Genetic diversity among goat populations using microsatellites has been earlier studied by [Cañón et al. \(2006\)](#). Recently, genetic diversity has been studied using microsatellite markers by various researchers ([Musthafa et al., 2018](#)).

In Saudi Arabia, goat population is very high and serves as an important source of meat and milk exceeding 2.2 million head producing 22,500 tons of meat and 76,500 tons of milk ([FAOSTAT, 2008](#)). They are very adapted to survive in low water resources, harsh climatic conditions and limited food availability. Ardi, Hollandi and Shami goats are three distinct Saudi Arabian goat breeds holding a special place in the Riyadh regional agribusiness economy ([Al-saeef, 2013](#)). The naming of these native goats suggests some variability.

In Saudi Arabia, Ardi goat population is very high than other breeds and plays a very important role in milk production. Hejazi (Pakistani, Indian or Hollandi) goats are usually black and long haired and used primarily for meat production. It is claimed that this breed was developed recently by the crossing Pakistani goat breed called “Kapla” breed native to a Pakistani province “Sindh” with two different local breeds, the “Ardi” and the “Cyprus Shami goats”. The Damascus goat, commonly known as Chami, Shami, Damascene, Baladi, Halep, Aleppo has unique shape of mouth and head raised originally in Lebanon, Cyprus and Syria and introduced to Saudi Arabia mainly for milk production. Few studies investigated the extent to which Saudi goat populations are genetically differentiated utilizing few or different microsatellite markers producing different outcomes, even with same breed ([Aljumaah et al., 2012](#)). Therefore, to preserve the genetic resources and to develop future comprehensive breeding programs in order to improve goat populations in Saudi Arabia, this study assessed the genetic variance between and within local goat populations in Riyadh region utilizing a set of 18 microsatellite markers.

2. Material and methods

2.1. Isolation of DNA and PCR amplification

Forty-eight Ardi (AR), 48 Hollandi (HO) and 36 Shami (SH) local goats were selected from a farm in Riyadh region of Saudi Arabia. About 10 ml of blood sample from each goat was collected aseptically and DNA was extracted from the sample using QIAgen DNeasy Kit according to the manufacturer's instructions. Nano-Drop 2000 spectrophotometer was used to analyze purity of extracted DNA and also quantified. A total of 18-microsatellite primer-pairs used to study the genetic variation which has been recommended previously to analyze the genetic variation of goats. A total of three markers were selected to study the level of polymorphism. DNA amplification was performed as described earlier by [Kumar et al.](#) The amplified DNA was analyzed using a spectrophotometer.

2.2. Statistical analyses

Heterozygosity analysis (H_E), observed heterozygosity (H_O), private alleles (N_p), effective number of alleles (N_e), observed number of alleles (N_a) and allele frequency were calculated for every population as suggested by [Kalinowski \(2007\)](#) using Cervus version 3.0.3 software. GenePop version 4.0.10 software was used to

analyze Wright's F-statistics, deviations from Hardy-Weinberg equilibrium (HWE), number of migrants exchanged between populations per generation as described previously by [Raymond and Rousset \(1995\)](#).

3. Results and discussion

Totally 18 microsatellites used in our study ([Table 1](#)) were found to be highly polymorphic with 3 (locus OarFCB128 of AR and HO, locus MAF209 of AR and locus OarJMP29 of SH) to 26 (locus MAF214 of HO) alleles per locus. This vast allelic diversity is very much useful for selection and show the suitability of these microsatellites for the analysis of diversity. The mean allelic value of 9.389 ([Table 2](#)) is higher than that reported for Iranian goats ([Mahmoudi, 2010](#)), in some Indian goat populations and lower than the value reported in some other goat populations in India ([Dixit et al., 2011](#)). For the same breeds, the mean number of alleles reported by [Al-Atiyat and Aljumaah \(2014\)](#) (8.25) for the Ardi goats and by [Al-Atiyat and Aljumaah \(2014\)](#) (7.25) for the Shami goats was lower than this of the same breeds reported in the present study (9.222 and 9.722 for Ardi and Shami goats, respectively). These differences may be due to different sample size and/or number and type of microsatellites used. Higher genetic diversity analyzed in this study might be also due to the larger effective number of alleles (5.343). This value compromised about 57% of the total number of alleles describing a large number of alleles at high frequency. A small number of effective alleles lead to a larger genetic drift from one generation to another. By population, SH had the highest number of private, effective and observed alleles as described in [Table 2](#). Also, Locus MAF214 had the highest number of observed, effective and private alleles. The percentage of loci with private alleles was 41.79 (112 out of 268). The private alleles accentuate the uniqueness of the population and help in genetic differentiation between populations. The Shannon's Information Index (I) assessments ([Table 3](#)) revealed that all the loci were very informative making them very useful for genetic diversity studies. The values varied as low as 0.0340 (OarJMP29 of SH) to as high as 2.881 (MAF214 of AR), indicating great heterogeneity for the population. The mean (I) value (1.751) observed in this study was found to be higher than that of results reported in Gaddi goats ([Singh et al., 2015](#)); in Korki Jonub Khorasan, and in Jakhrana goats, but lower than that reported in Mahabubnagar goats by [Raghavendra et al. \(2017\)](#).

Among various species and breeds or some times within breed heterozygosity level of a microsatellite varied considerably. All populations had qualitatively higher levels of genetic diversity confirmed by higher H_O and H_E values and described in [Table 3](#), making these microsatellites more suitable for analyzing genetic variation in goat population. Values of H_O were higher than of H_E values, indicating heterozygote excess in these populations, and belong to outbreeding systems as also indicated by the negative values of F_{is} ([Table 4](#)). In outbreeding, gene introgression and gene exchange is very common, which mainly increases heterogeneity in population ([Wang and Yue, 2008](#)). High rates of mutations at specific loci, and the large allele numbers detected could also be credited. The mean observed heterozygosity (0.913) was found to be higher than Asian goats populations, Korean goats, many Indian goats ([Raghavendra et al., 2017](#)); Sub-Saharan breeds, Spanish Guadarrama goats, West African dwarf goats, Croatian spotted breed and Albanian goats and was comparable with some Indian goat breeds and Sardinian goat population ([Sechi et al., 2005](#)). In the present study, the mean values of H_E for AR, HO and SH populations were 0.745, 0.774 and 0.751, respectively, with an overall mean of 0.757 throughout selected populations and markers. The highest value of 0.934 was for MAF214 locus in AR population

Table 1

Primers sequences and labels of the 18 primer pairs used to amplify microsatellite regions in the three Saudi local goat populations of the present study.

Locus Name	Sequences 5' → 3' Forward/Reverse	Label	Allele Size (bp)	Chromosomal Location
ILSTS005	GGAAGCAATGAAATCTATAGCC TGTTCTGTGAGTTTGAAGC	56FAM	174–218	7
MCM527	GTCCATTGCCTCAAATCAATC AAACCACTTGACTACTCCCAA	56-TAMN	165–187	5
SRCRSP5	GGACTCTACCAACTGAGCTACAAG GTTTCTTGAATGAAGCTAAAGCAATGC	5HEX	126–158	18
OarFCB128	ATTAAGCATCTTCTTTATTTCTCTGC CAGCTGAGCAACTAAGACATACATGCG	56FAM	96–130	2
HUJ616	TTCAAACACACATTGACAGGG GGACCTTTGGCAATGGAAGG	56-ROXN	114–160	13
OarHH47	TTTATTGACAAAACCTCTTCTAACTCCAC GTAGTTATTTAAAAAATATCATACTCTTAAGG	56-TAMN	130–152	18
ILSTS11	GCTTGCTACATGAAAGTGC CTAAAATGCAGAGCCCTACC	56FAM	256–294	9
DYMS1	AACAACATCAAACAGTAAGAG CATAGTAACAGATCTTCTTACA	56-TAMN	159–211	20
BM8024	CTCTATCTGTGGAAAAGGTGGG GGGGTTAGACTTCAACATACG	56-TAMN	110–130	1
OarFCB226	CTATATGTTGCCTTCCCTTCTCTGC GTGAGTCCCATAGAGCATAAGCTC	5-HEX	119–153	2
OarAE129	AATCCAGTGTGTGAAAGACTAATCCAG GTAGATCAAGATATAGAATATTTTCAACACC	56FAM	133–159	5
OarJMP29	GTATACACGTGGACACCGCTTTGTAC GAAGTGGCAAGATTGAGGGGAAAG	56-ROXN	96–150	24
SRCRSP9	AGAGGATCTGGAAATGGAATC GCACCTTTTTCAGCCCTAATG	56FAM	99–135	12
MAF214	GGGTGATCTTAGGAGGTTTGGAGG AATGCAGGAGATCTGAGGCAGGGACG	56FAM	174–282	16
OarCP34	GCTGAACAATGTGATATGTTTCAGG GGGACAATACTGTCTTAGATGCTGC	56-ROXN	112–130	3
OarFCB304	CCCTAGGAGCTTTCAATAAAGAATCCG CGCTGCTGTCAACTGGGTCAGGG	5-HEX	150–188	19
MAF209	GATCACAAAAGTTGGATACAACCGTGG TCATGCACTTAAGTATGTAGGATGCTG	5-HEX	109–135	17
MAF65	AAAGGCCAGATGCAATTAGGAG CCACTCCTCTGAGAATATAACATG	56-TAMN	123–163	15

Table 2

Number of different (Na), effective (Ne) and private (Np) alleles across 18 loci and three local goat populations.

Population*	AR			HO			SH			OVERALL		
	Na	Ne	Np	Na	Ne	Np	Na	Ne	Np	Na	Ne	Np
ILSTS005	6.000	3.454	0	4.000	3.034	1	15.000	5.378	9	8.333	3.955	3.333
MCM527	10.000	7.591	1	10.000	5.738	0	12.000	8.100	3	10.667	7.143	1.333
SRCRSP5	8.000	5.109	0	5.000	3.356	0	14.000	6.698	8	9.000	5.054	2.667
OarFCB128	3.000	2.593	0	3.000	2.571	0	7.000	4.334	7	4.333	3.166	2.333
HUJ616	4.000	2.768	0	5.000	3.060	0	10.000	3.661	7	6.333	3.163	2.333
OarHH47	14.000	10.716	3	11.000	8.056	0	4.000	2.114	0	9.667	6.962	1.000
ILSTS11	14.000	7.223	4	11.000	6.727	0	5.000	3.236	1	10.000	5.729	1.667
DYMS1	16.000	7.745	5	9.000	6.555	0	10.000	5.366	1	11.667	6.555	2.000
BM8024	10.000	5.103	2	8.000	5.969	0	5.000	3.951	0	7.667	5.008	0.667
OarFCB226	4.000	2.522	2	9.000	2.994	5	6.000	3.082	0	6.333	2.866	2.333
OarAE129	12.000	6.391	4	8.000	4.078	1	8.000	5.709	3	9.333	5.393	2.667
OarJMP29	6.000	2.783	1	6.000	2.793	1	3.000	1.185	1	5.000	2.253	1.000
SRCRSP9	10.000	5.696	0	11.000	8.502	1	9.000	6.014	1	10.000	6.737	0.667
MAF214	23.000	15.208	5	26.000	10.263	7	18.000	6.949	6	22.333	10.807	6.000
OarCP34	3.000	1.743	0	11.000	4.594	4	7.000	5.043	0	7.000	3.793	1.333
OarFCB304	13.000	8.113	1	10.000	6.518	0	15.000	6.968	2	12.667	7.199	1.000
MAF209	3.000	1.743	0	11.000	4.465	4	8.000	4.713	1	7.333	3.640	1.667
MAF65	7.000	4.024	0	8.000	4.876	0	19.000	11.368	10	11.333	6.756	3.333
Mean	9.222	5.585	1.556	9.222	5.230	1.333	9.722	5.215	3.333	9.389	5.343	2.074
SE	1.282	0.825	0.437	1.165	0.524	0.505	1.131	0.557	0.820	0.677	0.369	0.587

*AR: Ardi, HO: Hollandi and SH: Shami.

which was lowest (0.156) in OarJMP29 locus of SH population, and the. The expected heterozygosity values obtained in our study is similar with the findings of [Serrano et al. \(2009\)](#) for Spanish Guadarrama goats, in Namibian goats, [Agha et al. \(2008\)](#) in Egyptian goats, [Els et al. \(2004\)](#) in Namibian goats,

The F_{st} values of the goat population varied between 0.022 and 0.177, showing various degrees of variation in genetic level between the different populations. This differentiation either moderate or negligible. The mean value of F_{st} indicated that between population the variation is very low in genetic level (8.1%) and

Table 3
Shannon's Information Index (I), Observed (H_O) and Expected (H_E) heterozygosities across 18 loci and three Saudi local goat populations.

Population*	AR			HO			SH			ALL		
	I	H _O	H _E	I	H _O	H _E	I	H _O	H _E	I	H _O	H _E
ILSTS005	1.465	1.000	0.711	1.226	1.000	0.670	2.196	0.750	0.814	1.629	0.917	0.732
MCM527	2.119	1.000	0.868	1.939	0.833	0.826	2.269	0.889	0.877	2.109	0.907	0.857
SRCRSP5	1.790	0.896	0.804	1.306	0.938	0.702	2.221	0.944	0.851	1.772	0.926	0.786
OarFCB128	1.018	1.000	0.614	1.011	1.000	0.611	1.650	1.000	0.769	1.226	1.000	0.665
HUJ616	1.134	1.000	0.639	1.329	1.000	0.673	1.747	1.000	0.727	1.403	1.000	0.680
OarHH47	2.478	0.896	0.907	2.210	0.938	0.876	0.820	1.000	0.527	1.836	0.944	0.770
ILSTS11	2.205	0.958	0.862	2.121	1.000	0.851	1.268	1.000	0.691	1.865	0.986	0.801
DYMS1	2.286	0.771	0.871	2.005	0.917	0.847	1.944	0.972	0.814	2.078	0.887	0.844
BM8024	1.828	1.000	0.804	1.864	1.000	0.832	1.496	1.000	0.747	1.729	1.000	0.794
OarFCB226	1.044	0.917	0.604	1.582	0.896	0.666	1.352	1.000	0.676	1.326	0.938	0.648
OarAE129	2.067	0.958	0.844	1.637	1.000	0.755	1.897	1.000	0.825	1.867	0.986	0.808
OarJMP29	1.224	0.896	0.641	1.257	0.813	0.642	0.340	0.111	0.156	0.940	0.606	0.479
SRCRSP9	1.966	1.000	0.824	2.258	1.000	0.882	1.958	1.000	0.834	2.061	1.000	0.847
MAF214	2.881	0.958	0.934	2.757	0.958	0.903	2.318	1.000	0.856	2.652	0.972	0.898
OarCP34	0.751	0.542	0.426	1.841	0.896	0.782	1.751	0.917	0.802	1.448	0.785	0.670
OarFCB304	2.241	0.979	0.877	2.016	0.938	0.847	2.223	0.944	0.856	2.160	0.954	0.860
MAF209	0.751	0.542	0.426	1.826	0.917	0.776	1.736	0.917	0.788	1.438	0.792	0.663
MAF65	1.583	0.813	0.752	1.744	0.750	0.795	2.621	0.944	0.912	1.983	0.836	0.819
Mean	1.713	0.896	0.745	1.774	0.933	0.774	1.767	0.910	0.751	1.751	0.913	0.757
SE	0.147	0.034	0.037	0.104	0.017	0.022	0.133	0.049	0.041	0.073	0.021	0.019

*AR: Ardi, HO: Hollandi and SH: Shami.

Table 4
F-statistics analysis and estimate of number of migrants (Nm) for the 18 microsatellite loci used for genotyping the three Saudi local goat populations.

	*Fis	Fit	Fst	Nm
ILSTS005	-0.253	-0.131	0.097	2.327
MCM527	-0.059	-0.028	0.029	8.370
SRCRSP5	-0.179	-0.058	0.102	2.205
OarFCB128	-0.504	-0.238	0.177	1.164
HUJ616	-0.472	-0.242	0.156	1.355
OarHH47	-0.227	-0.079	0.121	1.821
ILSTS11	-0.231	-0.129	0.083	2.780
DYMS1	-0.050	0.000	0.048	4.971
BM8024	-0.259	-0.207	0.041	5.784
OarFCB226	-0.446	-0.354	0.064	3.676
OarAE129	-0.221	-0.172	0.040	5.962
OarJMP29	-0.265	-0.121	0.114	1.949
SRCRSP9	-0.181	-0.142	0.033	7.404
MAF214	-0.083	-0.054	0.027	8.932
OarCP34	-0.171	-0.013	0.135	1.602
OarFCB304	-0.109	-0.084	0.022	10.975
MAF209	-0.193	-0.062	0.110	2.021
MAF65	-0.020	0.038	0.057	4.133
Mean	-0.218	-0.115	0.081	4.302
SE	0.033	0.023	0.011	0.703

*Fis: the inbreeding coefficient for an individual relative to the total population; Fit: the inbreeding coefficient for an individual relative to a subpopulation; Fst: the inbreeding coefficient for a subpopulation relative to the total population. Nm: Gene flow estimated from F_{ST} , $Nm = 0.25(1 - F_{ST})/F_{ST}$.

within breeds, variation is very high in genetic level (91.9%) corresponding to differences among individuals. The high number of migrants (Nm) for some loci exchanged between populations per generation (Table 4), indicated some degree of gene flow corresponding to the lower individual loci values of Fst (Fst < 0.05) which may avoid genetic drift due to the results of genetic variation. The highest genetic differentiation Fst value of 0.177 (locus OarFCB128) was recorded at the least number of migrants (1.164). The highest Nm value (10.975) and the least genetic differentiation coefficient (0.022) were recorded for locus OarFCB304. The number of migrations (Nm) between AR and HO (8.474) was higher than that between HO and SH (3.519) and between AR and SH (2.518) as Fst values reached 0.029, 0.066 and 0.090, respectively (Table 6). However, the overall value of 4.302 for

Nm (Table 4) suggests some sharing in genetic level among goats population. This intermixing may be due to the results of these goats originate from the same region or bought from the same market or neighbor farm or some uncontrollable breeding among different breeds of the same farm. However, the three Saudi goat populations should all be considered as separate breeds since all pair wise Fst values were highly significant ($P < 0.001$). Most of goat studies indicated low genetic differentiation (Hoda et al., 2011).

Based on the analysis of Nei's unbiased genetic distance the pair wise population matrix (Table 5), goats from AR and HO are closely related (0.166), more than to goats from SH. The most distance was AR from SH population (0.789), with some relation between HO and SH populations (0.547). This was confirmed with the Nei's genetic identity matrix where goats sampled from AR population were 84.7% identical to those in HO population. AR and SH goats were only 45.4% identical to those in SH which is a moderate value though the least identical value. HO goats were 57.9% identical to those in SH population. The same results are reflected in the pair wise population Fst matrix represented in Table 6. All the populations were showed deviation from the equilibrium of Hardy-

Table 5
Pairwise Population Matrix of Nei's unbiased Genetic Distance (lower left diagonal) (Nei, 1972) and Genetic Identity (upper right diagonal) between the Three Saudi Local Goat Populations.

Population	AR	HO	SH
AR	-	0.847	0.454
HO	0.166	-	0.579
SH	0.789	0.547	-

*AR: Ardi, HO: Hollandi and SH: Shami.

Table 6
Pairwise Population Fst Values (lower left diagonal) and Number of Migrants (Nm) (upper right diagonal) between the Three Saudi Local Goat Populations.

Population	AR	HO	SH
AR	0.000	8.474	2.518
HO	0.029	0.000	3.519
SH	0.090	0.066	0.000

*AR: Ardi, HO: Hollandi and SH: Shami.

Weinberg (Table 7; $P < 0.05$) indicating a low level of inbreeding ($F_{is} < 0.05$). Only two loci (OarCP34 and MAF209) in AR and one locus (OarFCB226) in HO and (HUIJ616) in SH were found to be

Hardy–Weinberg equilibrium and was statistically significant ($p > 0.05$). Seven of the total fourteen microsatellite markers used in Aljumaah et al. (2012) study of Ardi goat population of Saudi

Table 7
Number of loci significantly deviating from Hardy–Weinberg equilibrium (HWE).

Population*	AR				HO				SH			
	Locus	DF	ChiSq	Prob	Signif [#]	DF	ChiSq	Prob	Signif	DF	ChiSq	Prob
ILSTS005	15	81.488	0.000	***	6	42.688	0.000	***	105	185.639	0.000	***
MCM527	45	172.173	0.000	***	45	232.103	0.000	***	66	133.275	0.000	***
SRCRSP5	28	125.273	0.000	***	10	30.563	0.001	***	91	304.306	0.000	***
OarFCB128	3	48.000	0.000	***	3	48.000	0.000	***	21	46.903	0.001	***
HUIJ616	6	48.000	0.000	***	10	48.000	0.000	***	45	48.754	0.325	ns
OarHH47	91	157.938	0.000	***	55	192.603	0.000	***	6	108.000	0.000	***
ILSTS11	91	189.482	0.000	***	55	201.392	0.000	***	10	23.117	0.010	*
DYMS1	120	293.155	0.000	***	36	135.102	0.000	***	45	198.296	0.000	***
BM8024	45	91.589	0.000	***	28	59.935	0.000	***	10	43.283	0.000	***
OarFCB226	6	34.367	0.000	***	36	31.596	0.678	ns	15	38.057	0.001	***
OarAE129	66	144.690	0.000	***	28	59.188	0.001	***	28	104.836	0.000	***
OarJMP29	15	70.754	0.000	***	15	39.858	0.000	***	3	36.132	0.000	***
SRCRSP9	45	81.767	0.001	***	55	136.029	0.000	***	36	86.771	0.000	***
MAF214	253	589.072	0.000	***	325	625.847	0.000	***	153	417.943	0.000	***
OarCP34	3	6.622	0.085	ns	55	147.092	0.000	***	21	53.294	0.000	***
OarFCB304	78	345.921	0.000	***	45	255.403	0.000	***	105	204.516	0.000	***
MAF209	3	6.622	0.085	ns	55	149.758	0.000	***	28	61.770	0.000	***
MAF65	21	38.755	0.010	*	28	55.091	0.002	**	171	288.310	0.000	***

*AR: Ardi, HO: Hollandi and SH: Shami.
Key: ns = not significant, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Fig. 1. Scatter-plot of the factorial correspondence analysis (FCA) based on allele frequency of goat populations. Ardi genotypes (closed yellow squares); Hollandi genotypes (closed blue squares); Shami genotypes (empty squares).




Fig. 2. Determination of the best number of clusters from STRUCTURE analysis for microsatellite loci in goat populations.




Fig. 3. Bar plots from inferred population structure using the Bayesian grouping admixture model-based program STRUCTURE (K = 3).

Arabia, and 6 out of 11 markers in the goat population of Moxotó dairy and Alpine Saanen goat in Brazil showed Hardy Weinberg Equilibrium (HWE). The huge difference from HWE observed on goats of the present study could be due to the presence of alleles, called “null” alleles, size of homoplasy of microsatellite loci, high mutation rate and also because of small sample size (Araújo et al., 2006). However, this deviation could be related to excess heterozygous individuals than homozygous ones as noticed from higher H_0 than H_E values reported in the present study. Factorial correspondence analysis (FCA) placed on 2-dimensional plane was used to analyze the relation between the individuals in genetic level used in the present study based on their allele frequencies and described in Fig. 1. The individuals of Shami, Hollandi and Ardi goat populations usually have their own groups. The first and second principle components (PC) explained 7.61% of the total difference and widely between these clusters. STRUCTURE analysis was carried out to elucidate the substructures of breeds within the population. Based on STRUCTURE analysis with no prior distribution specified, K = 3 had the highest ΔK ($\Delta K = 1821.25$) and 3 was optimal value for K (Fig. 2). This analysis showed a strong signature of genetic structure grouping the accessions into three well differentiated clusters (Fig. 3) corresponding to the 3 goat populations. Concerning population demography fluctuations, Bottleneck results showed significant heterozygote excess within AR and HO populations at IAM and TPM mutation models ($p < 0.01$) indicating and Cornuet and Luikart (1996) previously earlier reported bottle neck. TPM is considered more appropriate for microsatellite analysis. Our results displayed no significant genetic bottleneck effect in the population of SH goat.

4. Conclusion

The selected microsatellite markers are highly useful for the analysis of genetic variation in Saudi goats. Good breeding strategy is helpful to maintain heterozygosity in the goat population. The present findings displayed no significant genetic bottleneck effect in the population of SH goat in Saudi Arabia.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

The authors extend their appreciation to The Researchers Supporting Project number (RSP-2019/108) King Saud University, Riyadh, Saudi Arabia.

References

- Agha, S.H., Pilla, F., Galal, S., Shaat, I., D'Andrea, M., Reale, S., Abdelsalam, A.Z.A., Li, M.H., 2008. Genetic diversity in Egyptian and Italian goat breeds measured with microsatellite polymorphism. *J. Ani. Breed. Gen.* 125, 194–200.
- Al-Atiyat, R., Aljumaah, R., 2014. Genetic relatedness between Ardi, Black Bedouin and Damascus goat breeds. *Genet. Mol. Res.* 13, 4654–4665.
- Aljumaah, R.S., Musthafa, M.M., Al-Shaikh, M.A., Badri, O.M., Hussein, M.F., 2012. Genetic diversity of Ardi goat based on microsatellite analysis. *Afr. J. Biotech.* 11, 16539–16545.
- Al-Saeef, A., 2013. Genetic and phenotypic parameters of body weights in Saudi Aradi goat and their crosses with Syrian Damascus goat. *Small Rum. Res.* 112, 35–38.
- Araújo, A.M., Guimarães, S.E.F., Machado, T.M.M., Lopes, P.S., Pereira, C.S., Silva, F.R., Rodrigues, M.T., Columbiano, V., Fonseca, C.G., 2006. Genetic diversity between herds of Alpine and Saanen dairy goats and the naturalized Brazilian Moxotó breed. *Gen. Mol. Biol.* 29, 67–74.
- Blott, S., Williams, J., Haley, C., 1998. Genetic variation within the Hereford breed of cattle. *Ani. Genet.* 29, 202–211.
- Cañón, J., García, D., García-Atance, M.A., Obexer-Ruff, G., Lenstra, J.A., Ajmone-Marsan, P., Dunner, S., 2006. Geographical partitioning of goat diversity in Europe and the Middle East. *Ani. Genet.* 37, 327–334.
- Cornuet, J.M., Luikart, G., 1996. Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics* 144, 2001–2014.
- Dixit, S. et al., 2011. Genetic variability and bottleneck analyses of Kanniadu goat breed based on microsatellite markers. *Indian J. Anim. Sci.* 81, 40.
- Els, J., Kotze, A., Swart, H., 2004. Genetic diversity of indigenous goats in Namibia using microsatellite markers: preliminary results. *South Afr. J. Ani. Sci.* 34 (6), 65–67.
- FAOSTAT, 2008. <http://faostat.fao.org/default.aspx>.
- Galal, S., 2005. Biodiversity in goats. *Small Rumin. Res.* 60, 75–81.
- Groeneveld, L., Lenstra, J.A., Eding, H., Toro, M.A., Scherf, B., Pilling, D., Negrini, R., Finlay, E.K., Jianlin, H., Groeneveld, E., Weigend, S., 2010. Genetic diversity in farm animals—a review. *Anim. Genet.* 41, 6–31.
- Hoda, A., Hyka, G., Boxberger, S.D., Obexer-Ruff, G., Consortium, E., 2011. Genetic diversity of Albanian goat breeds based on microsatellite markers. *Arch. Zootec.* 60, 607–615.
- Kalinowski, S.T., 2007. How many alleles per locus should be used to estimate genetic distances? *Heredity* 88, 62–65.
- Mahmoudi, B., 2010. Genetic diversity of Lori goat population based on microsatellite marker. *Asian-Australas J. Anim. Sci.* 4 (1), 13–19.
- Musthafa, M.M., Hussain, T., Babar, M.E., Aljumaah, R.S., Alshaikh, M.A., Muritala, I., Landi, V., Martinez, A., Amills, M., Dadi, O., Delgado, J.V., Aina, A.B.J., Onasoga, A. A., Adebambo, O.A., Visser, C., Van Marle-Köster, E., Adebambo, A.O., Marikar, F. M.M.T., 2018. Usability of metadata analysis of goat genetic resources among five countries from Africa, Asia and Europe: metadata analysis of goat genetic. *Agr. Sci. Biotech.* 10 (3), 183–190.
- Raghavendra, C., Saireddy, S., Amareswari, P., Raghunandan, T., 2017. Molecular characterization of Mahabubnagar goats based on microsatellite markers. *The Pharma Innov. J.* 6 (5), 130–133.
- Raymond, M., Rousset, F., 1995. Genepop (Version-1.2) - population-genetics software for exact tests and ecumenicism. *J. Hered.* 86, 248–249.
- Sechi, T., Usai, M.G., Casu, S., Carta, A., 2005. Genetic diversity of Sardinian goat population based on microsatellites. *Ital. J. Anim. Sci.* 4, 58–60.
- Serrano, M., Calvo, J.H., Martínez, M., Marcos-Carcavilla, A., Cuevas, J., González, C., Jurado, J.J., de Tejada, P.D., 2009. Microsatellite based genetic diversity and population structure of the endangered Spanish Guadarrama goat breed. *BMC Genet.* 10, 61.
- Singh, G., Thakur, Y., Kour, A., Sankhyan, V., Katoch, S., 2015. Genetic characterization of Gaddi goat breed of Western Himalayas using microsatellite markers. *Vet. World* 8, 527–531.
- Wang, J.M., Yue, W.B., 2008. Genetic relationships of domestic sheep and goats in the lower reaches of the Yellow River based on microsatellite analysis. *Biodiv. Sci.* 16, 53–62.