



# A nonlocal diffusion problem that approximates the heat equation with Neumann boundary conditions<sup>☆</sup>



Cesar A. Gómez<sup>a,\*</sup>, Julio D. Rossi<sup>b</sup>

<sup>a</sup> Department of Mathematics, National University of Colombia, Bogotá, Colombia

<sup>b</sup> Departamento de Matemática, FCEyN, Universidad de Buenos Aires, Ciudad Universitaria. Pab 1, 1428 Buenos Aires, Argentina

## ARTICLE INFO

### Article history:

Received 1 August 2017

Accepted 19 August 2017

Available online 24 August 2017

### 2010 Mathematics Subject Classification:

45A05

45J05

35K05

### Keywords:

Nonlocal diffusion

Neumann boundary conditions

Heat equation

## ABSTRACT

In this paper we discuss a nonlocal approximation to the classical heat equation with Neumann boundary conditions. We consider

$$w_t^\epsilon(x, t) = \frac{1}{\epsilon^{N+2}} \int_{\Omega} J\left(\frac{x-y}{\epsilon}\right) (w^\epsilon(y, t) - w^\epsilon(x, t)) dy + \frac{c}{\epsilon} \int_{\partial\Omega} J\left(\frac{x-y}{\epsilon}\right) g(y, t) dS_y, \quad (x, t) \in \bar{\Omega} \times (0, T),$$

$$w(x, 0) = u_0(x), \quad x \in \bar{\Omega},$$

and we show that the corresponding solutions,  $w^\epsilon$ , converge to the classical solution of the local heat equation  $v_t = \Delta v$  with Neumann boundary conditions,  $\frac{\partial v}{\partial n}(x, t) = g(x, t)$ , and initial condition  $v(0) = u_0$ , as the parameter  $\epsilon$  goes to zero. The obtained convergence is in the weak star on  $L^\infty$  topology.

© 2017 The Authors. Production and hosting by Elsevier B.V. on behalf of King Saud University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

The nonlocal evolution equation

$$u_t(x, t) = \int_{\Omega} J(x-y)[u(y, t) - u(x, t)] dy, \quad (x, t) \in \Omega \times (0, T), \quad (1.1)$$

$$u(x, 0) = u_0(x), \quad x \in \Omega,$$

see (Fife (2003)), can be seen as similar to the local heat equation with homogeneous Neumann boundary conditions

$$\begin{aligned} v_t(x, t) &= \Delta v(x, t), \quad (x, t) \in \Omega \times (0, T), \\ \frac{\partial v}{\partial n}(x, t) &= 0, \quad (x, t) \in \partial\Omega \times (0, T), \\ v(x, 0) &= u_0(x), \quad x \in \Omega. \end{aligned} \quad (1.2)$$

Here,  $J$  is a symmetric continuous nonnegative real function defined on  $\mathbb{R}^N$ , compactly supported in the unit ball, and such that  $\int_{\mathbb{R}^N} J(x) dx = 1$ ,  $\Omega$  is a bounded smooth domain in  $\mathbb{R}^N$  and  $u_0$  stands

<sup>☆</sup> Supported by MEC MTM2010-18128 and MTM2011-27998 (Spain).

\* Corresponding author.

E-mail addresses: [cagomezsi@una.edu.co](mailto:cagomezsi@una.edu.co) (C.A. Gómez), [jrossi@dm.uba.ar](mailto:jrossi@dm.uba.ar) (J.D. Rossi).

Peer review under responsibility of King Saud University.



Production and hosting by Elsevier

for the initial condition. The problem is known as a nonlocal model since the diffusion of the density  $u$  at a point  $x$  and time  $t$  not only depends on  $u(x, t)$  locally, but also on all values of  $u$  through the convolution like term  $\int_{\Omega} J(x-y)u(y, t) dy$ . Following (Fife, 2003), the model (1.1) can be interpreted as follows: If  $u(x, t)$  is the density of a population at point  $x$  and time  $t$ , and  $J(x-y)$  is thought as the probability distribution of jumping from location  $y$  to location  $x$ , then the convolution  $\int_{\Omega} J(y-x)u(y, t) dy$ , is the rate at which the individuals are arriving to location  $x$  from all other places  $y \in \Omega$  (notice that no individuals may arrive to  $x$  coming from outside  $\Omega$ ). In the same way,  $-\int_{\Omega} J(y-x)u(x, t) dy$ , is the rate at which individuals are leaving the location  $x$  to travel to other sites  $y \in \Omega$  (notice that no individual can jump outside  $\Omega$ ). So, in absence of external or internal sources, the density  $u$  satisfies the nonlocal Eq. (1.1). Now we remark that the fact that there is no individuals that enter or leave the domain makes this problem a zero flux diffusion problem and therefore the total mass is preserved,  $\int_{\Omega} u(x, t) dx = \int_{\Omega} u_0(x) dx$ , as happens with solutions to (1.2). The model (1.1) shares more properties with (1.2) such as: bounded stationary solutions are constant, a maximum principle is satisfied and perturbations propagate with infinite speed (Fife, 2003). Concerning applications, (1.1) and some variants of it have been used, for instance, in Biology (Carrillo and Fife, 2005), image processing, Gilboa and Osher (2006), and particle systems, Bodnar and Velázquez (2006); see also (Alberti and Bellettini, 1998 and

Fournier and Laurencot, 2006). For the mathematical analysis of nonlocal models the list of references is large and we refer to Abu Arqub et al. (2015), El-Ajou et al. (2015a,b), Andreu et al. (2008), Bates et al. (1997), Bogoya and Gómez (2012), Caffarelli and Silvestre (2009), Chasseigne et al. (2006), Coville et al. (2008), Coville and Dupaigne (2007), Ignat and Rossi (2007), and to the book (Andreu-Vaillio et al., 2010) and references therein.

Moreover, when one rescales the kernel  $J$  considering  $J(\xi) = \frac{1}{\epsilon^{N+2}} J(\frac{\xi}{\epsilon})$ , it was shown in Cortazar et al. (2008) that the corresponding solutions to (1.1) with a fixed initial condition converge to the solution to (1.2) as  $\epsilon \rightarrow 0$ . In addition, concerning the non-homogeneous problem, that is, (1.2) with  $\frac{\partial v}{\partial n}(x, t) = g(x, t)$ , in Cortazar et al. (2008) it is proved that this problem can be approximated with

$$(u^\epsilon)_t(x, t) = \frac{1}{\epsilon^{N+2}} \int_{\Omega} J(\frac{x-y}{\epsilon}) (u^\epsilon(y, t) - u^\epsilon(x, t)) dy + \frac{K}{\epsilon^{N+1}} \int_{\mathbb{R}^N \setminus \Omega} J(\frac{x-y}{\epsilon}) g(y, t) dy, \tag{1.3}$$

$$u^\epsilon(x, 0) = u_0(x),$$

as  $\epsilon \rightarrow 0$  (with an appropriate choice of the constant  $K$ ). Notice that in this model there are individuals that enter the domain coming from outside (this is the meaning of the integral in  $\mathbb{R}^N \setminus \Omega$ ). This model has the following disadvantage: when one tries to approximate solutions to (1.2) with  $\frac{\partial v}{\partial n}(x, t) = g(x, t)$  the datum  $g(x, t)$  is given only for  $x \in \partial\Omega$  and therefore one has to extend it outside (to a strip around  $\partial\Omega$  inside  $\mathbb{R}^N \setminus \Omega$  of width of order  $\epsilon$ ) in order to consider (1.3). An easy way to obtain such extension is to consider  $g(\tilde{y} + n(\tilde{y})s) = g(\tilde{y})$  where  $\tilde{y} \in \partial\Omega$  and  $s \in [0, \delta]$  (with  $\delta$  small), being  $n$  the exterior normal vector to  $\partial\Omega$  at  $\tilde{y}$ . To overcome the fact that an extension of the Neumann datum  $g$  outside  $\Omega$  is needed, recently, following (Cortazar et al., 2007), the authors of Bogoya and Gómez (2012) have introduced the nonlocal diffusion model

$$u_t(x, t) = \int_{\Omega} J(x-y)(u(y, t) - u(x, t)) dy + \int_{\partial\Omega} G(x-y)g(y, t) dS_y, \tag{1.4}$$

$$(x, t) \in \bar{\Omega} \times (0, T),$$

$$u(x, 0) = u_0(x), x \in \bar{\Omega},$$

where  $\Omega$  is a bounded domain,  $J : \mathbb{R}^N \mapsto \mathbb{R}$  and  $G : \mathbb{R}^N \mapsto \mathbb{R}$  are continuous, nonnegative, radially symmetric functions compactly supported in the unit ball and such that  $\int_{\mathbb{R}^N} J(z) dz = 1$ ,  $\int_{\mathbb{R}^N} G(z) dz = 1$ , and  $g \in L^\infty_{loc}([0, \infty); L^1(\partial\Omega))$ .

**Remark 1.1.** The main advantage of the model given by (1.4) compared with (1.3) is that when one deals with a nonlinear datum of the form  $g(y, t) = f(u(y, t))$  it is necessary to use an extension of the solution  $u$  from  $\Omega$  to  $\mathbb{R}^N \setminus \Omega$  in the case of (1.3) (notice that such an extension is not trivial since  $g$  depends on the solution itself). However, it is not necessary to perform such extension when dealing with (1.6).

For the problem (1.4) in Bogoya and Gómez (2012) it is proved existence and uniqueness of solutions for  $u_0 \in L^1(\Omega)$ , that a comparison principle is satisfied and it is also studied the asymptotic behavior of the solutions as  $t \rightarrow \infty$ .

Our main goal in this paper is to study the behaviour of solutions to this nonlocal model when the involved kernels are rescaled appropriately. If we consider the new kernels

$$J(\xi) = \frac{1}{\epsilon^{N+2}} J\left(\frac{\xi}{\epsilon}\right), \tag{1.5}$$

$$G(\xi) = \frac{C_1}{\epsilon^N} J\left(\frac{\xi}{\epsilon}\right),$$

then we arrive to

$$w_t^\epsilon(x, t) = \frac{1}{\epsilon^{N+2}} \int_{\Omega} J\left(\frac{x-y}{\epsilon}\right) (w^\epsilon(y, t) - w^\epsilon(x, t)) dy + \frac{C_1}{\epsilon^N} \int_{\partial\Omega} J\left(\frac{x-y}{\epsilon}\right) g(y, t) dS_y, \tag{1.6}$$

$$w(x, 0) = u_0(x),$$

$$(x, t) \in \bar{\Omega} \times (0, T),$$

$$x \in \bar{\Omega},$$

We have the following existence and uniqueness result:

**Theorem 1.1.** For every  $u_0 \in L^1(\Omega)$  and every  $g \in L^\infty_{loc}([0, \infty); L^1(\partial\Omega))$  there exists a unique solution  $w^\epsilon \in C([0, \infty); L^1(\Omega))$  to problem (1.6).

The proof follow the same lines of Theorem 2.2 in Bogoya and Gómez (2012) and then we omit the details here. As we have mentioned, our main objective is to show that the solution of the non-homogeneous Neumann problem for the heat Eq. (1.2), can be approximated by solutions of (1.6) when the parameter  $\epsilon$  goes to zero. We have the following theorem:

**Theorem 1.2.** Let  $\Omega$  be a bounded  $C^{2+\alpha}$  domain,  $g \in C^{1+\alpha, 1+\frac{\alpha}{2}}(\bar{(\mathbb{R}^N \setminus \Omega)} \times [0, T])$ , let  $v$  the solution to (1.2) and assume that  $v \in C^{2+\alpha, \frac{\alpha}{2}}(\bar{\Omega} \times [0, T])$ , for some  $0 < \alpha < 1$ . Let  $w^\epsilon$  be the solution to (1.6). Then, for each  $t \in [0, T]$

$$w^\epsilon(x, t) \rightharpoonup v(x, t) \quad \text{as } \epsilon \rightarrow 0 \text{ * -weakly in } L^\infty(\Omega).$$

We remark that the obtained convergence is the weak-\* topology is the same that was obtained in Cortazar et al. (2008) for the problem (1.3).

## 2. Proof of Theorem 1.2. Weak-\* convergence in $L^\infty$

In this section we give the proof of Theorem 1.2. To this end we use a result proved in Cortazar et al. (2008) for the problem (1.3). More precisely, we will use the following theorem:

**Theorem 2.1.** Let  $\Omega$  be a bounded  $C^{2+\alpha}$  domain,  $g \in C^{1+\alpha, 1+\frac{\alpha}{2}}(\bar{(\mathbb{R}^N \setminus \Omega)} \times [0, T])$ ,  $v \in C^{2+\alpha, \frac{\alpha}{2}}(\bar{\Omega} \times [0, T])$  the solution to (1.2), for some  $0 < \alpha < 1$ . Let  $u^\epsilon$  be a solution to (1.3). Then, there is an adequate constant  $K$  such that, for each  $t \in [0, T]$ ,

$$u^\epsilon(x, t) \rightharpoonup v(x, t) \quad \text{* -weakly in } L^\infty(\Omega).$$

We fix  $K$  in such a way that the conclusion of Theorem 2.1 holds.

**Proof (Proof of Theorem 1.2).** With the aim to prove Theorem 1.2, we just want to obtain that  $w^\epsilon(x, t)$  the solution to (1.6) is close to  $u^\epsilon(x, t)$  the solution to (1.3). To this end we consider equation verified by the difference

$$u^\epsilon(x, t) - w^\epsilon(x, t) \tag{2.1}$$

that is,

$$(u^\epsilon(x, t) - w^\epsilon(x, t))_t = \frac{1}{\epsilon^{N+2}} \int_{\Omega} J\left(\frac{x-y}{\epsilon}\right) ((u^\epsilon(y, t) - w^\epsilon(y, t)) - (u^\epsilon(x, t) - w^\epsilon(x, t))) dy + A_\epsilon(x, t)$$

$$(u^\epsilon(x, 0) - w^\epsilon(x, 0)) = 0,$$

where, from the fact that  $w^\epsilon, u^\epsilon$  satisfies (1.6) and (1.3) respectively, we get that

$$A_\epsilon(x, t) = \frac{K}{\epsilon^{N+1}} \int_{\mathbb{R}^N \setminus \Omega} J\left(\frac{x-y}{\epsilon}\right) g(y) dy - \frac{C_1}{\epsilon^N} \int_{\partial\Omega} J\left(\frac{x-y}{\epsilon}\right) g(y) dS_y.$$

After integration in  $\Omega$  of (2.1), we consider  $A_\epsilon(x, t)$  and we decompose it as follows:

$$\int_{\Omega} A_{\epsilon}(x, t) = I_1 - I_2, \tag{2.2}$$

with

$$I_1 = \frac{K}{\epsilon^{N+1}} \int_{\Omega} \int_{\mathbb{R}^N \setminus \Omega} J\left(\frac{x-y}{\epsilon}\right) g(y) dy dx,$$

and

$$I_2 = \frac{C_1}{\epsilon^N} \int_{\Omega} \int_{\partial\Omega} J\left(\frac{x-y}{\epsilon}\right) g(y) dS_y dx.$$

Let  $\delta$  such that  $2\epsilon < \delta$ . We consider the following sets:

$$\Gamma_{\delta}^+ = \{\tilde{x} \in \Omega / d(\tilde{x}, \partial\Omega) \leq \delta\},$$

$$\Gamma_{\delta}^- = \{\tilde{x} \in (\mathbb{R}^N \setminus \Omega) / d(\tilde{x}, \partial\Omega) \leq \delta\},$$

and

$$\Gamma_{[-\delta, \delta]} = \{\tilde{x} \in \mathbb{R}^N / d(\tilde{x}, \partial\Omega) \leq \delta\} = \Gamma_{\delta}^+ \cup \Gamma_{\delta}^-.$$

Note that

$$I_1 = \frac{K}{\epsilon^{N+1}} \int_{\Gamma_{\epsilon}^+} \int_{\Gamma_{\epsilon}^-} J\left(\frac{x-y}{\epsilon}\right) g(y) dy dx.$$

When  $\tilde{y} \in \partial\Omega$  and  $s \in [-\delta, \delta]$ , we use the change of variables

$$y = \tilde{y} + n(\tilde{y})s$$

being  $n(\tilde{y})$  the normal vector at  $\tilde{y}$  to obtain

$$I_1 = \frac{1}{\epsilon^N} \int_{\Gamma_{\epsilon}^+} \left[ \frac{1}{\epsilon} \int_0^{\epsilon} \int_{\partial\Omega} KJ\left(\frac{x - (\tilde{y} + n(\tilde{y})s)}{\epsilon}\right) g(\tilde{y} + n(\tilde{y})s) |\bar{J}(\tilde{y}, s)| dS_y ds \right] dx, \tag{2.3}$$

where  $|\bar{J}(\tilde{y}, s)|$  is the Jacobian of the change of variable. In the same way we can transform  $I_2$  and obtain

$$I_2 = \frac{C_1}{\epsilon^N} \int_{\Gamma_{\epsilon}^+} \int_{\partial\Omega} J\left(\frac{x-\tilde{y}}{\epsilon}\right) g(\tilde{y}) dS_{\tilde{y}} dx. \tag{2.4}$$

Now we consider the extension of  $g$  given by

$$g(\tilde{y} + n(\tilde{y})s) = \mathbf{g}(\tilde{y})$$

and from (2.3) and (2.4) we get

$$I_1 - I_2 = \frac{1}{\epsilon^N} \int_{\Gamma_{\epsilon}^+} \left[ \int_{\partial\Omega} (I_3) d\tilde{y} \right] dx, \tag{2.5}$$

being

$$I_3 = \frac{1}{\epsilon} \int_0^{\epsilon} KJ\left(\frac{x-\tilde{y}}{\epsilon} - n(\tilde{y})\frac{s}{\epsilon}\right) g(\tilde{y} + n(\tilde{y})s) |\bar{J}(\tilde{y}, s)| ds - C_1 J\left(\frac{x-\tilde{y}}{\epsilon}\right) g(\tilde{y} + n(\tilde{y})s).$$

Using the change of variables

$$\tilde{s} = \frac{s}{\epsilon}$$

in the first integral we obtain

$$I_3 = \int_0^1 KJ\left(\frac{x-\tilde{y}}{\epsilon} - n(\tilde{y})\tilde{s}\right) g(\tilde{y} + n(\tilde{y})s) |\bar{J}(\tilde{y}, \epsilon\tilde{s})| d\tilde{s} - C_1 J\left(\frac{x-\tilde{y}}{\epsilon}\right) g(\tilde{y} + n(\tilde{y})s). \tag{2.6}$$

Taking into account (2.2), (2.5) and (2.6), making the change of variables

$$z = \frac{x-\tilde{y}}{\epsilon}$$

and considering that

$$x = \tilde{y} + \epsilon z$$

we obtain

$$\int_{\Omega} A_{\epsilon}(x, t) = \int_{\partial\Omega} \left[ \int_{\{z / (\tilde{y} + \epsilon z) \in \Omega\}} \left( \int_0^1 KJ(z - n(\tilde{y})\tilde{s}) |\bar{J}(\tilde{y}, \epsilon\tilde{s})| d\tilde{s} - C_1 J(z) \right) dz \right] g(\tilde{y}) dS_y. \tag{2.7}$$

Choosing  $C_1$  appropriately in such a way that

$$\lim_{\epsilon \rightarrow 0} \int_{\{z / (\tilde{y} + \epsilon z) \in \Omega\}} \left( \int_0^1 KJ(z - n(\tilde{y})\tilde{s}) |\bar{J}(\tilde{y}, \epsilon\tilde{s})| d\tilde{s} - C_1 J(z) \right) dz = 0,$$

we conclude

$$\lim_{\epsilon \rightarrow 0} \int_{\Omega} A_{\epsilon}(x, t) = 0. \tag{2.8}$$

With this estimate we can control

$$u^{\epsilon}(x, t) - w^{\epsilon}(x, t).$$

In fact, let  $z_{\epsilon}$  be a nonnegative solution of the problem

$$z_{\epsilon}^t(x, t) = \frac{1}{\epsilon^{N+2}} \int_{\Omega} J\left(\frac{x-y}{\epsilon}\right) (z^{\epsilon}(y, t) - z^{\epsilon}(x, t)) dy + |A_{\epsilon}(x, t)| \quad (x, t) \in \bar{\Omega} \times (0, T), \\ z(x, 0) = 0, \quad x \in \bar{\Omega}. \tag{2.9}$$

Integrating (2.9) in  $\Omega$  and using the symmetry of  $J$  we have

$$\frac{\partial}{\partial t} \int_{\Omega} z^{\epsilon}(x, t) dx = \int_{\Omega} |A_{\epsilon}(x, t)| dx$$

and therefore

$$\int_{\Omega} z^{\epsilon}(x, t) dx = \int_0^t \int_{\Omega} |A_{\epsilon}(x, t)| dx dt.$$

From (2.8) we obtain that

$$\int_{\Omega} z^{\epsilon}(x, t) dx \rightarrow 0 \quad \text{when } \epsilon \rightarrow 0. \tag{2.10}$$

Now, since

$$A_{\epsilon}(x, t) \leq |A_{\epsilon}(x, t)|,$$

we get that

$$|u^{\epsilon}(x, t) - w^{\epsilon}(x, t)| \leq z_{\epsilon}(x, t).$$

Hence, from (2.10), it holds that

$$\int_{\Omega} |u^{\epsilon}(x, t) - w^{\epsilon}(x, t)| dx \rightarrow 0, \quad \text{when } \epsilon \rightarrow 0. \tag{2.11}$$

To finish the proof of the theorem, let  $\theta \in L^{\infty}(\Omega)$ , then

$$\int_{\Omega} w^{\epsilon} \theta dx \leq \int_{\Omega} |w^{\epsilon}(x, t) - u^{\epsilon}(x, t)| |\theta(x)| dx + \int_{\Omega} u^{\epsilon}(x, t) \theta(x) dx.$$

Taking into account (2.11) and Theorem 2.1 we conclude that

$$\int_{\Omega} w^{\epsilon} \theta dx \rightarrow \int_{\Omega} v(x, t) \theta dx, \quad \text{when } \epsilon \rightarrow 0,$$

as we wanted to show.  $\square$

Remark that we obtained  $\|u^{\epsilon} - w^{\epsilon}\|_{L^{\infty}(0, T); L^1(\Omega)} \rightarrow 0$  in (2.11). This says that solutions to the nonlocal models (1.3) and (1.6) are close for  $\epsilon$  small in a topology that is stronger than weak- $*$  in  $L^{\infty}$ .

### References

Abu Arqub, O., El-Ajou, A., Momani, S., 2015. Constructing and predicting solitary pattern solutions for nonlinear time-fractional dispersive partial differential equations. *J. Comp. Phys.* 293, 385–399.

- El-Ajou, A., Abu Arqub, O., Momani, S., Baleanu, D., Alsaedi, A., 2015a. A novel expansion iterative method for solving linear partial differential equations of fractional order. *Appl. Math. Comp.* 257, 119–133.
- El-Ajou, A., Abu Arqub, O., Momani, S., 2015b. Approximate analytical solution of the nonlinear fractional KdV-Burgers equation: A new iterative algorithm. *J. Comp. Phys.* 293, 81–95.
- Alberti, G., Bellettini, G., 1998. A nonlocal anisotropic model for phase transition: asymptotic behaviour of rescaled. *European J. Appl. Math.* 9, 261–284.
- Andreu-Vailló, F., Mazón, J.M., Rossi, J.D., Toledo-Melero, J.J., 2010. Nonlocal diffusion problems. *Am. Math. Soc. Math. Surveys Monographs*, 165.
- Andreu, F., Mazón, J.M., Rossi, J.D., Toledo, J., 2008. The Neumann problem for nonlocal nonlinear diffusion equations. *J. Evol. Eq.* 8 (1), 189–215.
- Bates, P., Fife, P., Ren, X., Wang, X., 1997. Travelling waves in a convolution model for phase transitions. *Arch. Ration. Mech. Anal.* 138, 105–136.
- Bodnar, M., Velázquez, J.J.L., 2006. An integro-differential equation arising as a limit of individual cell-based models. *J. Diff. Eq.* 222, 341–380.
- Bogoya, M., Gómez, J., 2012. On a nonlocal diffusion model with Neumann boundary conditions. *Nonlinear Anal.* 75, 3198–3209.
- Caffarelli, L., Silvestre, L., 2009. Regularity theory for fully nonlinear integro-differential equations. *Comm. Pure Appl. Math.* 62, 597–638.
- Carrillo, C., Fife, P., 2005. Spatial effects in discrete generation population models. *J. Math. Biol.* 50 (2), 161–188.
- Chasseigne, E., Chaves, M., Rossi, J.D., 2006. Asymptotic behaviour for nonlocal diffusion equations. *J. Math. Pures Appl.* 86 (9), 271–291.
- Cortazar, C., Elgueta, M., Rossi, J.D., Wolanski, N., 2008. How to approximate the heat equation with Neumann boundary conditions by nonlocal diffusion problems. *Arch. Rat. Mech. Anal.* 187 (1), 137–156.
- Cortazar, C., Elgueta, M., Rossi, J.D., Wolanski, N., 2007. Boundary fluxes for nonlocal diffusion. *J. Diff. Eq.* 234, 360–390.
- Coville, J., Dávila, J., Martínez, S., 2008. Nonlocal anisotropic dispersal with monostable nonlinearity. *J. Diff. Eq.* 244, 3080–3118.
- Coville, J., Dupaigne, L., 2007. On a nonlocal equation arising in population dynamics. *Proc. Roy. Soc. Edinburgh Sect. A* 137, 1–29.
- Fife, P., 2003. Some nonclassical trends in parabolic and parabolic-like evolutions. *Trends in nonlinear analysis*. Springer, Berlin, pp. 153–191.
- Fournier, N., Laurencot, P., 2006. Well-posedness of Smoluchowski's coagulation equation for a class of homogeneous kernels. *J. Funct. Anal.* 233, 351–379.
- Gilboa, G., Osher, S., 2006. Nonlocal linear image regularization and supervised segmentation. *UCLA CAM Report* 06–47.
- Ignat, L.I., Rossi, J.D., 2007. A nonlocal convection-diffusion equation. *J. Funct. Anal.* 251 (2), 399–437.