

Supplementary table S4. Comparison of antimicrobial activity 5'-hydroxy-aurapten with other coumarin derivatives studied in the literature

Coumarin derivative	Microorganism	Antimicrobial activity	Refs
7-(5-Hydroxy-3,7-dimethylocta-2,6-dienyloxy)-chromen-2-one	<i>Aspergillus flavus</i>	Antifungal activity	This work
4-((5-Mercapto-4-phenyl-4H1,2,4-triazol-3-yl)-methoxy)-2H-chromen-2-one	<i>Aspergillus niger and Candida albicans</i>	Antifungal activity	(Al-Amiry et al. 2012)
4-((5-(Phenylamino)-1,3,4-thiadiazol-2-yl)methoxy)-2Hchromen-2-one	<i>Aspergillus niger and Candida albicans</i>	Antifungal activity	(Al-Amiry et al. 2012)
3-Acetyl-8-amino-4,7-dihydroxy-chromen-2-one	<i>Staphylococcus aureus, E. coli, and Bacillus cereus</i>	Antibacterial activity	(Behrami and Krasniqi 2012)
Cephalosporin	<i>Staphylococcus aureus, Escherichia coli, and Pseudomonas aeruginosa</i>	Antibacterial activity	(Bonsignore et al. 1998)
4-Methyl-3-phenyl-6-[4-(3-aryl-1-phenyl-1H-pyrazol-4-yl)-6-aryl-pyridin-2-yl]	<i>Escherichia coli, Bacillus subtilis</i>	Antibacterial activity	(Brahmbhatt et al. 2010)
4-Chloro-3-((substituted phenylamino) methyl)-2Hchromen-2-one	<i>Staphylococcus aureus, Bacillus subtilis, Escherichia coli, Aspergillus niger, and Candida albicans</i>	Antibacterial and antifungal activity	(Bairagi et al. 2009)
4-Aryloxymethylcoumarins	<i>Staphylococcus aureus, Streptococcus faecalis, Escherichia coli, Pseudomonas aeruginosa, Klebsiella pneumonia, Aspergillus favus, Aspergillus fumigates, Candida albicans, Penicillium notatum, Rhizopus</i>	Antibacterial and antifungal activity	(Barzon et al. 2000)
6',7'-Dihydroxybergamottin	<i>S. aureus, S. epidermidis, P. aeruginosa, E. cloacae, K. pneumonia, and E. coli</i>	Antibacterial activity	(Widelski et al. 2018)
4-Acetatecoumarin	<i>A. fumigates, A. favus</i>	Antifungal activity	(Guerra Felipe QS et al. 2018)
7-Hydroxy-6-nitro-2H-1-benzopyran-2-one	<i>A. fumigates, A. favus</i>	Antifungal activity	(Guerra Felipe Queiroga Sarmento et al. 2015)
7-(5-Hydroxy-3,7-dimethylocta-2,6-dienyloxy)-chromen-2-one	<i>Aspergillus flavus</i>	Antifungal activity	This work

References

- Al-Amiry AA, Kadhum AAH, Mohamad AB. 2012. Antifungal activities of new coumarins. *Molecules* 17:5713-5723.
- Bairagi S, Bhosale A, Deodhar MN. 2009. Design, synthesis and evaluation of Schiff's bases of 4-chloro-3-coumarin aldehyde as antimicrobial agents. *E-journal of Chemistry* 6.
- Barzon L, Bonaguro R, Palu G, Boscaro M. 2000. New perspectives for gene therapy in endocrinology. *Eur.J Endocrinol.* 143:447-466.
- Behrami A, Krasniqi I. 2012. Antibacterial activity of coumarine derivatives synthesized from 8-amino-4,7-dihydroxy-chromen-2-one and comparison with standard drug. *Journal of Chemical and Pharmaceutical Research* 4:2495-2500.
- Bonsignore L, Cottiglia F, Elkhaili H, Jehl F, Lavagna SM, Loy G, Manna F, Monteil H, Pompei D, Secci D. 1998. Synthesis and antimicrobial activity of coumarin 7-substituted cephalosporins and sulfones. II *Farmaco* 53:425-430.
- Brahmbhatt DI, Kaneria A, Patel AK, Patel NH. 2010. Synthesis and antimicrobial screening of some 3-[4-(3-aryl- 1-phenyl-IH-pyrazol- 4-yl)-6-aryl-pyridin-2-yl] and 4-methyl- 3-phenyl-6-[4-(3-aryl- 1-phenyl- IH-pyrazol- 4-yl)-6-aryl-pyridin-2-yl] coumarins. *Indian Journal of Chemistry - Section B Organic and Medicinal Chemistry* 49:971-977.
- Guerra FQ, Araújo RS, Sousa JP, Silva VA, Pereira FO, Mendonça-Junior FJ, Barbosa-Filho JM, Pereira JA, Lima EO. 2018. A new coumarin derivative, 4-acetatecoumarin, with antifungal activity and association study against *Aspergillus* spp. *brazilian journal of microbiology* 49:407-413.
- Guerra FQS, Araújo RSA, Sousa JPd, Pereira FdO, Mendonça-Junior FJ, Barbosa-Filho JM, de Oliveira Lima E. 2015. Evaluation of antifungal activity and mode of action of new coumarin derivative, 7-hydroxy-6-nitro-2h-1-benzopyran-2-one, against *Aspergillus* spp. *Evidence-Based Complementary and Alternative Medicine* 2015.
- Widelski J, Luca SV, Skiba A, Chinou I, Marcourt L, Wolfender J-L, Skalicka-Wozniak K. 2018. Isolation and antimicrobial activity of coumarin derivatives from fruits of *Peucedanum luxurians* Tamamsch. *Molecules* 23:1222.