

Dr. Nadiyah_JKSUS-D-22- 00151_R4

by Mohammed Mohammed

Submission date: 21-Apr-2022 03:04AM (UTC-0500)

Submission ID: 1350721294

File name: JKSUS-D-22-00151_R4.pdf (2.83M)

Word count: 6443

Character count: 35236

The synthesis of polysaccharide crude nanoparticles extracts from Taif rose petals and its effect on eggplant seedlings under drought and salt stress

--Manuscript Draft--

Manuscript Number:	JKSUS-D-22-00151R4
Article Type:	Full length article
Section/Category:	Environmental Sciences
Keywords:	Taif rose petals, Polysaccharides, Antioxidant enzymes, environmental factors, lipid peroxidation, total soluble sugar
Corresponding Author:	Nadiyah Alabdallah, PhD Imam Abdulrahman Bin Faisal University Dammam, SAUDI ARABIA
First Author:	Nadiyah Alabdallah, PhD
Order of Authors:	Nadiyah Alabdallah, PhD Saleh M Alluqmani
Abstract:	<p>Objectives: Natural biopolymer nanoparticles are an emerging area of research in green electronic devices and biomedical and agricultural fields. However, because drought and salt are two keys environmental aspects that affect agricultural crop yields around the world, a new strategy has been suggested to valorize the waste of Taif rose petals (<i>Rosa damascena</i> Miller var. <i>trigintipetala</i>) in the production of polysaccharide material on the nanoscale. In this study, we explore the effects of drought and salinity stress on the physiology of eggplants.</p> <p>Methods: The key objectives of this research were to investigate the effect of polysaccharide crude nanoparticles extract (PCE) in the alleviation of drought and salinity stress in eggplants.</p> <p>Results: Reduced chlorophyll content, chlorophyll fluorescence (Fv/Fm), and chlorophyll stability index (CSI) were seen in the plants that had been subjected to drought and salt stress. Under drought and salt stress, proline, total soluble protein, and sugar levels were all shown to be higher. However, polysaccharide crude nanoparticles extract reverses the negative effects of drought and salt stress and increases the chlorophyll content, chlorophyll fluorescence (Fv/Fm), and chlorophyll stability index (CSI) in the presence of drought and salt stress. When PCE supplemented, the accumulation of lipid peroxidation was reduced, and the activities of antioxidant enzymes (SOD, CAT, POX, GR) were raised.</p> <p>Conclusion: In summary, the results suggest that exogenous PCE can mitigate the detrimental effects of drought and salt stress in eggplants by modulating antioxidant systems. Furthermore, the findings of this study give experimental proof that PCE can be used to increase drought tolerance as well as production in eggplant plants.</p>
Suggested Reviewers:	<p>Sam Paul spaul@ju.edu.sn</p> <p>Vittal Malhotra vital.mal1@yahoo.com</p> <p>Mohammed Ayub ayub.mohammad@gmail.com</p>
Opposed Reviewers:	
Response to Reviewers:	Dear Editor, The Plagiarism of the revised manuscript JKSUS-D-22-00151_R4 was found to be 17%. I am attaching the plagiarism report for your reference. Additionally, we have followed the author guidelines mentioned in your email. Please do the needful.

To,
The Editor,

Journal of King Saud University Science

Subject: Resubmission of Review Article

Dear Editor,

I along with my co-author would like to resubmit the original Review article entitled “**The synthesis of polysaccharide crude nanoparticles extracts from Taif rose petals and its effect on eggplant seedlings under drought and salt stress**” (JKSUS-D-22-00151_R4) in Journal of King Saud University Science. We both the authors are thankful to the reviewers as well as editors for your valuable comments. We both the authors have justified the raised quires and resubmitting the manuscript. The Plagiarism was found to be less than 20%. References was arranged with EndNote software. Once again, we confirm this manuscript was not submitted anywhere else apart from your esteem journal. I as a corresponding author assure you to the publication fee once the manuscript gets accepted. Please acknowledge us.

Dr. Nadiyah Alabdallah

Corresponding author

The synthesis of polysaccharide crude nanoparticles extracts from Taif rose petals and its effect on eggplant seedlings under drought and salt stress

Nadiyah M Alabdallah^a, Saleh M. Alluqmani^b

^aDepartment of Biology, College of Science, Imam Abdulrahman Bin Faisal University, P.O. Box 1982, 31441, Dammam, Saudi Arabia; nmalabdallah@iau.edu.sa

^bDepartment of Physics, Faculty of Applied Science, Umm Al-Qura University, Makkah P.O. Box 21955, Saudi Arabia; smluqmani@uqu.edu.sa

Correspondence: nmalabdallah@iau.edu.sa

Abstract

Objectives: Natural biopolymer nanoparticles are an emerging area of research in green electronic devices and biomedical and agricultural fields. However, because drought and salt are two keys environmental aspects that affect agricultural crop yields around the world, a new strategy has been suggested to valorize the waste of Taif rose petals (*Rosa damascena* Miller var. *trigintipetala*) in the production of polysaccharide material on the nanoscale. In this study, we explore the effects of drought and salinity stress on the physiology of eggplants.

Methods: The key objectives of this research were to investigate the effect of polysaccharide crude nanoparticles extract (PCE) in the alleviation of drought and salinity stress in eggplants.

Results: Reduced chlorophyll content, chlorophyll fluorescence (F_v/F_m), and chlorophyll stability index (CSI) were seen in the plants that had been subjected to drought and salt stress. Under drought and salt stress, proline, total soluble protein, and sugar levels were all shown to be higher. However, polysaccharide crude nanoparticles extract reverses the negative effects of drought and salt stress and increases the chlorophyll content, chlorophyll fluorescence (F_v/F_m), and chlorophyll stability index (CSI) in the presence of drought and salt stress. When PCE was supplemented, the accumulation of lipid peroxidation was reduced, and the activities of antioxidant enzymes (SOD, CAT, POX, GR) were raised.

Conclusion: In summary, the results suggest that exogenous PCE can mitigate the detrimental effects of drought and salt stress in eggplants by modulating antioxidant systems. Furthermore, the findings of this study give experimental proof that PCE can be used to increase drought tolerance as well as production in eggplant plants.

Key words: Taif rose petals, Polysaccharides, Antioxidant enzymes, environmental factors, lipid peroxidation, total soluble sugar

Introduction

Natural polysaccharides are biomaterials and have received considerable interest due to their significant structural features, physical and chemical properties, and biological activities [1]. These biomaterials have the promising potential to be applied for green electronic devices, biomedical applications and agriculture [2, 3]. Polysaccharides can be found in plants, algae and animals. Among these natural sources of polysaccharides, floral waste consumed in social occasions and events were not fully examined as value-added product [4]. Studies on waste *Rosa damascena* petals reported they can serve as a source of polysaccharides material [5]. However, the functionality of extracted polysaccharide can be affected by their dimensionality, shape and purity during the optimal synthetic procedure, leading to an increase in the range of potential applications [6]. As a sustainable strategy could be suggested and targeted the application of polysaccharides in the protection of crop yields could be essential and gaining a great deal of attention.

³⁸ Eggplant (*Solanum melongena* L.) is a member of the Solanaceae family and one of the most common agricultural products, ranking third in terms of production after potato and tomato [7]. Eggplant fruits are known for their low-calorie content and it also contains carbohydrates, vitamins, and high protein content [8]. In addition, it possesses antioxidant, ant-proliferative, and anti-inflammatory properties [9]. However, in their full lifecycle, eggplants are extremely vulnerable to several abiotic stresses and biological stimuli that can restrict their growth. They are quite susceptible to water shortages, as they require a large volume of water for continued growth [10].

When water shortages rose from 20% to 40% of the field capacity, eggplant fruit yield declined by up to 60% [11]. Eggplant is also relatively susceptible to the effects of salinity [12]. The optimal electrical conductivity (EC) for eggplant growing is between 1.1 and 1.5 dSm⁻¹ [13]. Because it is such a valuable plant, understanding its sensitivity and tolerance to increased salinity and drought stress is crucial in order to build effective crop production techniques. Since eggplant is one of the most major crops in the Gulf, and changes in climate will increase the intensity and frequency of coupled pressures (drought and salt), addressing the problem of climate-resilient crops is essential.

Drought is the key environmental variables that that has a detrimental impact on crop yields around the world [14, 15]. Furthermore, rapid evaporation causes water loss, which exacerbates salt accumulation. Non-optimal irrigation practices can also cause soil salinity, resulting in agricultural production loss [6]. Both stress and anxiety generate physiological and ⁹ biochemical changes, such as the creation of reactive oxygen species (ROS) as a result of metabolic processes [16]. The production of suitable solutes, antioxidant components, and the induction of secondary protection against oxidative stress by crops are all crucial to mitigate cellular damage caused by ROS accumulation in the environment. It has been demonstrated that drought and salinity stress cause a variety of interacting processes, such as the suppression of enzyme activity in metabolic pathways. Plant responses to environmental stressors such as drought and salinity, on the other hand, have received little attention. When exposed to abiotic stress, many plant species have been reported to accumulate suitable solutes such as proline [17, 18]. They have been shown to have an important function in crop osmoregulation [19, 20].

⁴⁴ It has been demonstrated in previous studies that exogenous application of many plant hormones can ³ improve the abiotic stress tolerance of plants [19, 21, 22]. However, exogenous application of crude polysaccharides nanoparticles extracts from Taif rose petals (*Rosa damascena* Miller var. *trigintipetala*) (has never been reported) was not investigated. A polysaccharide is a carbohydrate macromolecule consisting of monosaccharide units that are covalently attached together by glycoside bonds, whether in a linear or branched setup. They ⁴² are essential for the survival and activity of all organisms, and play an important role in their metabolic processes [23]. The method used to extract polysaccharides from plants has a considerable impact on not only the extraction rate, but also the characteristics and bioactivities of the polysaccharides that have been extracted, all of which have an impact on the applications of the polysaccharides [24]. Despite the fact that numerous biochemical and physiological changes have been shown to be involved in the adaptation process of several crops to environmental challenges, however little understood about the sensitivities of eggplants to ROS production, which leads to oxidative stress. In the current research, we address the effects on plant development, antioxidant defense mechanisms, oxidative damage, and physiological and biochemical changes that occur ⁵ as a result of drought and salinity stress ⁵ in order to better understand the plant's adaptation to these stresses.

Materials and Methods

2.1 Extraction of crude polysaccharides nanoparticles

Taif roseus were purchased from Al Qurashi Rose Factory, Al Hada, Taif, Saudi Arabia. Taif roseus polysaccharides nanoparticles were extracted. A 5g of plant components were extracted in a ¹³ 30% (w/v) potassium hydroxide solution at 90°C for 120 minutes. The plant parts homogenate was combined with 100 mL of a 30% (w/v)

potassium hydroxide (KOH) solution, which was boiled in water at 100°C for 1 hour before being chilled and centrifuged. The polysaccharide concentration of the upper solution was tested at neutral pH. The residue was sifted using Whatman filter paper. The filtrate was condensed using a rotary extractor at 60°C and reduced pressure. The concentration was tripled by adding three times the amount of 98% ethanol. After that, the solution was left for 7 days to allow the polysaccharides to precipitate. The size of PCE crude nanoparticles was determined using TEM (2.33 to 4.36) and Fourier-transform infrared (FTIR) spectroscopy were used to confirm the isolation of polysaccharides from a particular plant. The FTIR samples were acquired in the band of 5000–400 cm⁻¹ utilizing Fourier transmission infrared spectrometer (Perkin Elmer).

2.2. Separation and quantification

An Agilent liquid chromatographic system (USA) with the column SB-C18 and G 4226A software was used to separate and quantify gallic acid, quercetin, and tannic acid in plant materials (1.8 m, 4.6 150 mm). Finally, Taif rose polysaccharide extracts (0, 15, 35, and 65mM) were obtained.

2.3. Chromatographic analysis of glucose and fructose

To identify glucose in the sample, 0.6% aqueous acetic acid solution (A) and methanol (B) (10: 90) (V/V) were used as the mobile phase. The samples were eluted using the gradient given below, with a flow rate of 0.700 mL/min and an injection volume of 11. The temperature in the column was kept constant at 25°C. A chromatogram was acquired at a wavelength of 274 nm based on the absorption maxima of the tested materials (Figure 2). The presence of glucose in the sample was determined by measuring its retention time and spiking it with glucose as a reference under the same conditions.

Acetonitrile solution (A) and methanol (B) were utilized as the mobile phase for identifying fructose sugar in the sample (70: 30). The following gradient was used to elute the samples: The injection volume was 10 l and the flow rate was 1000 mL/min. The column temperature was kept constant at 25° C. According to the absorption maxima of the examined samples, a chromatogram was acquired at a wavelength of 274 nm (Figure 7). Retention time measurements and D-fructose spikes were used to assess the presence of fructose in the sample.

2.4. Plant growth, treatments, and experimental design

The two pot trials were carried out at Imam Abdulrahman Bin Faisal University and Umm Al-Qura University in Saudi Arabia, in their laboratories and green houses. Eggplant seeds were purchased in Altuajri, Saudi Arabia, and sterilized in 0.35% sodium hypochlorite for ten minutes before being properly cleaned with distilled water. To germinate the seeds, three seeds were placed in each of the 20 cm plastic pots, which contained a one-to-one volume ratio of potting soil and compost. To determine each pot's field capacity, FC (100 %) had to soak the soil with distilled water. Before being reused, all of the pots were covered with plastic sheets and allowed to drain for 48 hours. After 48 hours of draining and 24 hours of drying at 105° C, the pot weights were recalculated. Field capacity was determined by comparing soil weight before draining and after drying to determine the soil moisture content. Drought stress was induced (50% FC, 35% FC) by allowing plants to dry out over a 15-day period by withholding water. Field capacity (FC) was maintained by weighing the pot. In order to impose salinity stress, NaCl at final concentrations of 0, 20, 50, 80, and 120 mM (at intervals of 3 days between each treatment) was supplied for a total of 15 days. Plants

were maintained on a 25/19^o C day/night temperature cycle with a 16/8 h light/dark cycle. At least three plants from each treatment were employed in each of the three biological replicates to assess growth and physiological parameters. The plants were harvested at 35 days.

2.5. Chlorophyll content determination

Chlorophyll content was extracted based on Aron method [25]. A minimum of 0.25g of leaf samples were taken and grounded with 5ml of acetone at room temperature, centrifugation was performed for 10 minutes at 3000rpm at 400^o C. The supernatant absorbs at 663 and 645nm was used to measure the amounts of chlorophyll a and b

2.6. Chlorophyll fluorescence (Fv/Fm) and Chlorophyll stability index measurements (CSI)

At midday, chlorophyll fluorescence (Fv/Fm) in eggplant seedlings was measured using a chlorophyll fluorometer (OS-30p, Opti-Science, Inc., Hudson, USA). The chlorophyll stability index (CSI) was calculated using the Murphy method (1962).

The CSI was calculated using the following formula:

$$CSI (\%) = 1 - \frac{\text{total chlorophyll of heated samples}}{\text{Total chlorophyll of non - heated sample}} \times 100$$

2.7. Lipid peroxidation

Using the thiobarbituric acid (TBA) method devised by Heath and Packer [26], it was determined that the products of lipid peroxidation (which were indicative of MDA concentration) were present. The findings for these assays were obtained by combining 0.5% TBA with 1 ml of trichloroacetic acid (TCA) extract (prepared in 20% TCA). The mixture was incubated for 30 minutes at 95° C before being chilled in ice for 10 minutes. To deliver the best results result, the specific absorbance of the objects was measured at 532 nm, and the nonspecific background absorbance at 600 nm was subtracted from the values. The MDA content was calculated using the molar extinction coefficient of 155 mM⁻¹ cm⁻¹.

2.8. Proline content

Bates et al. [27] used a modified technique to determine the amount of proline in the sample. It was necessary to homogenize fresh leaf samples (0.1 g, for example) from each treatment in the mortar and pestle in order to obtain the homogenous solution. An equal amount of each homogenate was mixed with an equal amount of glacial acetic acid, which was then mixed with an equal amount of ninhydrin. Each homogenate was present in roughly 0.2 ml of the final mixture. For 30 minutes, the reaction mixture was heated to a boil in a 1000° C water bath before being quickly cooled in an ice bath. Toluene (400μl) was added to the reaction mixture after the mixture had cooled. Chromophore-containing toluene was fully mixed before a spectrophotometer equipped with a UV–visible detector (Chemito Spectrascan, UV 2600) was used to measure the absorbance of red color created against a toluene blank.

2.9. Total soluble sugar (TSS)

Cock et al. [28] discovered a method for measuring the amount of soluble sugars available in eggplant leaves. The total soluble sugar analysis was carried out using 0.5 mL of soluble sugar extract and 4.5 mL of 85% ethanol in a test tube. Slowly add 10 mL of anthrone reagent to the sample tubes in a cold-water bath until they are completely filled. The tubes were immersed in a boiling water bath for 7.5 minutes before being swiftly cooled in an ice water bath. The absorbance at 630 nm was measured one hour after cooling.

2.10. Total soluble protein (TSP)

500 mg homogenized mixture was centrifuged for 25 minutes at 16000 x g and 4°C in 5 ml potassium phosphate buffer (pH 7.0, 10 mM, 4 percent polyvinylpyrrolidone, PVP). The absorbance at 595 nm was measured after adding 980 ml of Bradford solution to the supernatant (20 ml). Finally, to calculate the TSP content, a standard curve was constructed using various amounts of bovine serum albumin [29].

2.11. Antioxidant enzyme activity

The extraction of antioxidant enzymes was proceeded as per the previous studies [30]. The leaves were extracted with 10 mL of phosphate buffer and 0.5 g of freshly chosen material (pH 7). Afterwards, the homogenate was centrifuged for 10 minutes at 15,000x g at 40°C. Antioxidant enzyme activity was measured by heating the supernatant to 200°C.

2.11.1. Superoxide dismutase (SOD) activity determination

SOD activity was measured using the nitro-blue-tetrazolium (NBT) reduction method [31]. The reaction mixture contains roughly 3 ml of 50 ml enzyme extract, 150 ml

riboflavin (13 mM), 2.5 ml methionine (13 mM), 250 ml NBT (63 mM), and 50 ml phosphate buffer (50 mM, pH 7.8). The absorbance at 560 nm was measured using a spectrophotometer.

2.11.2 Catalase (CAT) activity determination

The Aebi [32] protocol was used to measure CAT activity. In a 40 mL container, the enzyme extract was combined with 0.016 mL H₂O₂ (30%) and a 10 mM phosphate buffer solution (pH 7). Finally, using a spectrophotometer, the absorbance at 240 nm was determined (LKB-Biochrom 4050).

2.11.3 Peroxidase (POX) activity determination

Hammerschmidt et al [33] established methods for measuring the activity of the peroxidase (POX) enzyme. In the solution, there was 0.25% (v/v) guaiacol and 100 mM H₂O₂ in a buffer solution of 100mM sodium phosphate (pH 6.0). For three minutes, 100 liters of crude enzyme extract were combined, and every 30 seconds, the absorbance at 470 nm was measured. g⁻¹ FW was used to quantify POX activity.

2.11.4 Measuring glutathione reductase (GR) activity

Glutathione reductase (GR) activity was assessed using the Foyer and Halliwell procedures in a mixture of sodium phosphate buffer (pH 7.8), NAD phosphate (0.01 mM), and 0.1 mM glutathione reductase in an assay mixture [34].

2.12 Statistical evaluation

The data were analyzed using analysis of variance (ANOVA) in the MINITAB 17 statistical tool, with the treatment mean±standard error (n=3) provided [35]. At the

p<0.05 level, the LSD test indicates that bars with dissimilar letters are statistically significantly different.

Results

PCE nanoparticle was derived from the Taif rose. TEM analysis was employed to explore the structure and distribution of polysaccharide nanoparticles. Figure 1 shows the TEM image and we clearly observe the spherical shape of polysaccharide nanoparticle with a size on the nanoscale. The FTIR spectra of a basic polysaccharide are shown in Figure 2. Furthermore, we have investigated the chromatographic technique to analyze the glucose and fructose (Figure 3-4).

We examined at a variety of physiological and biochemical parameters to see if PCE had any favorable impacts on eggplant seedlings. When eggplants were exposed to drought stress (50% and 35% FC) and varied salinity concentrations (20, 50, 80, and 120 mM), the levels of Chla, Chlb, carotenoids, total chl, and chlorophyll fluorescence (Fv/Fm) were drastically reduced (p<0.001; Figure 5). However, when eggplants were subjected to drought and salinity stress, PCE treatment increased these photosynthetic metrics dramatically.

Drought treatments (50% and 35% FC) resulted in substantial (p<0.001) decreases in CSI in eggplant seedlings of 42% and 71%, respectively, as compared to control seedlings. Externally administered PCE (15, 35, and 65mM) significantly increased CSI by 36%, 49%, 64%, and 33%, 47%, and 72%, respectively, as compared to drought stressed (50% and 35% FC) drought plants alone (Figure 3). In contrast, salt treatments of 20, 50, 80, and 120 mM reduced CSI in eggplants by 17%, 33%, 61%, and 84%, respectively, as compared to control seedlings that did not receive PCE (Figure 3).

Despite this, the exogenous infusion of PCE considerably (p<0.001) boosted the CSI.

Drought stress (50% and 35% FC) significantly ($p < 0.001$) elevated MDA in eggplant seedlings by 13% and 46%, respectively, as compared to controls. Nonetheless, pretreatment with PCE (15%, 35%, and 65%) reduced MDA at 50% FC by 14%, 48%, and 70%, and at 35% FC by 13%, 41%, and 60%, respectively, compared to drought stressed drought plants alone (Figure 6). The salt treatments (20, 50, 80, and 120 mM) raised MDA content in eggplant seedlings by 42%, 55%, 58%, and 63%, respectively.

However, ⁶ the exogenous application of PCE decreased MDA content significantly ($p \leq 0.001$), in comparison with those in the salt-stressed eggplant (Figure 7).

Drought treatments (50% and 35% ⁶ FC) significantly ($p < 0.001$) raised proline content by 44% and 27%, respectively, and TSP by 37% and 26% which in turn strikingly increased the TSS by 23% and 23% relative to the control (Figure 6-7). However, pretreatment of PCE (15, 35, 65mM) decreased proline content by 10%, 15%, 22% and 7%, 14%, 31%, TSP by 14%, 48%, 70% and 13%, 41%, 60%, TSS by 16%, 27%, 43% and 4%, 11%, 14%, when eggplant seedlings exposed to 50% and 35% FC. In response to salt stress (20, 50, 80, 120 mM), the content of proline, TSP, and TSS increased by 4%, 12%, 30%, 28%, 42%, 55%, 58%, 63%, and 4%, 39%, 49%, 51%, respectively. However, the external supply of PCE considerably ($p < 0.001$) reduced proline, TSP, and TSS content (Figure 7-8).

Drought treatments (50% and 35% FC) resulted in ⁶ significant ($p < 0.001$) increases in SOD activity of 52% and 56%, CAT activity of 29% and 47%, and POX activity of 43% and 55%, respectively, which increased GR activity by 30% and 40% relative to the control (Figure 9-10). However, PCE (15, 35, 65mM) increased SOD activity by 7%, 11%, 15%, and 17%, 20%, 25%, CAT activity by 15%, 32%, 42% and 12%, 22%,

36%, POX activity by 11%, 29%, 44% and 8%, 25%, 44%, whereas GR activity increased by 4%, 25%, 28% and 4%, 22%, 31%, respectively, ¹ in comparison to that ⁴¹ observed in the drought-stressed (50% and 35% FC) plants. SOD; CAT; POX; and GR activities ⁴¹ were increased in salt-stressed (20, 50, 80, 120 mM) eggplants by 15%, 42%, 47%, 48%; 4%, 27%, 32%, 39%; 8%, 19%, 31%, 38%; and 8%, 11%, 21%, 25%, respectively, compared to untreated plants. However, when eggplants were exposed to PCE treatments, their ⁵¹ SOD, CAT, POX, and GR activities increased when ⁵¹ compared to salt-stressed (20, 50, 80, and 120 mM) seedlings (Figure 9-10).

Discussion

¹⁹ This study examined the impact of drought and salinity stress on the physiology of eggplant. Chlorophyll is a vital component that allows plants to undergo photosynthetic activity. Photosynthesis-related features and chlorophyll pigments, which are the major factors of plant growth, often influence plant growth performance. In this ³ study, we discovered that when eggplant leaves were subjected to dryness and salt, they lost chlorophyll. Drought and salt stress increase the ⁹ generation of reactive oxygen species (ROS), which ²⁵ causes lipid peroxidation and, as a result, a decrease in plant chlorophyll content [36]. A decrease in chlorophyll content due to saltwater stress is common in drought and salt-sensitive plant species. This is due to dehydration and salt toxicity, which frequently results in burning of leaves or other fleshy parts, as well as the destruction of other pigments. PCE derived from the Taif rose, on the other hand, was found to be effective in the restoration of chlorophyll in eggplants. PCE may aid in the recovery of photosynthetic regulation by enhancing metabolites for eggplant growth, which in turn aids in the regeneration of chlorophyll. Furthermore, drought and salt stress reduced the chlorophyll stability index (CSI) as well as the maximal quantum yield of PSII (Fv/Fm) in the eggplant.

Chlorophyll fluorescence (Fv/Fm) is related to photosynthetic efficiency, and when plants are stressed, their photosynthetic efficiency decreases. As a result, chlorophyll fluorescence has become a key parameter for assessing plant chlorophyll content, classifying plants, and identifying abiotic stress in plants. Chlorophyll content, on the other hand, is directly related to chlorophyll fluorescence [21, 37].

Drought and salt stress cause a decrease in Fv/Fm value, indicating photoinhibition, indicating that the ²⁸ light-harvesting complex of PSII in stressed eggplant is disrupted. According to the outcomes of our study, PCE aids in the alleviation of photoinhibition in eggplants by increasing the Fv/Fm value. It is possible that the aforementioned traits are to blame for ³⁹ the significant increases in total soluble sugars and protein levels found in eggplants when treated ⁵ to drought and salt stress. Proline buildup in stressed plants has been demonstrated to be an important defense response for maintaining osmotic pressure within the cell. Proline accumulation potential under stress conditions has been linked ¹⁶ to stress tolerance in a range of plant species. More importantly, applying PCE to eggplants grown ⁴³ in drought and salt conditions has been shown to increase their tolerance to drought and salinity stress through biosynthetic modulation of osmolytes. According to the previous findings, the beneficial effect of osmolytes on stressed eggplants can be associated to the involvement of osmotic adjustment in membrane integrity and the prevention of physiological dehydration in plant cells. Under a wide range of stress conditions, a similar pattern in the accumulation of osmolytes was observed [38]. Despite the fact that proline concentration increased under both drought and salinity stress conditions, it was unable to offset the oxidative damage caused by the increase in H₂O₂ and the subsequent increase in lipid peroxidation. It has already been established that lipid peroxidation is connected with cellular injury induced by a variety of environmental

stressors. In this research, we found that stress treatments significantly enhance lipid peroxidation, and that this increase is closely linked to the generation of ROS, which promotes membrane instability. Nevertheless, MDA content decreased by exogenous application of PCE, which may be accomplished via limiting ROS generation.

SOD, CAT, POX, and GR antioxidant enzymes were all more effective in eggplants that had been treated to drought and salt stress. When there is a lot of drought and salt stress, these enzymes become more active, which helps them resist the detrimental effects of stress by getting rid of reactive oxygen species [22]. Our findings are comparable with those of other researchers who have studied a wide range of plant species under a variety of environmental stresses [39, 40]. This study found that exogenous PCE modifies ⁴ antioxidant enzyme activity and keeps reactive oxygen species levels tolerable, protecting eggplants from oxidative damage in a stress environment. The crucial involvement of PCE in ⁴ increasing the activities of SOD, CAT, POX, and GR enzymes could be ascribed to PCE's role in attempting to control ⁴ oxidative stress by reducing ion toxicity and the development of nucleoproteins that play a part in plant stress tolerance parameters. Taking into account all of the findings of this research, we infer that increasing drought and salinity stress treatments severely inhibited eggplant development by modifying multiple biochemical reactions in a differentiated manner. We found a clear relationship between stress-induced oxidative damage (caused by high MDA levels) and the accumulation of osmoprotectants, with stress resistance being regulated by the ability to effectively scavenge ROS from the environment.

Conclusion

The data gained from the study, we suggest that PCE obtained from Taif rose petals was advantageous and ²⁵ plays a crucial role in mitigating ⁴ the harmful impacts of drought and salt stress on eggplant physiological and biochemical features. It triggered the stimulation of antioxidative metabolism mechanisms in eggplants, resulting in increased drought and salt tolerance. All this data brings a clear piece of knowledge on the physiological and biochemical pathways that underpin the response of eggplants to drought and salinity. Finally, we can conclude that using PCE to alleviate drought and salinity stress while also increasing water usage efficiency and crop yield in dry environments where drought-tolerant eggplant is grown.

References

1. Xia, S., et al., *Application of polysaccharide biopolymer in petroleum recovery*. Polymers, 2020. **12**(9): p. 1860.
2. Bouissil, S., et al., *Use of alginate extracted from Moroccan brown algae to stimulate natural defense in date palm roots*. Molecules, 2020. **25**(3): p. 720.
3. Torres, F.G. and G.E. De-la-Torre, *Polysaccharide-based triboelectric nanogenerators: a review*. Carbohydrate Polymers, 2021. **251**: p. 117055.
4. Dutta, S. and M.S. Kumar, *Potential of value-added chemicals extracted from floral waste: A review*. Journal of Cleaner Production, 2021. **294**: p. 126280.
5. Vov, A., et al., *Combined recovery of polysaccharides and polyphenols from Rosa moscena wastes*. Industrial Crops and Products, 2017. **100**: p. 85-94.
6. Torres, F.G., et al., *Natural polysaccharide nanomaterials: an overview of their immunological properties*. International journal of molecular sciences, 2019. **20**(20): p. 5092.
7. Rotino, G., T. Sala, and L. Toppino, *Alien gene transfer in crop plants*. 2014, Springer Berlin, Germany:.
8. Saedifar, F., P. Ziarati, and Y. Ramezan, *Nitrate and Heavy Metal Contents in Eggplant (Solanum melongena) cultivated in the farmlands in the south of Tehran-Iran*. International Journal of Farming and Allied Sciences, 2014. **3**(1): p. 60-65.
9. Gürbüz, N., et al., *Health benefits and bioactive compounds of eggplant*. Food chemistry, 2018. **268**: p. 602-610.
10. Fu, Q., et al., *Leaf morphological and ultrastructural performance of eggplant (Solanum melongena L.) in response to water stress*. Photosynthetica, 2013. **51**(1): p. 109-114.
11. Karam, F., et al., *Yield and water use of eggplants (Solanum melongena L.) under full and deficit irrigation regimes*. Agricultural Water Management, 2011. **98**(8): p. 1307-1316.
12. Abbas, W., M. Ashraf, and N. Akram, *Alleviation of salt-induced adverse effects in eggplant (Solanum melongena L.) by foliar-applied natural and synthetic glycinebetaine*. Sci. Hort, 2010. **125**: p. 188-195.
13. Ünlükara, A., et al., *Effects of salinity on eggplant (Solanum melongena L.) growth and evapotranspiration*. Irrigation and Drainage: The Journal of the International Commission on Irrigation and Drainage, 2010. **59**(2): p. 203-214.
14. Hasan, M.M., et al., *Insights into 28-homobrassinolide (HBR)-mediated redox homeostasis, AsA-GSH cycle, and methylglyoxal detoxification in soybean under drought-induced oxidative stress*. Journal of Plant Interactions, 2020. **15**(1): p. 371-385.
15. Alabdallah, N.M., et al., *Green synthesized metal oxide nanoparticles mediate growth regulation and physiology of crop plants under drought stress*. Plants, 2021. **10**(8): p. 1730.
16. Alabdallah, N.M. and M.M. Hasan, *Plant-based green synthesis of silver nanoparticles and its effective role in abiotic stress tolerance in crop plants*. Saudi Journal of Biological Sciences, 2021. **28**(10): p. 5631-5639.
17. Hasan, M., et al., *EFFECTS OF MAGNETIZED WATER ON PHENOLIC COMPOUNDS, LIPID PEROXIDATION AND ANTIOXIDANT ACTIVITY OF MORINGA SPECIES UNDER DROUGHT STRESS*. JAPS: Journal of Animal & Plant Sciences, 2018. **28**(3).
18. Krasensky, J. and C. Jonak, *Drought, salt, and temperature stress-induced metabolic rearrangements and regulatory networks*. Journal of experimental botany, 2012. **63**(4): p. 1593-1608.

19. Hasan, M., et al., *Spermine: its emerging role in regulating drought stress responses in plants*. Cells, 2021. **10**(2): p. 261.
20. Hasan, M., et al., *GABA: A Key Player in Drought Stress Resistance in Plants*. International Journal of Molecular Sciences, 2021. **22**(18): p. 10136.
21. Jahan, M.S., et al., *Melatonin-mediated photosynthetic performance of tomato seedlings under high-temperature stress*. Plant Physiology and Biochemistry, 2021. **167**: p. 309-320.
22. Hasan, M.M., et al., *ABA-induced stomatal movements in vascular plants during dehydration and rehydration*. Environmental and Experimental Botany, 2021. **186**: p. 104436.
23. Barbosa, J.R. and R.N. de Carvalho Junior, *Polysaccharides obtained from natural edible sources and their role in modulating the immune system: Biologically active potential that can be exploited against COVID-19*. Trends in Food Science & Technology, 2021. **108**: p. 223-235.
24. Zhu, Z.-y., et al., *Effects of extraction methods on the yield, chemical structure and anti-tumor activity of polysaccharides from Cordyceps gunnii mycelia*. Carbohydrate Polymers, 2016. **140**: p. 461-471.
25. Arnon, D.I., *Copper enzymes in isolated chloroplasts. Polyphenoloxidase in Beta vulgaris*. Plant physiology, 1949. **24**(1): p. 1.
26. Heath, R., *Packer (1968) Photooxidation in isolated chloroplasts*. Arch Biochem Biophys. **125**: p. 189-198.
27. Bates, L.S., R.P. Waldren, and I. Teare, *Rapid determination of free proline for water-stress studies*. Plant and soil, 1973. **39**(1): p. 205-207.
28. Cock, J., S. Yoshida, and D.A. Forno, *Laboratory manual for physiological studies of rice*. 1976: Int. Rice Res. Inst.
29. Bradford, M.M., *A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding*. Analytical biochemistry, 1976. **72**(1-2): p. 248-254.
30. Mukherjee, S. and M. Choudhuri, *Implications of water stress-induced changes in the levels of endogenous ascorbic acid and hydrogen peroxide in Vigna seedlings*. Physiologia plantarum, 1983. **58**(2): p. 166-170.
31. Hasanuzzaman, M., M.A. Hossain, and M. Fujita, *Nitric oxide modulates antioxidant defense and the methylglyoxal detoxification system and reduces salinity-induced damage of wheat seedlings*. Plant Biotechnology Reports, 2011. **5**(4): p. 353-365.
32. Aebi, H., *[13] Catalase in vitro*, in *Methods in enzymology*. 1984, Elsevier. p. 121-126.
33. Hammerschmidt, R., E. Nuckles, and J. Kuć, *Association of enhanced peroxidase activity with induced systemic resistance of cucumber to Colletotrichum lagenarium*. Physiological Plant Pathology, 1982. **20**(1): p. 73-82.
34. Foyer, C.H. and B. Halliwell, *The presence of glutathione and glutathione reductase in chloroplasts: a proposed role in ascorbic acid metabolism*. Planta, 1976. **133**(1): p. 21-25.
35. Hasan, M., et al., *Spermine-Mediated Tolerance to Selenium Toxicity in Wheat (Triticum aestivum L.) Depends on Endogenous Nitric Oxide Synthesis*. Antioxidants, 2021. **10**(11): p. 1835.
36. Foyer, C.H., M. Lelandais, and K.J. Kunert, *Photooxidative stress in plants*. 1994, Wiley Online Library.
37. Urban, L., J. Aarrouf, and L.P. Bidel, *Assessing the effects of water deficit on photosynthesis using parameters derived from measurements of leaf gas exchange and of chlorophyll a fluorescence*. Frontiers in plant science, 2017. **8**: p. 2068.
38. Vighi, I., et al., *Functional characterization of the antioxidant enzymes in rice plants exposed to salinity stress*. Biologia plantarum, 2017. **61**(3): p. 540-550.

39. El-Banna, M. and K.A. Abdelaal, *Response of strawberry plants grown in the hydroponic system to pretreatment with H₂O₂ before exposure to salinity stress*. Journal of Plant Production, 2018. **9**(12): p. 989-1001.
40. Abdulmajeed, A.M., et al., *Alleviation of copper phytotoxicity by acetylsalicylic acid and nitric oxide application in mung bean involves the up-regulation of antioxidants, osmolytes and glyoxalase system*. Journal of Plant Interactions, 2021. **16**(1): p. 201-212.




Figure 1. Images of polysaccharide crude extract derived from Taif rose.




Figure 2: FTIR spectra identify a basic polysaccharide backbone with an intense peak: (3449 cm^{-1}). The broad peak signifies the stretching vibrations of the OH group [4]. (1651 cm^{-1}) represent the carbonyl group of a carboxylic acid group. The region below 800 cm^{-1} is known as the "skeletal region" and it is related to the carbohydrate skeletal vibrations.




Figure 3. HPLC chromatograms of glucose.




Figure 4. HPLC chromatograms of fructose.




Figure 5: Effect of polysaccharide crude extract, PCE (0, 15, 35, and 65mM) on Chla, Chlb and carotenoids in the leaf of eggplant seedlings under drought and salt stress. The data displayed are the means (\pm SE) of three replicates, and bars of dissimilar letters differ significantly at the $p = 0.05$ level.




Figure 6. Effect of polysaccharide crude extract, PCE (0, 15, 35, and 65mM) on total chl, chlorophyll fluorescence (Fv/Fm), and chlorophyll stability index (CSI) in the leaf of eggplant seedlings under drought and salt stress. The data displayed are the means (\pm SE) of three replicates, and bars of dissimilar letters differ significantly at the $p = 0.05$ level.




Figure 7. Effect of polysaccharide crude extract, PCE (0, 15, 35, and 65Mm) on MDA and proline content in the leaf of eggplant seedlings under drought and salt stress. The data displayed are the means (\pm SE) of three replicates, and bars of dissimilar letters differ significantly at the $p = 0.05$ level.




Figure 8. Effect of polysaccharide crude extract, PCE (0, 15, 35, and 65Mm) on total soluble protein (TSP) and total soluble sugar (TSS) in the leaf of eggplant seedlings under drought and salt stress. The data displayed are the means (\pm SE) of three replicates, and bars of dissimilar letters differ significantly at the $p = 0.05$ level.




Figure 9. Effect of polysaccharide crude extract, PCE (0, 15, 35, and 65Mm) on SOD and CAT activity in the leaf of eggplant seedlings under drought and salt stress. The data displayed are the means (\pm SE) of three replicates, and bars of dissimilar letters differ significantly at the $p = 0.05$ level.




Figure 10. Effect of polysaccharide crude extract, PCE (0, 15, 35, and 65Mm) on POX and GR activity in the leaf of eggplant seedlings under drought and salt stress. The data displayed are the means (\pm SE) of three replicates, and bars of dissimilar letters differ significantly at the $p = 0.05$ level.

Conflict of Interest

Conflict of Interest: None

Reviewers comments

Dear Editor,

The Plagiarism of the revised manuscript JKSUS-D-22-00151_R4 was found to be 17%. I am attaching the plagiarism report for your reference. Additionally, we have followed the author guidelines mentioned in your email. Please do the needful.

ORIGINALITY REPORT

17%

SIMILARITY INDEX

12%

INTERNET SOURCES

13%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	Saleh M Alluqmani, Nadiyah M Alabdallah. "Dry waste of red tea leaves and rose petals confer salinity stress tolerance in strawberry plants via modulation of growth and physiology", Journal of the Saudi Society of Agricultural Sciences, 2022 Publication	1%
2	www.tandfonline.com Internet Source	1%
3	mdpi-res.com Internet Source	1%
4	www.mdpi.com Internet Source	1%
5	"Drought Stress Tolerance in Plants, Vol 2", Springer Science and Business Media LLC, 2016 Publication	1%
6	www.agriculturejournals.cz Internet Source	1%

7	Saleh M. Alluqmani, Nadiyah M. Alabdallah. "Dry waste of red tea leaves and rose petals confer salinity stress tolerance in strawberry plants via modulation of growth and physiology", Journal of the Saudi Society of Agricultural Sciences, 2022	1 %
Publication		
8	Cervilla, Luis M., Begoña Blasco, Juan J. Rios, Miguel A. Rosales, Eva Sánchez-Rodríguez, Maria M. Rubio-Wilhelmi, Luis Romero, and Juan M. Ruiz. "Parameters Symptomatic for Boron Toxicity in Leaves of Tomato Plants", Journal of Botany, 2012.	1 %
Publication		
9	link.springer.com	1 %
Internet Source		
10	"The Plant Family Fabaceae", Springer Science and Business Media LLC, 2020	1 %
Publication		
11	Submitted to King Saud University	<1 %
Student Paper		
12	dx.doi.org	<1 %
Internet Source		
13	pericles.pericles-prod.literatumonline.com	<1 %
Internet Source		
14	ir.lib.uwo.ca	
Internet Source		

<1 %

15

Submitted to Universidad Estadual Paulista

Student Paper

<1 %

16

mafiadoc.com

Internet Source

<1 %

17

Submitted to Dr Rajendra Central Agricultural University

Student Paper

<1 %

18

arjournals.org

Internet Source

<1 %

19

etd.uwc.ac.za

Internet Source

<1 %

20

ajas.ubt.edu.al

Internet Source

<1 %

21

Duan, Zhen, Daiyu Zhang, Jianquan Zhang, Hongyan Di, Fan Wu, Xiaowen Hu, Xuanchen Meng, Kai Luo, Jiyu Zhang, and Yanrong Wang. "Co-transforming bar and CsALDH Genes Enhanced Resistance to Herbicide and Drought and Salt Stress in Transgenic Alfalfa (*Medicago sativa* L.)", *Frontiers in Plant Science*, 2015.

Publication

<1 %

22

Submitted to University of Southampton

Student Paper

<1 %

23

Liangliang Cai, Shanshan Zou, Dengpan Liang, Libiao Luan. "Structural characterization, antioxidant and hepatoprotective activities of polysaccharides from Sophorae tonkinensis Radix", Carbohydrate Polymers, 2018

Publication

<1 %

24

api.intechopen.com

Internet Source

<1 %

25

"Reactive Oxygen, Nitrogen and Sulfur Species in Plants", Wiley, 2019

Publication

<1 %

26

Chan-Su Rha, Hyeong Geun Kim, Nam-In Baek, Dae-Ok Kim, Cheon-Seok Park. "Amylosucrase from Deinococcus geothermalis can be modulated under different reaction conditions to produce novel quercetin 4'-O- α -D-isomaltoside", Enzyme and Microbial Technology, 2020

Publication

<1 %

27

Mohammad Shah Jahan, Md. Mahadi Hasan, Fahad S. Alotaibi, Nadiyah M. Alabdallah et al. "Exogenous Putrescine Increases Heat Tolerance in Tomato Seedlings by Regulating Chlorophyll Metabolism and Enhancing Antioxidant Defense Efficiency", Plants, 2022

Publication

<1 %

28

Wael M. Semida, Abdelsattar Abdelkhalik, Gamal. F. Mohamed, Taia A. Abd Abd El-Mageed et al. "Foliar Application of Zinc Oxide Nanoparticles Promotes Drought Stress Tolerance in Eggplant (*Solanum melongena* L.)", *Plants*, 2021

Publication

<1 %

29

repository.uwc.ac.za

Internet Source

<1 %

30

upcommons.upc.edu

Internet Source

<1 %

31

zenodo.org

Internet Source

<1 %

32

Luke O. Omoarelojie, Manoj G. Kulkarni, Jeffrey F. Finnie, Tomáš Pospíšil, Miroslav Strnad, Johannes Van Staden. "Synthetic strigolactone (rac-GR24) alleviates the adverse effects of heat stress on seed germination and photosystem II function in lupine seedlings", *Plant Physiology and Biochemistry*, 2020

Publication

<1 %

33

Nounjan, N.. "Exogenous proline and trehalose promote recovery of rice seedlings from salt-stress and differentially modulate antioxidant enzymes and expression of

<1 %

related genes", Journal of Plant Physiology,
20120415

Publication

34

academic.oup.com

Internet Source

<1 %

35

lipidworld.biomedcentral.com

Internet Source

<1 %

36

Mohammad Shah Jahan, Shirong Guo, Jin Sun, Sheng Shu et al. "Melatonin-mediated photosynthetic performance of tomato seedlings under high-temperature stress", Plant Physiology and Biochemistry, 2021

Publication

<1 %

37

journals.plos.org

Internet Source

<1 %

38

krishi.icar.gov.in

Internet Source

<1 %

39

Khaled A. A. Abdelaal, Yasser S.A. Mazrou, Yaser M. Hafez. "Silicon Foliar Application Mitigates Salt Stress in Sweet Pepper Plants by Enhancing Water Status, Photosynthesis, Antioxidant Enzyme Activity and Fruit Yield", Plants, 2020

Publication

<1 %

40

Nandita Bharadwaj, Sharmistha Barthakur, Akash Deep Biswas, Monoj Kumar Das, Manpreet Kour, Anand Ramteke, Nirmali

<1 %

Gogoi. "Transcript expression profiling in two contrasting cultivars and molecular cloning of a SKP-1 like gene, a component of SCF-ubiquitin proteasome system from mungbean *Vigna radiate* L.", *Scientific Reports*, 2019

Publication

41

Nihal E. Nahhas, Khaled A.A. Abdelaal, Muneera D.F. AlKahtani, Latifa Al Husnain et al. "Biochar and jasmonic acid application attenuate antioxidative systems and improves growth, physiology, nutrient uptake and productivity of faba bean (*Vicia faba* L.) irrigated with saline water", *Plant Physiology and Biochemistry*, 2021

Publication

<1 %

42

Qiyu Luo, Shu Chen, Jiazheng Zhu, Laihua Ye, Nathan Daniel Hall, Suma Basak, J. Scott McElroy, Yong chen. " Overexpression of confers paraquat-tolerance in rice L.) by promoting polyamine pathway ", *Cold Spring Harbor Laboratory*, 2020

Publication

<1 %

43

Sónia Silva, Maria Celeste Dias, Artur M. S. Silva. "Titanium and Zinc Based Nanomaterials in Agriculture: A Promising Approach to Deal with (A)biotic Stresses?", *Toxics*, 2022

Publication

<1 %

- | | | |
|----|--|------|
| 44 | academic-accelerator.com
Internet Source | <1 % |
| 45 | edepot.wur.nl
Internet Source | <1 % |
| 46 | v3r.esp.org
Internet Source | <1 % |
| 47 | www.scipress.com
Internet Source | <1 % |
| 48 | www.wjgnet.com
Internet Source | <1 % |
| 49 | "Introducing Article Numbering to Journal of King Saud University - Science", Journal of King Saud University - Science, 2021
Publication | <1 % |
| 50 | Nadiyah M. Alabdallah, Md. Mahadi Hasan, Abdalrhaman M. Salih, S.S. Roushdy et al. "Silver nanoparticles improve growth and protect against oxidative damage in eggplant seedlings under drought stress", Plant, Soil and Environment, 2021
Publication | <1 % |
| 51 | Neveen B. Talaat. "Effective Microorganisms Improve Growth Performance and Modulate the ROS-Scavenging System in Common Bean (<i>Phaseolus vulgaris</i> L.) Plants Exposed to | <1 % |

Salinity Stress", Journal of Plant Growth Regulation, 2014

Publication

52

Sameer Gupta, Chandana Haldar, Raise Ahmad. "Photoperiodic regulation of nuclear melatonin receptor ROR α in lymphoid organs of a tropical rodent *Funambulus pennanti*: Role in seasonal oxidative stress", Journal of Photochemistry and Photobiology B: Biology, 2015

Publication

<1 %

53

Yang, F.. "Salt stress responses in *Populus cathayana* Rehder", Plant Science, 200905

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Dr. Nadiyah_JKSUS-D-22-00151_R4

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19

PAGE 20

PAGE 21

PAGE 22

PAGE 23

PAGE 24

PAGE 25

PAGE 26

PAGE 27

PAGE 28

PAGE 29

PAGE 30

PAGE 31

PAGE 32

PAGE 33

PAGE 34

PAGE 35
