

# UPDATED Manuscript Ni and Cd (1) (5).docx

*by Dr Anish Khan*

---

**Submission date:** 08-May-2022 07:48PM (UTC+0800)

**Submission ID:** 1831087330

**File name:** UPDATED\_Manuscript\_Ni\_and\_Cd\_1\_5.docx (48.28K)

**Word count:** 3523

**Character count:** 21350

32

**An in-vitro anti-inflammatory and anti-microbial essential on Ni(II), Cd(II) mixed ligand complexes by using 2,4-dinitrophenyl hydrazine and dimethylglyoxime**

**Abstract**

The objective of the present work deals among synthesis as well as characterization of Ni(II), Cd(II) mixed ligand complex were derived from 2,4 di-nitrophenyl hydrazine and dimethyl glyoxime. The mixed ligand metal complexes were prepared via 2,4-dinitrophenyl hydrazine as well as dimethylglyoxime among Ni(II) and Cd(II) in molar ratio (M: L1: L1) 1:1:1. CHNO analysis, conductivity measurement, PXRD, SEM, UV-visible, infrared spectrum, and magnetic susceptibility, thermal analysis were used to characterized the synthesized compounds. According to the study of the results, all of the metal complexes have octahedral geometry. The molar conductance value indicate that non-electrolytic character of compounds Ni(II), Cd(II). According to the powder X-ray diffraction (PXRD) patterns, all of the compounds are crystalline character. Moreover the Ni(II), Cd(II) complexes were very stable in terms of thermal stability. Metal complexes have been tested against Gram-positive (*S. aureus*, *B. subtilis* and *A. niger*) and Gram-negative (*P. aeruginosa*, *E. coli* and *C. albicans*) pathogenic bacteria as well as fungus. As a result, the metal complexes have reported strong antimicrobial efficiency.

**Conclusions;** Furthermore, applications of metal compounds were tested for anti-inflammatory efficiency utilize egg albumin. Result shows that the activities of metal complexes are good.

**Keywords:** Ni(II)-DNPH-DMG complex; Cd(II)-DNPH-DMG complex; thermal analysis of mixed ligand metal complexes, Antimicrobial; and anti-inflammatory assay

## 1. Introduction

Many researchers are focusing on mixed chelating metal complexes because they are more important than ever in pharmacological applications such as antimicrobial and fungal, antitumor, anti-inflammatory, and anticancer (Siddiki et al., 2021). Recent evaluations in the fields of bioinorganic and medicinal chemistry contain enhanced the coordination chemistry of transition metal ions using several variety of ligands (E.-Z. Hossain MS, Banu LA, 2019). In an important field of research, transition elements play a vital part in the use of transition metals complexes as medicines for the treatment of numerous disorders (Muthuppalani et al., 2022). Metal complexes that are stable and safe among active metal sites are useful as biological probes (Raman et al., 2012). The environment roughly the metals centre, such as coordination structures, ligands, and donor classes, is one of the most important aspects for metallo enzymes to accomplish their specific physiological function (Sujamol et al., 2010). The 2,4-dinitrophenylhydrazine (DNPH) is an important component in biological and pharmacological processes (Devi et al., 2010). It is a more over essential class of medicines counterpart of  $H_2NNH_2$  (hydrazine) (Nasiru, 2012). N (Nitrogen) and O (Oxygen) can be brought together since they are both beneficial to bio-inorganic processes (Journal, 2011). Dimethylglyoxime was among the most widely utilised ligand for coordinating nickel ions in priority, which also forms Ni-DMG compounds due to attribute specificity and consistency (M.R.K. and S.L.B. Padma Rao, Ch. V., Praveen K., Kishorebabu B., Padma M., Anna Sudheer K., Sandhya Rani K., Koteswarao K., Suseelabai G., 2013). The oxime molecule ( $>C=N-OH$ ) is amphoteric, with slightly basic nitrogen as well as moderately acidic hydroxyl groups, and is thought to be formed from oxy-imine. Many transition metals in the Periodic Table react on dioxime ligands to generate extremely reliable complexes. Furthermore, the Dioximes can coordinate through the oxime groups' N, N or N, O sites. As a result, some Dioximes compounds showed considerable anti - carcinogenic and therapeutic efficacy (Shaker, 2010).

The current research has developed Ni(II) and Cd(II) compounds utilizing 2,4-dinitrophenylhydrazine (DNPH) with dimethyl glyoxime (DMG), moreover physico-chemical characteristics and also to investigate the biological capability of DNPH with compounds of Ni(II), Cd(II).

## 2. Experimental

## 2.1. Materials

All of the chemicals have been obtained from commercial sources and used without additional purified process ( $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ ,  $\text{CdCl}_2 \cdot 6\text{H}_2\text{O}$  and dimethyl glyoxime (DMG) from Merck as well as 2,4-dinitrophenylhydrazine from Sigma Aldrich, and also solvents such as dimethyl sulfoxide (DMSO), dimethyl formaldehyde (DMF), Ethanol, Acetone, as well as Methanol, utilized as an AnalaR grad.

## 2.2. physicochemical characterization

The Elementar Vario EL III CHNO analyzer was used to accomplish the elemental analysis. A conductivity evaluation became observed from (Equip – Tronics, eq-661A) analyser at room temperature in DMF medium ( $10^{-3}\text{M}$ ). Infra-red spectrum was obtained utilising Agilent spectrophotometer in the wavelength from 4000 to  $400\text{ cm}^{-1}$  based on the KBr pellet method. UV-visible absorption spectra were collected to use a JASCO V-630 with a 1nm optical resolution with wavelengths range between 300 to 800 nm, whereas PXRD patterns were obtained using a Shimadzu PW3050/60 with  $\text{CuK}\alpha_1 = 1.5406$  and  $\text{CuK}\alpha_2 = 1.5444$ . Thermal analysis was carried out (Perkin Elmer thermal analyzer) and the SEM photographs were taken with a ZEISS apparatus. Furthermore the magnetic behavior evaluated from (Model MsB-MK1) using Guey's Balance.

## 2.3. Preparation of Ni(II), and Cd(II) complexes

Mixed ligand Ni(II) as well as Cd(II) complexes have been obtained via adding a hot DMF solution of (10mM) DNPH dropwise to an EtOH solution of Nickel(II), Cadmium(II) chloride (10mM) and ethanol solution of (10mM) DMG mixture (1:1:1) ratios. At  $60^\circ\text{C}$ , the reaction mixture was stirred in addition to refluxed for four hours. Finally, a colored complexes scheme – 1 precipitation was obtained and rinsed with EtOH solutions before even being dried under vacuum onto anhydrous  $\text{CaCl}_2$  desiccators.

## 2.4. In-vitro antimicrobial studies

Disc dispersal was utilized to investigate the antimicrobial potential of Ni(II) and Cd(II) complexes against a variety of gram (positive, negative) bacteria as well as fungals (*C. albicans*, *A. niger*, *B. subtilis*, *S. aureus*, *P. aeruginosa* and *E. coli*). Ni(II), Cd(II) components was diluted in DMSO. Individually, the solution variations ( $30, 60\ \mu\text{g/ml}$ ) were set up. The prepared disks were placed in petri plates containing nutrient medium seeded through each bacterial and fungal serum individually

after being dipped in a specific variant of complexes. Every plat was incubated at 37<sup>0</sup>C for 24 and 48 hours for bacterial and fungal growth, as well as the zone of inhibition value was recorded (Aly and Fathalla, 2020; Shukla and Mishra, 2019).

#### 2.4.1. Microorganisms

The microorganisms species employed in the biological tests included E. coli (MTCC 732), Staphylococcus aureus (MTCC 3160), Pseudomonas aeruginosa (MTCC 424), and Bacillus subtilis (MTCC 441), while the fungal species are Aspergillus niger (MTCC 10180), as well as Candida albicans (MTCC 10183). The Institute of Molecular Biology and biotechnology (IMTECH) in Chandigarh, India, provided the MTCC (microbial variety cultured collections).

#### 2.4.2. Preparation of dried filter paper discs

Filter papers (No. 1) have been used to form disc with a 6 mm diameter that were sterilised in warm air. Upon sterilisation, discs was supplied containing 30µg/ml and 60µg/ml from each sample, with a 60µg/ml chloramphenicol as well as fluconazole standard stock solution and a 60µg/ml control solution being used correlate the sample solution. They are preserved cold even though they were being used in the test.

#### 2.4.3. In-vitro Antimicrobial evaluations

Antimicrobial was accomplished via disc dispersal technique utilising tests. 30 ml of Nutrient agar (NA) medium as well as PDA (Potato dextrose agar) standard was poured into petri plates. Using a micropipette, the experiment organism were injected on consolidated agar platter and distributed for 10 mints before drying. Bacteria from a broth culture were used to isolate the surface areas of the medium. Before evenly inoculating the whole facade of the NA/PDA platter, a sterilized cotton brush is soaked within a standardised microbiological experiment mixture. In a nutshell, inoculum of microbe's embrace were diffuse on agar platter. The germ-free filter sheets (six millimeter width) comprising disks have been loaded to 30µg/ml as well as 60µg/ml of every sample, 60µg/ml of control, and 60µg/ml from each sample utilising sterile forceps. A standard solution was placed on the area of the infused agar platter. Bacterial strains were infused on 37 degrees Celsius at 24 hours, whereas fungal strains were incubated for 48 hours. Each sample was examined three times.

#### 2.4.4. Measurement of inhibition zones

The antibacterial property of the evaluation substances had been calculated using the mean diameter of the inhibition zones around the disc in mm. A millimeter scale was used to assess the inhibition zones of the microorganisms evaluated by the substances.

## 2.5. Anti-inflammatory properties of metal complexes

### 2.5.1. Inhibition of Albumin Denaturation:

0.2 ml new hen's egg albumin, 2.8 ml phosphate buffered saline (PBS, pH 6.4), as well as 2 ml of different metal compound ratios (100, 200, 300, 400, and 500 µg/ml) were included in the reaction mixture (5 ml). As a control, similar amounts of double-distilled water were used. The mixes were then heated on 70 degrees Celsius for five minutes behind being incubated at (37°C) in an incubator for 15 minutes. After cooling, the absorbance was in use at 660 nm to use a solution as a control. For the purposes of evaluating absorbance, diclofenac sodium was used as a reference drug, with final concentrations ranging from 100 to 500µg/ml. The percentage inhibitions of protein denaturation were determined using the method (Hasan, 2019):

$$\text{Percentage of inhibition (\%)} = [1 - \frac{V_t}{V_c}] \times 100 \quad (1)$$

wherever,

$V_c$  = absorbance of control

$V_t$  = absorbance of experiment sample.

Three different tests were conducted in triplicate. The quantity of sample required inhibiting concentration by 50%, or  $IC_{50}$ , was graphically calculated using Ms-Windows based graph pad Instate software using a linear regression technique. The result was represented graphically as average standard deviation.

## 3. Result and discussion

The physical characteristics of metal complexes, together with their analytical dates, are given through Table - 1. A Ni(II), Cd(II) compounds were also stabilised at room temperature. The complexes are solubility using DMF and DMSO, but not in other organic solvents. The empirical values of C.H.N.S. analysis of ligand and metal compounds are in good agreement with theoretical computed values, with complexes ratios of 1:1:1 between metal and ligands.

27

### 3.1. Molar conductance

Molar conductance values of the metal complex were measured at  $1 \times 10^{-3}$  M accumulation of DMF solvent as well as entire compounds showed conductance in the range of  $25.47 - 20.39 \Omega^{-1} \text{ cm}^2 \text{ mol}^{-1}$  at  $37^\circ\text{C}$  was shown in (Table – 2). The lowest conductivity value suggested that Ni(II) and Cd(II) complexes were non-electrolytes (Manimaran et al., 2021).

18

### 3.2. FT-IR Spectra Analysis of Nickel(II) and Cd(II) mixed ligand Complexes

FTIR spectrum of the Ni(II) and Cd(II) mixed ligand complexes, to investigated mode of bonding DNP and DMG ligands to Ni(II), Cd(II) ion. Figure – 1 shows the characteristic bands of a infrared spectra of mixed ligand metal complexes. FTIR spectrum of DNP (2,4-dinitrophenyl hydrazine) revealed sharp peaks at  $3324 \text{ cm}^{-1}$  due to  $\nu(\text{N-H})$  wavelength shifted to the a higher frequency on  $3350 - 3310 \text{ cm}^{-1}$  shows nitrogen was involved in the coordination of the center metal atom. The  $\nu(\text{NO})$  band was shows on  $1320 \text{ cm}^{-1}$  were observed from DNP, furthermore these band was shift toward the lower in  $25 - 21 \text{ cm}^{-1}$  in the complexes. These shifts be able to explained by the ligand was coordinated with metal ions through oxygen (Anantha Lakshmi et al., 2008). At lower wavelength  $3100 - 3092 \text{ cm}^{-1}$  the wide band about  $3211 \text{ cm}^{-1}$ , which was assigned to  $\nu(\text{O-H})$  stretching modes in DMG, was detected on metal complexes spectrum. The impact of coordination of the oxygen atom of the OH group following deprotonation was attributed to this observation (Devi et al., 2018). The lower frequency some new band appear in metal complexes observed at  $475 - 416 \text{ cm}^{-1}$  and  $559 - 530 \text{ cm}^{-1}$  that has related to M – N and also M – O significantly (Adly et al., 2020; Ferraz et al., 2009).

34

### 3.3. Electronic Spectral data and magnetic moments

The electronic spectrum of Ni(II) and Cd(II) mixed ligand complexes data illustrated in table – 2 and Figure – 2. The electronic spectra of the Ni(II) complex exhibits a strong absorption band at 384, 465 nm which is according to  $n \rightarrow \pi^*$  and  ${}^3\text{A}_2\text{g} - {}^3\text{T}_1\text{g}(\text{P})$ . Transitions imply that the metal ion is surrounded by an octahedral geometry. The magnetic moment of the Ni(II) complex measured 3.14 B.M, confirming the metal complex's octahedral geometry (Anantha Lakshmi et al., 2008).

21

The absorption band at 379, 452 nm in the electronic spectra of the Cd(II) complex may be ascribed to the  $n \rightarrow \pi^*$  and LMCT. Transitions imply that the metal ion is surrounded by an octahedral

geometry. The diamagnetic nature of the Cd(II) complex may be seen in its magnetic moment value (Sulekh Chandra and Monika Tyagi, 2009).

### 3.4. PXRD analysis

Figure – 3a&b shows the PXRD (powder X-ray diffraction) study of Ni(II), Cd(II) compounds in the range of  $2\theta = 10 - 80^\circ$ . It's described as prominent crystalline peaks seen in metal complexes that reveal the complexes' crystalline nature. Debye Scherrer's equation was used to compute the crystallite size of the complexes (Netalkar et al., 2014).

$$D = 0.9\lambda / \beta \cos\theta \quad (2)$$

Where,

Constant as 0.9,  $\lambda = (\text{Cu K}\alpha = 1.5406 \text{ \AA})$  wavelength of X-ray radiation,  $\beta = \text{Full-width half maximum (FWHM)}$  and  $\theta = \text{diffraction angle}$ . The average crystallite sizes of the Ni(II) and Cd(II) complexes are around 56.74 and 69.34 nm respectively.

### 3.5 Morphological observations

The SEM micrographs revealed a Ni(II) and Cd(II) complex are given in Figure – 3c&d, spherical-like structure with particle diameter of estimated 55 and 65  $\mu\text{m}$ , with in the scale bar 1  $\mu\text{m}$  respectively. All these metal complexes surfaced to have showed an aggregated rod-like arrangement and slightly cluster nature.

### 3.6 Thermal property of metal complexes

TG (Thermogravimetric) as well as DTG (differential thermogravimetric) assessment of Ni(II), Cd(II) complexes curves are shown in Figure – 4a&b. Thermal dissolution of complexes of characteristic  $[\text{M}(\text{L})_2]$  consists of three phases:  $\text{M} = \text{Ni(II), Cd(II)}$ , and  $n = 1, 2, 3$ . The first process corresponds to the loss of moisture in the temperature varied 30-100°C; the important phase of decomposition takes place among 200 & 157°C, with a mass reduction of estimated 75.4% , 80.3% of the ligand; and the final phase occurs up to 201-340°C and 201-400, with the residual ligands being dropped. After several stages of decomposition, the complexes were transformed to their correlating metal oxides at extreme temps.

### 3.7. In-vitro Antimicrobial efficiency of Nickel and Cadmium complexes



The result of antimicrobial behavior of Ni(II) and Cd(II) complexes by using disc diffusion method is summarized table – 3. Synthesized metal complexes good active screened against gram positive and gram negative bacterial and fungal. The Ni(II) complex good active against the bacteria B. subtilis and E.coli in 30 as well as 60  $\mu\text{g/ml}$  and less active against fungal species. Furthermore, the Cd(II) complex high activities against bacteria p. aeruginosa and fungi c. albicans in 60  $\mu\text{g/ml}$ . The complexes had stronger antimicrobial property in higher concentration are shows in Figure – 5a. The higher activities of metal compounds are explain using the chelation theory(Gupta and Fahmi, 2016; Manimohan et al., 2020), which states that chelation reduces the polarity of metal ions by exchanging a portion of their positive charge by the ligand. This enhances lipid solubility, making it easier for it to permeate into microorganisms normal cells(Saddam Hossain, 2018; Tadavi et al., 2020). Organic ligands have donor sites which are active against the bio-potential activities the enhanced biological activities due to the presence of oxygen as well as nitrogen donor sites in DMG, DNPH.

### 3.8. *In-vitro anti-inflammatory activities*

An anti-inflammatory behavior of Ni(II), Cd(II) compounds was investigated in protein (bovine serum albumin) denaturation technique used with various concentration (100, 200, 300, 400 & 500 $\mu\text{g/ml}$ ), diclofenacsodium used as a reference drug. Table – 4 as well as Figure– 5b indicate the % of anti-inflammatory efficiency was computed moreover compared to the reference drug. Three part trial were carried out as well as  $\text{IC}_{50}$  was calculated. The results indicating the diclofenac sodium (96.05%)  $\text{IC}_{50}$  is 230.75. A Nickel compound (84.56%) $\text{IC}_{50}$  value is 257.31 superior protein denaturation compound compare to the Cadmium complex (79.55 %)  $\text{IC}_{50}$  value is 270.83. the metal complexes lower activity comparing to standard drug. Due to the occurrence of donor ('N', 'O') places into the DNPH and DMG contained in the compounds which demonstrate the good potential of complexes. The order of anti-inflammatory activity diclofenacsodium > Ni(II) complex> Cd(II) complexes (Szczepaniak and Fichna, 2019).

## 4. Conclusion

In this artwork, Ni(II), Cd(II) mixed ligand complexes were satisfactorily synthesised and characterised employing a multitude of physicochemical approaches. All complexes exhibited 1:1

9  
electrolyte in essence and metal complexes was initiated to have an octahedral geometry based on magnetic moment, colour, UV-vis spectral, TG inferences. And powder XRD analysis, which shows that the average crystalline diameters are between 56.74 and 69.34 nm. The mixed ligand complexes were extremely stable in terms of thermal stability. Thus the study found that the mixed ligand Ni(II) and Cd(II) complexes have significant antibacterial action when compared to conventional antibiotics. Furthermore the Ni (II), Cd(II) complexes have showed excellent inflammation properties compared to the standard diclofinac.

#### Acknowledgements

1  
The Deanship of Scientific Research (DSR) at King Abdulaziz University, Jeddah Saudi Arabia has funded this project, under grant number (KEP-40-130-42).

#### References:

- Adly, O.M.I., Shebl, M., Abdelrhman, E.M., El-Shetary, B.A., 2020. Synthesis, spectroscopic, X-ray diffraction, antimicrobial and antitumor studies of Ni(II) and Co(II) complexes derived from 4-acetyl-5,6-diphenyl-3(2H)-pyridazinone and ethylenediamine. *J. Mol. Struct.* 1219, 128607. <https://doi.org/10.1016/j.molstruc.2020.128607>
- Aly, S.A., Fathalla, S.K., 2020. Preparation, characterization of some transition metal complexes of hydrazone derivatives and their antibacterial and antioxidant activities. *Arab. J. Chem.* 13, 3735–3750. <https://doi.org/10.1016/j.arabjc.2019.12.003>
- Anantha Lakshmi, P.V., Saritha Reddy, P., Jayatyaga Raju, V., 2008. Synthesis and structural studies of first row transition metal complexes of N-(2-nitro)-benzilidine-3-hydrazino quinoxaline-2-one. *Bull. Chem. Soc. Ethiop.* 22. <https://doi.org/10.4314/bcse.v22i3.61215>
- Devi, J., Yadav, M., Sharma, S., 2018. Synthesis and Characterization of Transition Metal(II) Schiff Bases Complexes Derived from 2,5-Dihalosalicylaldehyde and 4-Methyl-3-thiosemicarbazide. *Asian J. Chem.* 30, 2445–2449. <https://doi.org/10.14233/ajchem.2018.21438>
- Devi, T.U., Lawrence, N., Babu, R.R., Ramamurthi, K., Bhagavannarayana, G., 2010. Structural and Optical Characterization Studies on 2, 4- dinitrophenylhydrazine Single Crystal. *J. Miner. Mater.*

Charact. Eng. 09, 321–330. <https://doi.org/10.4236/jmmce.2010.94023>

E.-Z. Hossain MS, Banu LA, H.M.M., 2019. Synthesis, Characterization and Biological Activity Studies of Mixed Ligand Complexes with Schiff base and 2,2'-Bipyridine. *Int. J. Appl. Sci. - Res. Rev* 6, 1–7.

Ferraz, K.O., Wardell, S.M.S.V., Wardell, J.L., Louro, S.R.W., Beraldo, H., 2009. Copper(II) complexes with 2-pyridineformamide-derived thiosemicarbazones: Spectral studies and toxicity against *Artemia salina*. *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.* 73, 140–145. <https://doi.org/10.1016/j.saa.2009.02.020>

Gupta, B., Fahmi, N., 2016. Co(II) and Ni(II) complexes with Schiff base ligands: Synthesis, characterization, and biological activity. *Russ. J. Gen. Chem.* 86, 1182–1190. <https://doi.org/10.1134/S1070363216050339>

Hasan, A.U.H., 2019. Evaluation of in vitro and in vivo therapeutic efficacy of *Ribes alpestre* Decne in Rheumatoid arthritis. *Brazilian J. Pharm. Sci.* 55. <https://doi.org/10.1590/s2175-97902019000217832>

Journal, B.S., 2011. The preparation and characterization of some metal complexes with tridentate ONO ligand derived from phenyl hydrazine. *Baghdad Sci. J.* 8, 796–805. <https://doi.org/10.21123/bsj.8.3.796-805>

M.R.K. and S.L.B. Padma Rao. Ch. V., Praveen K., Kishorebabu B., Padma M., Anna Sudheer K., Sandhya Rani K., Koteswarao K., Suseelabai G., V.R.B., 2013. Zinc dimethylglyoxime complexes. *Der Pharma Chem.* 5, 280–284.

Manimaran, P., Balasubramaniyan, S., Azam, M., Rajadurai, D., Al-Resayes, S.I., Mathubala, G., Manikandan, A., Muthupandi, S., Tabassum, Z., Khan, I., 2021. Synthesis, Spectral Characterization and Biological Activities of Co(II) and Ni(II) Mixed Ligand Complexes. *Molecules* 26, 823. <https://doi.org/10.3390/molecules26040823>

Manimohan, M., Pugalmani, S., Sithique, M.A., 2020. Synthesis, Spectral Characterisation and Biological Activities of Novel Biomaterial/N, N, O Donor Tridentate Co (II), Ni (II) and Zn (II) Complexes of Hydrazide Based Biopolymer Schiff Base Ligand. *J. Inorg. Organomet. Polym. Mater.* 30, 4481–4495. <https://doi.org/10.1007/s10904-020-01578-7>

- Muthuppalani, M., Otaibi, A. Al, Balasubramanian, S., Manikandan, S., Manimaran, P., Mathubala, G., Manikandan, A., Arshad, M.N., Puttegowda, M., Alorfi, H.S., Khan, A., Asiri, A.M., Rahman, M.M., 2022. Synthesis, Characterization and Bio-Potential Activities of Co(II) and Ni(II) Complexes with O and N Donor Mixed Ligands. *Crystals* 12, 326. <https://doi.org/10.3390/cryst12030326>
- Nasiru, N.P.N. and Y.P., 2012. Complexes Of Cobalt(Ii), Nickel(Ii) And Zinc(Ii) With Schiff Bases Derived From 4-Anisaldehyde. *Int. J. Pharm. Sci. Res* 3, 5116–5120.
- Netalkar, P.P., Netalkar, S.P., Revankar, V.K., 2014. Nickel(II) complexes of thiosemicarbazones: synthesis, characterization, X-ray crystallographic studies and in vitro antitubercular and antimicrobial studies. *Transit. Met. Chem.* 39, 519–526. <https://doi.org/10.1007/s11243-014-9827-8>
- Raman, N., Sobha, S., Selvaganapathy, M., Mahalakshmi, R., 2012. DNA binding mode of novel tetradentate amino acid based 2-hydroxybenzylidene-4-aminoantipyrine complexes. *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.* 96, 698–708. <https://doi.org/10.1016/j.saa.2012.07.051>
- Saddam Hossain, M., 2018. Metal Complexes as Potential Antimicrobial Agent: A Review. *Am. J. Heterocycl. Chem.* 4, 1. <https://doi.org/10.11648/j.ajhc.20180401.11>
- Shaker, S.A., 2010. Preparation and Spectral Properties of Mixed-Ligand Complexes of VO(IV), Ni(II), Zn(II), Pd(II), Cd(II) and Pb(II) with Dimethylglyoxime and N -acetyl glycine. *E-Journal Chem.* 7, S580–S586. <https://doi.org/10.1155/2010/125136>
- Shukla, S., Mishra, A.P., 2019. Metal complexes used as anti-inflammatory agents: Synthesis, characterization and anti-inflammatory action of VO(II)-complexes. *Arab. J. Chem.* 12, 1715–1721. <https://doi.org/10.1016/j.arabjc.2014.08.020>
- Siddiki, A.K.M.N.A., Islam, S., Begum, S., Salam, M.A., 2021. Synthesis, spectral characterization, thermal behavior and biological activities study of ternary metal complexes of alanine and 1,8-diaminonaphthalene with Co(III), Ni(II), Cu(II), Zn(II) and Cd(II). *Mater. Today Proc.* 46, 6374–6381. <https://doi.org/10.1016/j.matpr.2020.06.126>
- Sujamol, M.S., Athira, C.J., Sindhu, Y., Mohanan, K., 2010. Synthesis, spectroscopic characterization, electrochemical behaviour and thermal decomposition studies of some transition metal complexes

with an azo derivative. *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.* 75, 106–112.

<https://doi.org/10.1016/j.saa.2009.09.050>

Sulekh Chandra and Monika Tyagi, 2009. Synthesis and antimicrobial activities of n-s donor ligand and its transition metal complexes. *Int. J. Chem. Sci* 7, 116–124.

Szczepaniak, A., Fichna, J., 2019. Organometallic Compounds and Metal Complexes in Current and Future Treatments of Inflammatory Bowel Disease and Colorectal Cancer—a Critical Review. *Biomolecules* 9, 398. <https://doi.org/10.3390/biom9090398>

Tadavi, S.K., Bendre, R.S., Patil, S.V., Gaguna, S., Rajput, J.D., 2020. Synthesis, crystal structures and antimicrobial activity of palladium metal complexes of sulfonyl hydrazone ligands. *Eur. J. Chem.* 11, 377–384. <https://doi.org/10.5155/eurjchem.11.4.377-384.2040>

ORIGINALITY REPORT

15%

SIMILARITY INDEX

10%

INTERNET SOURCES

13%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1%
2	Walaa H. Mahmoud, Reem G. Deghadi, Gehad G. Mohamed. "Preparation, geometric structure, molecular docking thermal and spectroscopic characterization of novel Schiff base ligand and its metal chelates", Journal of Thermal Analysis and Calorimetry, 2016 Publication	1%
3	<a href="http://www.derpharmachemica.com">www.derpharmachemica.com</a> Internet Source	1%
4	M. Muthuppalani, Ahmed Al Otaibi, S. Balasubramanian, S. Manikandan et al. "Synthesis, Characterization and Bio-Potential Activities of Co(II) and Ni(II) Complexes with O and N Donor Mixed Ligands", Crystals, 2022 Publication	1%
5	Submitted to University of Central Lancashire Student Paper	1%
6	<a href="http://www.chemijournal.com">www.chemijournal.com</a> Internet Source	1%

---

7 Muhammad Furqan Akhtar, Syed Ahmad Raza, Ammara Saleem, Irfan Hamid et al. "Appraisal of Anti-Arthritic and Anti-Inflammatory Potential of Folkloric Medicinal Plant Peganum harmala", Endocrine, Metabolic & Immune Disorders - Drug Targets, 2022  
Publication 1 %

---

8 S.A. Aly, S.K. Fathalla. "Preparation, characterization of some transition metal complexes of hydrazone derivatives and their antibacterial and antioxidant activities", Arabian Journal of Chemistry, 2020  
Publication 1 %

---

9 [www.science.gov](http://www.science.gov)  
Internet Source 1 %

---

10 Laijin Tian, Bin Zhao, Zhengyu Zhou, Qingsen Yu, Wentao Yu. "SYNTHESIS AND CHARACTERIZATION OF TRANSITION METAL TERNARY COMPLEXES OF GLYOXYLIC ACID THIOSEMICARBAZONE AND 4,5-DIAZAFLUOREN-9-ONE", Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry, 2002  
Publication 1 %

---

11 [aip.scitation.org](http://aip.scitation.org)  
Internet Source 1 %

---

12	<a href="http://innovareacademics.in">innovareacademics.in</a> Internet Source	<1 %
13	<a href="http://link.springer.com">link.springer.com</a> Internet Source	<1 %
14	K. Velsankar, G. Parvathy, S. Mohandoss, R. Mohan Kumar, S. Sudhahar. "Green synthesis and characterization of CuO nanoparticles using Panicum sumatrense grains extract for biological applications", Applied Nanoscience, 2022 Publication	<1 %
15	J. Jeevitha Rani, A. Mary Imelda Jayaseeli, S. Rajagopal, S. Seenithurai, Jeng-Da Chai, J. Dhaveethu Raja, R. Rajasekaran. "Synthesis, characterization, antimicrobial, BSA binding, DFT calculation, molecular docking and cytotoxicity of Ni(II) complexes with Schiff base ligands", Journal of Molecular Liquids, 2021 Publication	<1 %
16	<a href="http://www.gssrr.org">www.gssrr.org</a> Internet Source	<1 %
17	"Promising Antimicrobials from Natural Products", Springer Science and Business Media LLC, 2022 Publication	<1 %



18 Asma I. El - Said. "Binary and Mixed Ligand Complexes of Nifuroxime, Spectral, and Thermal Studies", Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry, 2003  
Publication <1 %

---

19 Pragnesh K. Panchal, D. H. Patel, M. N. Patel. "Preparation and Thermal, Spectroscopic, and Antibacterial Studies of Some Mixed - Ligand Complexes", Synthesis and Reactivity in Inorganic and Metal-Organic Chemistry, 2004  
Publication <1 %

---

20 Ravinder, Dipak Kumar Das, Vinod Kumar Vashistha, Anuj Kumar. "Synthesis, Electrochemical and Antibacterial Studies of Hexa-aza-macrocyclic Complexes of Ni(II) and Cu(II) Ions", Nano LIFE, 2022  
Publication <1 %

---

21 [research.asu.edu.eg](http://research.asu.edu.eg)  
Internet Source <1 %

---

22 Amna Zidan. "Mixed Ligand Complexes of Nickel (II) Dialkyldithiophosphates with 2-Acetylpyridine Semicarbazone and 2-Acetylpyridine Benzoylhydrazone", Phosphorus, Sulfur, and Silicon and the Related Elements, 2003  
Publication <1 %

---

23

Moamen S. Refat. "Spectroscopic, thermal and biocidal studies on Mn(II), Co(II), Ni(II) and Cu(II) complexes of tridentate ligand having semicarbazone moieties", Journal of Thermal Analysis and Calorimetry, 08/28/2009

Publication

<1 %

24

[ijppjournal.com](http://ijppjournal.com)

Internet Source

<1 %

25

[journalrepository.org](http://journalrepository.org)

Internet Source

<1 %

26

Abbas Ali Salih Al-Hamdani, Amal S. Sadiq, Tamador A. Mahmood. "Preparation and Spectroscopic Studies of New Transition Metal Complexes of 1-(2, 2-Dicyclohexylethyl)-3-(1, 5-Dimethyl-3-Oxo-2-Phenyl-2,3- Dihydro-1H-pyrazol-4-ylimino) Indolin-2-One", Journal of Al-Nahrain University-Science, 2015

Publication

<1 %

27

Dhanaraj, Chellaian Justin, and Jijo Johnson. "Studies on some metal complexes of quinoxaline based unsymmetric ligand: Synthesis, spectral characterization, in vitro biological and molecular modeling studies", Journal of Photochemistry and Photobiology B Biology, 2016.

Publication

<1 %

28

M.M. Omar, Hanan F. Abd El-Halim, Eman A.M. Khalil. "Synthesis, characterization, and biological and anticancer studies of mixed ligand complexes with Schiff base and 2,2'-bipyridine", Applied Organometallic Chemistry, 2017

Publication

&lt;1 %

29

[chemrj.org](http://chemrj.org)

Internet Source

&lt;1 %

30

[coek.info](http://coek.info)

Internet Source

&lt;1 %

31

[nanobioletters.com](http://nanobioletters.com)

Internet Source

&lt;1 %

32

[worldwidescience.org](http://worldwidescience.org)

Internet Source

&lt;1 %

33

[www.abap.co.in](http://www.abap.co.in)

Internet Source

&lt;1 %

34

[www.sdiarticle2.org](http://www.sdiarticle2.org)

Internet Source

&lt;1 %

35

Basma A. Ismail, Doaa A. Nassar, Zeinab H. Abd El-Wahab, Omya A.M. Ali. "Synthesis, characterization, thermal, DFT computational studies and anticancer activity of furfural-type schiff base complexes", Journal of Molecular Structure, 2021

Publication

&lt;1 %

36

Nurin Sakinatul Hayati Haji Damit, Malai Haniti Sheikh Abdul Hamid, Nur Sabrina Rahayu Haji Abdul Rahman, Siti Nor Hashimah Haji Ilias et al. "Synthesis, structural characterisation and antibacterial activities of lead(II) and some transition metal complexes derived from quinoline-2-carboxaldehyde 4-methyl-3-thiosemicarbazone", *Inorganica Chimica Acta*, 2021

Publication

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On