

11.1%

Date: 2022-07-26 04:50 UTC

\* All sources 100 | Internet sources 35 | Plagiarism Prevention Pool 3


<input checked="" type="checkbox"/>	[1]	<a href="http://www.iraqi-datepalms.net/assets/uploads/2018/11/RPW-Book-Rabab-El-Mergawi-1.pdf">www.iraqi-datepalms.net/assets/uploads/2018/11/RPW-Book-Rabab-El-Mergawi-1.pdf</a>	8.2%	89 matches
<input checked="" type="checkbox"/>	[2]	<a href="http://www.theses.mu.tn/fr/dynamique/uploads/669e7fe20f7624ec3f2b28be90235ec6.pdf">www.theses.mu.tn/fr/dynamique/uploads/669e7fe20f7624ec3f2b28be90235ec6.pdf</a>	5.4%	56 matches
<input checked="" type="checkbox"/>	[3]	<a href="http://www.intechopen.com/chapters/73021">www.intechopen.com/chapters/73021</a>	4.1%	58 matches
<input checked="" type="checkbox"/>	[8]	<a href="http://www.ncbi.nlm.nih.gov/pmc/articles/PMC8301634/">www.ncbi.nlm.nih.gov/pmc/articles/PMC8301634/</a>	2.6%	57 matches
<input checked="" type="checkbox"/>	[12]	<a href="http://www.cabi.org/fisc/datasheet/47472">www.cabi.org/fisc/datasheet/47472</a>	4.2%	52 matches
<input checked="" type="checkbox"/>	[14]	<a href="http://researcherslinks.com/current-issues/Effect-of-Sub-Lethal-Doses-of-Beauveria-bassiana/20/1/2032/html">researcherslinks.com/current-issues/Effect-of-Sub-Lethal-Doses-of-Beauveria-bassiana/20/1/2032/html</a>	3.4%	51 matches
<input checked="" type="checkbox"/>	[15]	<a href="http://www.ncbi.nlm.nih.gov/pmc/articles/PMC8301634/">www.ncbi.nlm.nih.gov/pmc/articles/PMC8301634/</a>	2.5%	45 matches 1 documents with identical matches
<input checked="" type="checkbox"/>	[22]	<a href="http://scialert.net/fulltext/?doi=jbs.2015.56.67">scialert.net/fulltext/?doi=jbs.2015.56.67</a>	3.6%	38 matches
<input checked="" type="checkbox"/>	[23]	<a href="http://gd.eppo.int/taxon/RHYCFE/download/datasheet_pdf">gd.eppo.int/taxon/RHYCFE/download/datasheet_pdf</a>	3.2%	40 matches 1 documents with identical matches
<input checked="" type="checkbox"/>	[25]	<a href="http://bioone.org/journals/florida-entomologist/volume-105/issue-1/024.105.0109/Effects-of-Trip-Locations-Pheromone-Source-and-Temperature-on-Red/10.1653/024.105.0109.full">bioone.org/journals/florida-entomologist/volume-105/issue-1/024.105.0109/Effects-of-Trip-Locations-Pheromone-Source-and-Temperature-on-Red/10.1653/024.105.0109.full</a>	3.1%	39 matches
<input checked="" type="checkbox"/>	[26]	<a href="http://journals.plos.org/plosone/article/file?type=printable&amp;id=10.1371/journal.pone.0255029">journals.plos.org/plosone/article/file?type=printable&amp;id=10.1371/journal.pone.0255029</a>	1.8%	41 matches
<input checked="" type="checkbox"/>	[28]	<a href="http://link.springer.com/article/10.1007/s42690-020-00148-1">link.springer.com/article/10.1007/s42690-020-00148-1</a>	2.9%	36 matches
<input checked="" type="checkbox"/>	[31]	<a href="http://ejar.journals.ekb.eg/article_68566_4b895ee522e26917df7bc712d9f63993.pdf">ejar.journals.ekb.eg/article_68566_4b895ee522e26917df7bc712d9f63993.pdf</a>	1.6%	43 matches
<input checked="" type="checkbox"/>	[32]	<a href="http://researcherslinks.com/base/downloads.php?jid=20&amp;aid=4626&amp;acid=8&amp;path=pdf&amp;file=1643183428PJZ_54_2_927-936.pdf">researcherslinks.com/base/downloads.php?jid=20&amp;aid=4626&amp;acid=8&amp;path=pdf&amp;file=1643183428PJZ_54_2_927-936.pdf</a>	1.9%	33 matches 1 documents with identical matches
<input checked="" type="checkbox"/>	[38]	<a href="http://bmcmicrobiol.biomedcentral.com/articles/10.1186/1471-2180-14-136">bmcmicrobiol.biomedcentral.com/articles/10.1186/1471-2180-14-136</a>	1.3%	29 matches
<input checked="" type="checkbox"/>	[39]	<a href="http://www.ncbi.nlm.nih.gov/pmc/articles/PMC7853973/">www.ncbi.nlm.nih.gov/pmc/articles/PMC7853973/</a>	1.0%	32 matches
<input checked="" type="checkbox"/>	[40]	<a href="http://link.springer.com/article/10.1017/S1742758418000048">link.springer.com/article/10.1017/S1742758418000048</a>	1.2%	28 matches
<input checked="" type="checkbox"/>	[41]	<a href="http://www.academia.edu/6044530/Portable_acoustic_device_for_detection_of_coconut_palms_infested_by_Rhynchophorus_ferrugineus_Coleoptera_Curculionidae">www.academia.edu/6044530/Portable_acoustic_device_for_detection_of_coconut_palms_infested_by_Rhynchophorus_ferrugineus_Coleoptera_Curculionidae</a>	2.0%	24 matches
<input checked="" type="checkbox"/>	[42]	<a href="http://bioone.org/journals/florida-entomologist/volume-100/issue-2/024.100.0220/The-Invasion-of-Tunisia-by-Rhynchophorus-ferrugineus-Coleoptera--Curculionidae/10.1653/024.100.0220.full">bioone.org/journals/florida-entomologist/volume-100/issue-2/024.100.0220/The-Invasion-of-Tunisia-by-Rhynchophorus-ferrugineus-Coleoptera--Curculionidae/10.1653/024.100.0220.full</a>	1.0%	29 matches
<input checked="" type="checkbox"/>	[43]	<a href="http://www.frontiersin.org/articles/10.3389/fevo.2022.826808/full">www.frontiersin.org/articles/10.3389/fevo.2022.826808/full</a>	0.6%	30 matches
<input checked="" type="checkbox"/>	[45]	<a href="http://en.wikipedia.org/wiki/Rhynchophorus_ferrugineus">en.wikipedia.org/wiki/Rhynchophorus_ferrugineus</a>	0.9%	29 matches
<input checked="" type="checkbox"/>	[46]	<a href="http://link.springer.com/article/10.1007/BF02980931">link.springer.com/article/10.1007/BF02980931</a>	1.3%	24 matches
<input checked="" type="checkbox"/>	[47]	<a href="http://www.nature.com/articles/srep31659">www.nature.com/articles/srep31659</a>	0.5%	27 matches
<input checked="" type="checkbox"/>	[54]	<a href="http://pubmed.ncbi.nlm.nih.gov/20337561/">pubmed.ncbi.nlm.nih.gov/20337561/</a>	0.2%	23 matches
<input checked="" type="checkbox"/>	[57]	<a href="http://www.mapa.gob.es/ministerio/pags/Biblioteca/Revistas/pdf_Plagas/BSVP_32_04_2_631_641.pdf">www.mapa.gob.es/ministerio/pags/Biblioteca/Revistas/pdf_Plagas/BSVP_32_04_2_631_641.pdf</a>	0.5%	19 matches
<input checked="" type="checkbox"/>	[59]	<a href="http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3649424/">www.ncbi.nlm.nih.gov/pmc/articles/PMC3649424/</a>	0.8%	19 matches
<input checked="" type="checkbox"/>	[62]	<a href="http://www.sciencedirect.com/science/article/abs/pii/S0261219409002257">www.sciencedirect.com/science/article/abs/pii/S0261219409002257</a>	0.5%	23 matches
<input checked="" type="checkbox"/>	[63]	<a href="http://link.springer.com/article/10.1017/S1742758411000099">link.springer.com/article/10.1017/S1742758411000099</a>	0.3%	18 matches
<input checked="" type="checkbox"/>	[64]	<a href="http://www.mdpi.com/2075-4450/13/3/245">www.mdpi.com/2075-4450/13/3/245</a>	0.2%	14 matches
<input checked="" type="checkbox"/>	[65]	<a href="http://www.tandfonline.com/doi/abs/10.1080/03235400601121521">www.tandfonline.com/doi/abs/10.1080/03235400601121521</a>	0.4%	12 matches
<input checked="" type="checkbox"/>	[72]	<a href="http://journals.plos.org/plosone/article/authors?id=10.1371/journal.pone.0251278">journals.plos.org/plosone/article/authors?id=10.1371/journal.pone.0251278</a>	0.7%	8 matches
<input checked="" type="checkbox"/>	[79]	from a PlagScan document dated 2022-01-27 09:49	0.6%	8 matches
<input checked="" type="checkbox"/>	[81]	from a PlagScan document dated 2018-07-27 13:17	0.0%	12 matches

[88]  getd.libs.uga.edu/pdfs/boozar\_whitney\_e\_201108\_ms.pdf  
0.2% 8 matches


---

[90]  journals.plos.org/plosone/article/authors?id=10.1371/journal.pone.0245928  
0.7% 5 matches


---

[92]  www.mdpi.com/1424-8220/20/20/5918/htm  
0.2% 6 matches

---

[95]  pubmed.ncbi.nlm.nih.gov/28364201/  
0.4% 4 matches

---

[104]  from a PlagScan document dated 2021-12-09 13:03  
0.0% 8 matches

---

17 pages, 5302 words

**PlagLevel: 11.1% selected / 76.1% overall**

329 matches from 106 sources, of which 40 are online sources.

#### Settings

Data policy: *Compare with web sources, Check against organization repository, Check against the Plagiarism Prevention Pool*

Sensitivity: *High*

Bibliography: *Bibliography excluded*

Citation detection: *Highlighting only*

Whitelist: *--*

<sup>[14]</sup>1 Semi-artificial diet developed for the successful rearing of red palm weevil:

2 *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) in the laboratory

3 Abdulrahman S. Aldawood<sup>a\*</sup>, Khawaja G. Rasool<sup>2</sup>, Sukirno Sukirno<sup>b</sup>, Mureed Husain<sup>3</sup>, Koko D.

4 Sutanto<sup>3</sup>, Mona Abdullah Alduailej<sup>c</sup>

5

6<sup>a</sup> Department of Plant Protection, College of Food and Agriculture Sciences, King Saud University,

7 Riyadh, Saudi Arabia, P.O. Box 2460, Riyadh 11451

8

9<sup>b</sup> Entomology Laboratory, Faculty of Biology, Universitas Gadjah Mada, Yogyakarta 55281,

10 Indonesia

11<sup>c</sup> College of Computer and Information Sciences, Princess Nourah Bint Abdulrahman University,

12 Riyadh, Saudi Arabia

13

14\* Corresponding author: Abdulrahman S. Aldawood

15 Department of Plant Protection, College of Food and Agriculture Sciences,

16 King Saud University, Riyadh, Saudi Arabia, P.O. Box 2460, Riyadh 11451

17 e-mail=aldawood@ksu.edu.sa

18 Tel: <sup>[1]</sup>+966-114-678426

19 Abstract

20 Objectives

21 The red palm weevil (RPW), *Rhynchophorus ferrugineus* (Coleoptera: Dryophthoridae) is the most  
22 destructive insect pest of several palm species, worldwide. The maintaining of RPW laboratory  
23 colonies for experimentation has been very challenging. We synthesized a standardized semi-artificial  
24 diet which is easy to prepare and economical for the rearing of RPW.

25 <sup>[25]</sup> Methods

26 The effects of semi-artificial diet on biological traits related to RPW development were studied at the  
27 Economic Entomology Research Unit (EERU), Department of Plant Protection, College of Food and  
28 Agriculture Sciences, King Saud University, Riyadh, Saudi Arabia. The diet was synthesized by  
29 combining shredded date palm petiole tissues, corn flour, wheat flour, ascorbic acid, sodium benzoic,  
30 sorbic acid, agar, and distilled water. <sup>[31]</sup> Three replicates, each containing fifty newly hatched larvae,  
31 were reared on the semi-artificial diet. The developmental time of RPWs fed on the newly synthesized  
32 diet was compared to the growth rate of conspecifics fed on date palm and previously published  
33 artificial diets.

34 Results

35 Fully-grown larval weight ranged from 4 – 6 g ( $5 \pm 0.19$  g), and larval duration ranged from 46 to 57  
36 days ( $53 \pm 0.3$  days). Average adult emergence was 58-98% ( $78 \pm 0.6$ ), with an average weight of 0.90  
37 g for males and 0.97 g for females. Complete life span was between 48 and 98 days ( $77 \pm 0.9$  days).  
38 On present semi artificial diet, several larvae had 11 larval instars. The diet has a shelf life of at least  
39 seven days and can last up to two weeks if properly kept and saved.

40 Conclusions

41 In conclusion, the present study proved that semi artificial diet is quite capable for the successful  
42 rearing of RPW. The growth and development of RPW larvae reared in this experiment were  
43 comparable to those fed on a natural diet.

44 Key words: Red palm weevil, *Rhynchophorus ferrugineus*, artificial diet, date palm.

45 Abbreviations

46 Red Palm Weevil (RPW), Economic Entomology Research Unit (EERU), completely randomized

47 design (CRD), Duncan's multiple range test (DMRT),

48 1.<sup>[1]</sup> Introduction

49 Red Palm Weevil (RPW) has been the most destructive insect pest of date palm in the Middle East

50 since the mid-1980's (El Ezaby et al., 1998, Ferry and Gomez, 2002). Recently, RPW has become an

51 invasive insect pest in several countries around the globe (Murphy and Briscoe, 1999; Malumphy and

52 Moran 2007, Bozbuga and Hazir, 2008). Due to its economic importance, many studies have

53 attempted to construct control strategies using biological control agents (Llacer et al., 2008; Dembilio

54 et al., 2010; Güerri-Agulló et al., 2011), baited trap systems (Abbas et al., 2006, Al-Saoud et al., 2010,

55 Faleiro et al., 2011), sterile male techniques (Al-Ayedh and Rasool, 2010), microwave heating (Massa

56 et al., 2011), chemical insecticides (Abraham et al., 1975), and integrated pest management (Tapia et

57 al., 2011). Although many techniques have been proposed to control the pest, none are satisfactory

58 due to its unique biological behavior, which includes the concealed nature RPW. Larvae, pupae and

59 even adults remains inside the tree trunk and larvae feed on the soft tissues.

60 Several techniques for the detection of RPW infestation on palms have been established, such as using

61 acoustic sensors (Gutiérrez et al., 2010, Siriwardena et al., 2010), sniffing dogs (Nakash et al., 1999),

62 and X-ray CT (computed tomography) scans (Ma et al., 2012), but they are not sufficiently accurate.

63 Determining RPW presence in tree trunks usually encounters difficulties that affect the accuracy of

64 results.<sup>[1]</sup> From above mentioned evidences regarding huge numbers of experiments, it is obvious that,

65 mass rearing of the RPW is very important to provide RPW different developmental stages to be used

66 in experimentation/bioassays. In lab-based biological assays, the insects used must be homogeneous

67 and available in large numbers.<sup>[14]</sup> In this regard, the present study objective was to develop a cost

68 effective and nutritious, artificial diet with enhanced shelf life for mass rearing of the RPW for the

69 successful experimentation.

70 Artificial diets should contain enough nutrition for the growth and development of the insects, and

71 have both physical and chemical properties that are palatable, digestible and absorbable by the insect

72 (Coudron et al., 2004). Attempts to mass rear RPWs have been reported.<sup>[3]</sup> Rearing of the weevil can be

73 achieved using date palms trunk (Al-Ayedh, 2008, Aldawood and Rasool, 2011), or sugarcane as host  
74 plants (Prabhu and Patil, 2009). Additionally, vegetables-based diets (Alfazairy, 2011) and date palm  
75 frond semi-synthetic diets (Al-Ayedh, 2011) can be used. Using natural diets increases the risk of  
76 uncontrolled contamination and has a larger space requirement.<sup>(1)</sup> On the other hand, the use of artificial  
77 diets is more practical, as the risk of contamination can be controlled and less space is needed  
78 (Situmorang, 1997). This study aimed to develop an artificial diet with a longer shelf life that satisfies  
79 the nutritional requirements of RPW larval stages, and evaluate this through its effects on various  
80 RPW development stages.

812.<sup>[25]</sup> Materials and Methods

82The experiment was conducted at the economic entomology research unit (EERU) Plant Protection  
83Department, College of Food and Agriculture Sciences, King Saud University. The parent generation  
84of RPWs was collected from date palm fields in the Riyadh Region (24.4164°N, 46.5765°E) and  
85comprised both adults and larvae. Adults were kept in plastic boxes (size, 1 Kg) and were provided  
86with cotton saturated with 10% (w/v) bee pollen solution. The laid eggs were collected and kept in  
87petri dishes (d: 8 cm; h: 2.5 cm) containing wet filter paper. Newly hatched larvae were then  
88maintained on the artificial diet. Therefore, in the present experiments, second-generation individuals  
89were reared on the present semi artificial diet.

90<sup>[8]</sup>The present semi artificial diet synthesized was modified from Al-Ayedh (2011) by adding  
91preservatives, antioxidants, and agar in order to improve the texture and to enhance the shelf life of  
92the diet. The artificial diet was mainly composed of shredded date palm petioles. To prepare the  
93ground petioles, date palm petioles were cut into pieces (3 cm × 3.5 cm) and air dried for three days to  
94reach 35 – 40% water content. Then, 500 g of ground petiole, 250 g wheat flour, 250 g corn flour, and  
95two g ascorbic acid were homogenized (mix 1.). 1.6 g potassium benzoate, 1.6 g sorbic acid, 20 g  
96agar, and two liters distilled water were boiled, then cooled until reaching 60 °C. After this, mix. 1  
97was homogenized with agar solution.

98In this experiment, three replicates with 50 larvae per replicate were used. Newly hatched larvae were  
99maintained in a plastic cup (d: 5 cm; h: 3 cm) and provided with 60 g of the diet until the 5<sup>th</sup> instar.  
100After 5<sup>th</sup> instar, they were transferred into a plastic jar (d: 6 cm; h: 8 cm) with 250 g of the diet until  
101full grown. During the last instar, the larvae starts wandering to find fiber for pupation, these full  
102grown wandering larvae were transferred into a box (l: 17 cm; w: 11 cm; h: 7 cm) and provided with  
10310 cm long sugar cane piece/stick for pupation. Emerged adults had a 1:1 sex ratio. Then, laid eggs  
104were observed to measure fecundity. Diets were changed every seven to 10 days. <sup>[1]</sup>As a control, 10  
105larvae maintained on the previously published artificial diet and 10 larvae reared on date palm bolts  
106(one-meter-long date palm trunk) were used, with three replicates for each. All colonies were  
107maintained in a growth chamber set at 70% RH and 25 °C temperature.

108 Larval growth and development were monitored daily by recording the exuviae or head capsules,  
109 while weight was measured two days after molting. Measured parameters were: larval instar  
110 development, larval weight, adult weight, fecundity, and egg hatchability. These parameters are  
111 common in published papers within the field, and within the literature.

#### 112 2.1. Experimental Design and Analysis

113 A completely randomized design (CRD) was used in this experiment. Analysis of variance ( $\alpha = 0.05$ )  
114 was conducted to compare the developmental stages of RPWs reared on the diets, and this was  
115 followed by Duncan's multiple range test (DMRT) to separate means. Statistical analyzes were  
116 completed using SPSS version 13.0 (SPSS Inc 2005).



### 1173. Results

#### 1183.1. Larval development

119 On newly synthesized semi artificial diet, RPW larvae completed ten to eleven instars (Table 1).

120 Results indicated significant differences ( $F = 48.9$ ;  $df = 2, 74$ ;  $P = 0.001$ ) in larval weight at the full-  
121 grown stage. Larval weights developed from 0.001 g for the 1<sup>st</sup> instar, to an average of 5.5 g (range: 4  
122- 7 g) for the tenth instar, to an average of 5 g (range: 4 to 6 g) for the eleventh instar. The total RPW  
123 larval duration was 53 days (range: 46-57) days. Mortality during larval stage was nil. Only 7% larvae  
124 completed eleven instars while 93% larvae pupated after 10<sup>th</sup> instar (Table 1).

#### 1253.2. Pupal development

126 After larvae were fully grown and stopped feeding, they were transferred onto sugar cane for  
127 pupation. Mortality during the pupation stage was very low, at roughly 2%. The pupal period varied  
128 from 10-49 days, with an average of 25 days (Table 2).

#### 1293.3. Adult development

130 Red palm weevil larval stages fed on newly synthesized semi artificial diet successfully developed  
131 into adults. An average of 78% successfully emerged, with a range of 58 to 98%. More males emerged  
132 than females, resulting in a sex ratio of 1.00 males to 0.79 females (Table 2).

133 RPW adults provided with pollen solution varied in longevity. Longevity of males ranged from 33 to  
134 72 days, while for females the range was 15 to 72 days. There was a difference in adult longevity  
135 between females and males, with males having shorter life spans than females. On average, males  
136 survived 49 days post-emergence, while females survived 52 days on average.

137 Adult male and female weights ranged from 0.6 to 1.21 g, and 0.7 to 1.24 g respectively, with an  
138 average of 0.90 g and 0.97 g respectively. Additionally, we observed that females could lay between  
139 48 and 193 eggs, with an average of 96 eggs, in a 35 to 61-day period (Table 2).

140 The egg laying period for RPWs fed on newly synthesized semi artificial diet ranged from 35 to 61  
141 days, with an average of 48 days. Total development time for RPWs from egg to full size ranged from  
142 48 to 98 days, with an average of 77 days on present semi artificial diet (Table 2).

### 143 Discussion

144 In the present research, we studied RPW larval growth and development on our newly synthesized  
145 semi artificial diet and compared its results with other diets claimed to be suitable for red palm weevil  
146 rearing. The diet succeeded in supporting RPW larval growth and development.

147 RPW larvae completed ten to eleven instars on newly synthesized semi artificial diet. Around 93 % of  
148 the population fed on the newly synthesized semi artificial diet had ten instars, while the remaining  
149 had eleven instars. This result was similar to that observed when larvae are fed on chopped date palm  
150 fronds (Al-Ayedh, 2011). In our study, most larvae had a tenth instar for 4-13 days (mean = 8.9 days),  
151 and about 7% of the population needed an additional five days to develop an eleventh instar.

152 Larval weights developed from 0.001 g for the 1<sup>st</sup> instar, to an average of 5.5 g (range: 4-7 g) for the  
153 tenth instar and an average of 5 g (range: 4-6 g) for the eleventh instar. In comparison to larvae fed on  
154 sukary date palm (Al-Ayedh, 2008), for which larval weight averaged 2.73 g, larvae fed on this  
155 artificial diet were heavier. There were significant differences in larval weight at the full-grown stage.  
156 If we compare it with other diets, larval weight on date palm bolt was significantly lower than on  
157 other diets. On the Al-Ayedh, (2011) diet, larval instar development was faster compared to other  
158 treatments until the 6<sup>th</sup> instar, but was slower than other diets in the advanced stages of larval  
159 development.

160 RPW average larval duration was 53 days (range: 46-57) days. RPW larval duration was shorter than  
161 observed by Al-Ayedh (2011), Alsuhaibani et al. (2001), Nirula (1956), and Prabhu and Patil (2009),  
162 who reported 66, 182, 55, and 56 days, respectively, but longer than observed by Ghosh (1912),  
163 Ghosh (1912) Watanapongsiri (1966), and Abd El-Fattah et al. (2009), which reported 35, 35 to 38,  
164 and 45 days, respectively.

165 The use of an artificial diet for RPW rearing during larval stages is more suitable for developing  
166 insects for use in experimentation. Larval growth and development can be observed more easily than  
167 in larvae fed on a natural diet. This is because the texture of the diet makes collection of the desired  
168 larval stage easier, reducing mishandling effects. Additionally, less space is required for rearing. It is  
169 easy to control contamination by adding preservatives and anti-fungal agents. The percentage of  
170 RPWs reaching full-grown development ranged from 58 to 98%, with an average of 78%.

171The pupal period varied from 10-49 days, with an average of 25 days. When compared to  
172observations made by Prabhu and Patil (2009), Nirula (1956) and Alsuhaibani et al. (2001), in which  
173pupal duration averaged 16, 16, and 23 days respectively, pupal duration in this study was shorter.  
174Abd El-Fattah et al. (2009) developed an artificial diet, and larvae fed on it required 22.64 days to  
175develop into adults from the pre-pupae stage.  
176Through continuously rearing RPWs on the diet, it was found that rough chopped petioles could be  
177used as a medium for cocoon spinning, as they provide fibers for cocooning. Fresh rough ground date  
178palm petioles should be combined with 1% ascorbic acid solution (w/v) as an antioxidant to overcome  
179oxidation and provide humidity for the fibers.  
180Red palm weevil larval stages fed on present semi artificial diet were successfully developed into  
181adults. An average of 78% successfully emerged, with a range of 58 to 98%. On a diet of sukary date  
182palm trunk, almost 72% of the population developed into adults (Al-Ayedh, 2008), while on  
183sugarcane this figure rose to 87% (Prabhu and Patil, 2009) (Table 1). More males emerged than  
184females, resulting in a sex ratio of 1.00 males to 0.79 females. Al-Ayedh (2008) and Nirula (1956)  
185also observed the same phenomenon, with ratios of 0.51 : 0.49 and 1.3 : 1, respectively.  
186RPW adults provided with pollen solution varied in longevity. Longevity of males ranged from 33 to  
18772 days, while for females the range was 15 to 72 days. There was a difference in adult longevity  
188between females and males, with males having shorter life spans than females. On average, males  
189survived 49 days post-emergence, while females survived 52 days on average. These life span  
190averages were shorter than for larvae fed on sugar cane (Prabhu and Patil, 2009), date palm  
191(Alsuhaibani et al. 2001), artificial diet (Abd El-Fattah et al. 2009), and coconut (Nirula, 1956). Male  
192life spans for these diets averaged 71, 161, 98 and 84 days, respectively, whilst females survived for  
193an average of 69, 112, 95 and 60 days, respectively. Deprived of food and humidity, the weevil can  
194survive for no more than six days, while if only humidity is provided it may survive for three to six  
195weeks (Alsuhaibani et al. 2001). When fed on date palm bolts, females survived up to 172 days, while  
196males were capable of surviving for up to 176 days (Al-Ayedh, 2008).  
197There were significant effects of diet on adult performance. Male weevils' weight differed  
198significantly between diets, with males fed on date palm bolt being lighter than those reared on

199artificial diets. On the contrary, weight of females reared on the present semi artificial diet was lighter  
200than on the Al-Ayedh (2011) diet. The present semi artificial diet improved fecundity, which was  
201significantly higher than on the Al-Ayedh (2011) diet and date palm bolt treatments, with higher  
202hatchability percentage.

203Adult male and female weights ranged from 0.6 to 1.21 g, and 0.7 to 1.24 g respectively, with an  
204average of 0.90 g and 0.97 g respectively. <sup>[1]</sup>Prabhu and Patil (2009) observed that RPW weight ranged  
205from 0.53 to 1.09 g and 0.57 to 1.43 g for males and females respectively when fed on sugarcane. We  
206also observed that females could lay between 48 and 193 eggs, with an average of 96 eggs, in a 35 to  
20761-day period. This result was similar to Faghih (1996), which recorded that female RPWs could lay  
208between 127 and 276 eggs. On sugar cane, adult females laid 278 eggs. Each female laid up to 412  
209eggs when reared on date palm trunk. The egg laying period for females living on this tree ranges  
210from 54 to 137 days (Alsuhaibani et al., 2001). On coconut trees, RPW females lay eggs for between  
21125 and 63 days (Nirula, 1956). Weevils fed on an artificial diet by Abd El-Fattah et al. (2009) and  
212Salama and Abdel-Razek, (2002), produced 275 and 184 eggs, respectively. Fertile laid eggs required  
213between 2-6 days (mean: 4.5 days) to hatch. This range was similar to those observed by Ghosh  
214(1912), Nirula (1956), Watanapongsiri (1966), and Faghih (1996), Alsuhaibani et al. (2001), and  
215Prabhu and Patil (2009).

216The oviposition period for RPWs fed on present semi artificial diet ranged from 35 to 61 days, with an  
217average of 48 days. This duration was shorter than in those fed on date palm. On date palm trees, egg  
218laying duration varied from 54 to 137 days, and the average was 96 days (Alsuhaibani et al. 2001).  
219RPWs fed on coconut laid eggs for between 25 and 63 days (Nirula, 1956). Total development time  
220for RPWs from egg to full size ranged from 48 to 98 days, with an average of 77 days on present semi  
221artificial diet. This complete life cycle period was shorter than for those fed on sugar cane (Prabhu  
222and Patil, 2009), on date palm (Alsuhaibani et al., 2001), and on coconut (Nirula, 1956); average life  
223cycle periods on those diets were found to be 82, 216, and 81 days, respectively.

#### 2245. Conclusions

225In conclusion, the present semi artificial diet was successful in supporting growth and development  
226during RPW mass rearing in the laboratory. The growth and development of RPW larvae reared in this

227 experiment were comparable to those fed on a natural diet. This study also observed that roughly  
228 grinded date palm fronds provide satisfactory results for pupation with no need for the addition of  
229 sugarcane during laboratory rearing.

#### 230 Acknowledgments

231 The authors would like to thank the Researchers Supporting Project, King Saud University, Riyadh,  
232 Saudi Arabia for financial support of the present work through project number RSP-2021/107.

233 Disclosure of funding

234 The authors declare that they don't have any particular funding for this study.

235 Disclosure of any conflict of interest

236 The authors declare that they don't have any type of conflict of interest which might affect the  
237 present work.

238References

- 239Abbas, M.S.T., Hanounik, S.B., Shahdad, A.S., Al-Bagham, S.A., 2006. <sup>[1]</sup>Aggregation pheromone  
240 traps, a major component of IPM strategy for the red palm weevil, *Rhynchophorus*  
241 *ferrugineus* in date palms (Coleoptera: Curculionidae). *J. Pest Sci.* 79, 69-73. DOI:  
242 10.1007/s10340-005-0113-6.
- 243Abd El-Fattah, S.M., El-Zohairy, M.M., Kelany, E.M., El-Dieeb, M.A., 2009. <sup>[1]</sup>Biological studies on  
244 the red palm Weevil, *Rhynchophorus ferrugineus*, Olivier, Coleoptera: Curculionidae reared  
245 on natural and artificial diets. *Zagazig J. Agri. Res.* 36, 1297-1314.
- 246Abraham, V.A., Abdulla, K.M., Kurian, C.H., 1975. <sup>[1]</sup>Evaluation of seven insecticides for control of red  
247 palm weevil *Rhynchophorus ferrugineus* Fabr. *J. Plants Crops.* 3, 71-72.
- 248Al-Ayedh, H., 2008. <sup>[1]</sup>Evaluation of date palm cultivars for rearing the red date palm weevil,  
249 *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae). *Fla. Entomol.* 91, 353-358. DOI:  
250 10.1017/S1742758411000063.
- 251Al-Ayedh, H., 2011. <sup>[1]</sup>Evaluating a semi-synthetic diet for rearing the red palm weevil *Rhynchophorus*  
252 *ferrugineus* (Coleoptera: Curculionidae). *Int. J. of Trop Insect Sci.* 1, 1-9.
- 253Al-Ayedh, H.Y., Rasool, K.G., 2010. <sup>[1]</sup>Determination of the optimum sterilizing radiation dose for  
254 control of the red date palm weevil *Rhynchophorus ferrugineus* Oliv. (Coleoptera:  
255 Curculionidae). *Crop Prot.* 29, 1377-1380. DOI: <https://doi.org/10.1016/j.cropro.2010.06.019>.
- 256Al-Saoud, A., Al-Deeb, M., Murchie, A., 2010. <sup>[1]</sup>Effect of color on the trapping effectiveness of red  
257 palm weevil pheromone traps. *J. Entomol.* 7, 54-59. DOI: 10.3923/je.2010.54.59.
- 258Aldawood, A., Rasool, K., 2011. Rearing optimization of red palm weevil: *Rhynchophorus*  
259 *ferrugineus* (Coleoptera: Curculionidae) on date palm: *Phoenix dactylifera*. *Fla. Entomol.* 94,  
260 756-760. DOI: <https://doi.org/10.1653/024.094.0404>.
- 261Alfazairy, A.A., 2011. A simple feeding diet for evaluating the efficacy of microbial or chemical  
262 control agents against larval stage of curculionid red palm weevil, *Rhynchophorus*  
263 *ferrugineus* (Olivier.). *Egypt. J. Biol. Pest Co.* 21, 101.

- 264 Alsuhaibani, A.M., Aldafer, H.M., Alahmadi, A.Z., 2001.<sup>[1]</sup> Behavioral and biological studies on the red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera, Curculionidae) in Riyadh, Saudi Arabia. Res. Bult. Res. Cent. Coll. of Agri., King Saud Univ. 107, 5–30.
- 267 Bozbuga, R., Hazir, A., 2008. Pests of the palm (Palmae sp.)<sup>[28]</sup> and date palm (*Phoenix dactylifera*) determined in Turkey and evaluation of red palm weevil (*Rhynchophorus ferrugineus* Olivier) (Coleoptera: Curculionidae). EPPO Bull. 38, 127–130.
- 270 Coudron, T.A., Goodman, C. L., Jones, W.A., Leopold, R., 2004. Development of an artificial diet and evaluation of artificial ovipositional substrates for the in vitro rearing of *Gonatocerus* spp. parasitoids of the eggs of the glassy-winged sharpshooter. USDA ARS Bulletin. Columbia.
- 273 Dembilio, O., Quesada-Moraga, E., Santiago-Álvarez, C., Jacas, J.A., 2010.<sup>[1]</sup> Potential of an indigenous strain of the entomopathogenic fungus *Beauveria bassiana* as a biological control agent against the Red Palm Weevil, *Rhynchophorus ferrugineus*. J. Invertebr. Pathol. 104, 214–221. DOI: 10.1016/j.jip.2010.04.006.
- 277 El Ezaby, F.A., Khalifa, O., Assal, A.E., 1998.<sup>[1]</sup> Integrated pest management for the control of red palm weevil *Rhynchophorus ferrugineus* Oliv. in The United Arab Emirates, Eastern Region, Al Ain. In Rahman-Al Afifi, M. A. and Al-Sherif Al-Badawy, A. (Eds) Proceedings of the First International Conference on Date Palms. Al-Ain, UAE, 8–10 March 1998. Faculty of Agricultural Sciences, UAE University, pp. 269–281.
- 282 Faghih, A.A., 1996. The biology of red palm weevil, *Rhynchophorus ferrugineus* Oliv. (Coleoptera, Curculionidae) in Savaran region (Sistan province, Iran). Appl. Entomol. Phytopathol. 63, 16–86.
- 285 Faleiro, J., El-Saad, M.A., Al-Abbad, A.H., 2011. Pheromone trap density to mass trap *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae/Rhynchophoridae/ Dryophthoridae) in date plantations of Saudi Arabia. Int. J. Trop. Insect Sci. 31, 75–77. DOI: 10.1017/S1742758411000099.
- 289 Ferry, M., Gomez, S., 2002. The Red Palm Weevil in the Mediterranean Area. Palms 46, 172–178.

- 290 Ghosh, C.C., 1912. Life Histories of Indian Insects- III, The Rhinoceros Beetle (*Oryctes rhinoceros*)  
291 and the red or palm weevil (*Rhynchophorus ferrugineus*). Memoirs of the Dept. of Agric.  
292 India. (Ent. Ser.) II 10: 205–217.
- 293 Güerri-Agulló, B., López-Follana, R., Asensio, L., Barranco, P., Lopez-Llorca, L., 2011. <sup>[23]</sup> Use of a  
294 solid formulation of *Beauveria bassiana* for biocontrol of the red palm weevil  
295 (*Rhynchophorus ferrugineus*) (*Coleoptera*: Dryophthoridae) under field conditions in SE  
296 Spain. Fla. Entomol. 94, 737–747. DOI: 10.1653/024.094.0402.
- 297 Gutiérrez, A., Ruiz, V., Moltó, E., Tapia, G., Téllez, M.M., 2010. <sup>[12]</sup> Development of abioacoustic sensor  
298 for the early detection of Red Palm Weevil (*Rhynchophorus ferrugineus* Olivier). Crop Prot.  
299 29, 671–676. DOI: 10.1016/j.cropro.2010.02.001.
- 300 Llacer, E., 2008. <sup>[1]</sup> Prospect for the biological control of the red palm weevil, *Rhynchophorus*  
301 *ferrugineus* (*Coleoptera*: Curculionidae), in mediterranean. Proceedings of ISBCA 3- P.G.  
302 Mason, D. R. Gillespie and C. Vincent Eds. (2008).
- 303 Ma, A.K.W., Alghamdi, A.A., Tofailli, K., Spyrou, N.M., 2012. X-ray CT in the detection of palm  
304 weevils. J. Radioanal. Nucl. Ch. 291, 353–357. DOI: 10.1007/s10967-011-1202-z.
- 305 Malumphy, C., Moran, H., 2007. Plant Pest Notice: Red Palm Weevil, *Rhynchophorus ferrugineus*.  
306 <sup>[28]</sup> Department for Environment, Food and Rural Affairs, Central Science Laboratory.
- 307 Massa, R., Caprio, E., De Santis, M., Griffo, R., Migliore, M.D., Panariello, G., Pinchera, D., Spigno,  
308 P., 2011. Microwave treatment for pest control: <sup>[22]</sup> the case of *Rhynchophorus ferrugineus* in  
309 *Phoenix canariensis*. EPPO Bull. 41, 128–135. DOI: 10.1111/j.1365-2338.2011.02447.x.
- 310 Murphy, S., Briscoe, B., 1999. The red palm weevil as an alien invasive: <sup>[1]</sup> biology and the prospects for  
311 biological control as a component of IPM. Biocontrol News Inform. 20, 35–46.
- 312 Nakash, J., Osem, Y., Kehat, M., 1999. <sup>[1]</sup> A suggestion to use dogs for detecting red palm weevil  
313 (*Rhynchophorus ferrugineus*) infestation in date palms in Israel. Phytoparasitica. 28, 153–  
314 155. DOI: 10.1007/BF02981745.
- 315 Nirula, K.K., 1956. <sup>[22]</sup> Investigations on the pests of coconut palm, (*Rhynchophorus ferrugineus* F.).  
316 Indian Coconut J. 9, 229–247.



- 317 Prabhu, S.T., Patil, R.S., 2009.<sup>[1]</sup> **Studies on the biological aspects of red palm weevil, *Rhynchophorus***  
318 ***ferrugineus* (Oliv.)**. Karnataka J. Agric. Sci. 22, 732–733.
- 319 Salama, H.S., Abdel-Razek, A.S., 2002.<sup>[1]</sup> **Development of the red palm weevil, *Rhynchophorus***  
320 ***ferrugineus* (Olivier), (Coleoptera, Curculionidae) on natural and synthetic diets**. Anzeiger  
321 Schädlingsskd. 75, 137–139. DOI: 10.1046/j.1472-8206.2002.02039.x.
- 322 Siriwardena, K., Siriwardena, K.A.P., Fernando, L.C.P., Nanayakkara, N., Perera, K.F.G., Kumara,  
323 A.D.N.T., Nanayakkara, T., 2010.<sup>[2]</sup> **Portable acoustic device for detection of coconut palms**  
324 **infested by *Rhynchophorus ferrugineus* (Coleoptera: Curculionidae)**. Crop Prot. 29, 25–29.  
325 DOI: 10.1016/J.CROPRO.2009.09.002.
- 326 Situmorang, J., 1997. Optimasi susunan pakan buatan untuk pertumbuhan dan kualitas kokon ulat  
327 sutera atakas, *Attacus atlas* (Linn.) (Lepidoptera: Saturniidae). Research report Faculty of  
328 Biology Gadjah Mada University Indonesia.
- 329 Tapia, G., Ruiz, M., Téllez, M., 2011. Recommendations for a preventive strategy to control red palm  
330 weevil (*Rhynchophorus ferrugineus*, Olivier) based on the use of insecticides and  
331 entomopathogenic nematodes. EPPO Bull. 41, 136–141. DOI: [https://doi.org/10.1111/j.1365-](https://doi.org/10.1111/j.1365-2338.2011.02448.x)  
332 [2338.2011.02448.x](https://doi.org/10.1111/j.1365-2338.2011.02448.x).
- 333 Watanapongsiri, A., 1966. A revision of the genera *Rhynchophorus* and *Dynamis* (Coleoptera:  
334 *Curculionidae*). Department of Agriculture Science Bulletin Vol.1 No.1. Bangkok. Thailand.

335 Table 1. <sup>[1]</sup> Growth and development of red palm weevil larval stages fed on semi synthetic diet (Al-Ayedh, 2011), date palm, and present semi

336 artificial diet

Parameters	Semi-Artificial Diet			Date Palm			Al-Ayedh, 2011			F-test	p-value	
	N	Range	Mean ± SEM	N	Range	Mean ± SEM	N	Range	Mean ± SEM			
Instar duration (d):												
1 <sup>st</sup> instar	102	3 – 4	3.42±0.05 a	30	2–5	3.47±0.16 a	30	1–4	2.73±0.12 b	16.353	3.48E-07	P 0.0001
2 <sup>nd</sup> instar	102	3 – 6	4.41±0.05 a	30	2–5	3.23±0.13 b	30	2–5	3.30±0.16 b	57.675	2.20E-16	P 0.0001
3 <sup>rd</sup> instar	102	4 – 6	4.67±0.05 a	30	2–5	3.66±0.12 b	30	2–5	3.36±0.15 b	64.844	2.20E-16	P 0.0001
4 <sup>th</sup> instar	102	4 – 6	4.51±0.05 a	30	3–6	3.86±0.17 b	30	2–8	3.66±0.21 b	17.478	1.38E-07	P 0.0001
5 <sup>th</sup> instar	102	5 – 8	6.16±0.07 a	30	3–6	4.30±0.12 b	30	2–5	3.66±0.11 c	182.35	2.20E-16	P 0.0001
6 <sup>th</sup> instar	102	4 – 7	4.96±0.07 a	30	3–7	4.86±0.23 a	30	2–8	4.16±0.28 b	6.7058	0.0016	P 0.01
7 <sup>th</sup> instar	102	4 – 6	5.05±0.06 b	30	3–8	4.60±0.21 c	30	3–8	5.53 ±0.22 a	7.8524	0.0005595	P 0.001
8 <sup>th</sup> instar	102	4 – 8	5.27±0.09 b	30	2–13	5.43±0.39 b	30	2–11	7.26±0.42 a	19.61	2.44E-08	P 0.0001
9 <sup>th</sup> instar	102	4 – 8	5.09±0.08 b	30	4–26	8.10±0.78 a	30	5–10	7.16±0.24 a	30.278	7.22E-12	P 0.0001
10 <sup>th</sup> instar	95	4 – 13	8.99±0.26 b	30	12–39	23.16±1.37 a	30	5–19	9.86±0.69 b	137.24	2.20E-16	P 0.0001
11 <sup>th</sup> instar	7	0-5	0.49±0.14 b	-	-	-	30	10–28	17.86±0.94 a	591.83	2.20E-16	P 0.0001
Larval duration (d)	102	46-57	53.05±0.27 c	30	48–89	64.7±1.95 b	30	47-83	68.6±1.43 a	97.085	2.20E-16	P 0.0001

337

338

31  
32

16

Table 2. Comparison of Red palm weevil performance on different diets

Parameters	Nirula (1956)	Alsuhaibani et al. (2001)	Prabhu and Patil (2009)	Abd El -Fattah et al. (2009)			Al-Ayedh (2011)		Salama and Abdel-Razek (2002)	Aldawood et al. semi-artificial diet
	Coconut	Date palm	Sugarcane	Sugarcane	Date palm	Artificial diet	Artificial diet	Date palm		
<b>Egg</b>										
Fecundity	76-355 (204)	55-412 (227)	211-380 (278)	275±24	259±26	222±22.74	35.2 ±7	32 ± 8	184±19	48-193 (96 ) ±15
Oviposition period	25-63 (45)	54-137 (96)	-	-	-	-	-	-	-	35-61 (48) ± 3
Hatchability %	-	-	70-90 (83)	84±5	87±5	829±5	21-71 (40) ± 5	0-61 (30) ± 5	94-100	43-100 (73) ± 8
Incubation period	2-5 (3)	2-5 (4)	2-4 (3)	3.16±0.17	3.16±0.16	3.09±0.17	-	-	-	2-6 (4.5)
<b>Larva</b>										
# instars	-	-	8	-	-	-	11	10	-	10-11
Duration	35-71 (55)	120-241 (182)	32-65 (56)	68±4	37±2	45±3	45-81 (66) ± 1	48-89 (65) ± 2	45	46-57 (53) ± 0.3
Head Capsule width	-	-	-	-	-	-	5.16	5.69	-	-
Weight	-	-2.73	4-6.4 (4.95)	-	-	-	3-6 (5) ± 0.1	0.1-5 (2) ± 0.2	-	4-6 (5) ± 0.2
<b>Pupa</b>										
Duration	12-21 (16)	21-26 (23)	10-12 (16)	-	-	22-64	-	-	-	10-49 (25) ± 0.7
Weight	-	-	1-3 (2)	-	-	-	1-3 (2) ± 0.1	1-3 (2) ± 0.2	-	-
<b>Adult</b>										
Emergence %	-	-	85-90 (87)	-	-	-	-	-	-	58-98 (78) ±0.6
Male longevity	48-113 (84)	76-257 (161)	62-78 (71)	92.4±6.84	93.4±5.37	98±5.96	-	-	95.2	33-72 (49) ± 3
Female longevity	43-76 (60)	70-335 (112)	59-75 (69)	82±6	91 ± 6	95 ± 7	-	-	98	15-72 (52) ± 3
Sex ratio	1.3:1	0.8:1	1:1	0.53:0.47	-	-	0.51: 0.49	-	0.53 : 0.47	1 :0.79
Male weight	-	-	0.53-1 (0.9)	-	-	-	0.5-1.4 (1) ± 0	0.65-.96 (1) ± 0.1	-	0.6-1.21 (0.90) ± 0.1
Female weight	-	-	0.6-1.43 (1)	-	-	-	0.6-1.4(1) ± 0	0.6-1 (1) ± 0.1	-	0.7-1.24 (0.97) ±0.1
<b>Total Life cycle development</b>										
Egg to Adult	54-120 (81)	208-223 (216)	58-97 (82)	165-195	-	-	-	-	-	48-98 (77±0.9)

339

33

34