

Chemical characterization and Larvicidal Potential of Thuja orientalis Leaves and Fruits Extracts against Culex pipiens (Diptera: Culicidae)

By Hanan Bosly

Chemical characterization and Larvicidal Potential of *Thuja orientalis* Leaves and Fruits Extracts against *Culex pipiens* (Diptera: Culicidae)

ABSTRACT

Botanical pesticides targeted to avoid the pesticide resistance to synthetic ones that conventionally affecting ecosystems diversity. The study was designed to evaluate the larvicidal potentials of leaves and fruits extracts (methanol, acetone, hexane, and aqueous) from *Thuja orientalis* (Pinales: Cupressaceae) against *Culex pipiens* 3rd instar larvae and to identify the extracts compounds by gas chromatography-mass spectrometry (GC-MS) method of analysis. Leaves and fruits extracts were used in concentrations, 25, 50, 100, 200, and 400 ppm against the 3rd larval instar of *Cx. pipiens* in five replicates and larval mortalities recorded after 24 and 48 h post exposure. The larvicidal potentials of extracts showed concentration dependent and varied between leaves and fruits extracts. At 400 ppm concentration, leaves extracts showed 100% larval mortality except in hexane extract exhibited 98%. Acetone, methanol, aqueous and hexane leaves extracts recorded LC₅₀ values, 58.04, 70.20, 77.19 and 84.25 ppm, respectively. Fruits extracts by hexane and methanol exhibited 100% larval mortality and LC₅₀, 68.26 and 83.21 ppm, respectively, while, acetone and aqueous fruits extracts showed 98% and 96% larval mortality and LC₅₀, 92.81 and 102.97 ppm, respectively. Terpenoids and sesquiterpenoids, fatty acid esters mainly identified in both extracts. The present study showed that extracts from *T. orientalis* leaves and fruits acquired promising larvicidal potential for the control of *Cx. pipiens* larvae with the role of their chemical constituents.

Keywords: *Thuja orientalis*; leaves; Fruits; *Culex pipiens*; Larvicidal, GC-MS.

1. Introduction

Mosquitoes are considered a burden to human health by transmitting diseases including malaria, filariasis, dengue, and leishmaniasis (Wilson et al., 2020). *Culex pipiens* mosquitoes is common house mosquito and one of the most widely distributed mosquitoes worldwide due to its adaptation to human environments and its mode of feeding on birds and mammals reflected by its role in transmission of the West Nile virus and other pathogens (Farajollahi et al., 2011) that urged researches of its control.

Vector control is the main way to reduce public concerns about mosquito-borne diseases including filariasis, dengue, malaria, and leishmaniasis (Wilson et al., 2020). Control of mosquito larvae in aquatic phases is an effective method for reducing mosquito spread (WHO, 2013). Excessive use of synthetic insecticides, with a complete lack of awareness of the strategy of changing the pesticides, led to resistance to pesticides along with ¹⁴ environmental pollution and health risks to humans and non-target biota. Therefore, ¹⁴ the search for environmentally friendly alternatives as plants or oils rich in secondary metabolites is a recent trend since they are more efficient, less toxic, biodegradable, and capable of insect decrease plant resistance to these natural compounds (Mouden et al., 2017; Ahmed et al., 2021) besides serving as larvicides, adult pesticides, insect repellents and deterrents (Govindarajan et al., 2016) they ⁵ destroy only the insects they are meant to kill, leaving no residue on food or in the environment.

T. orientalis is a dense, evergreen and coniferous tree belongs to the family Cupressaceae growing in Saudi Arabia (Elsharkawy et al., 2017; Elsharkawy and Ali, 2019). In folk medicine *T. orientalis* used as herbal medicine for treatment of psoriasis, amenorrhea, enuresis, rheumatism, cystitis, bronchial catarrh and uterine carcinomas (Srivastava et al., 2012). Studies evaluated different activities of *T. orientalis* like; antimicrobial (Choi et al., 2021), antifungal activity (Caruntu et al., 2020), antioxidant (Nizam and Mushfiq, 2007), anticancer ³⁵ (Elsharkawy et al., 2017), anti-inflammatory (Darwish et al., 2021; Shin et al., 2015), hair growth promotion (Zhang et al., 2013). The aforementioned activities were proposed to the active constituents of *T. orientalis* that were mainly phenolic, flavonoid and terpenoid compounds extracted from areal parts, leaves and fruits by different methods of extraction as steam distillation for oil extraction or successive extractions by water, methanol and petroleum ether or simply by water and ethanol extract.

It is well known that the solvent extracting material is an important factor affecting the yield of the plant chemical constituents and the extract performance for the consequent larvicidal activity. The present study used methanol, hexane, acetone and water that they differ in polarity from the most polar was water followed by methanol followed the median polar acetone to the non-polar was hexane. The extractions were performed using direct soxhlet extraction by single solvent (Azwanida, 2015).

¹³ The present study was designed to investigate the larvicidal activities of the methanol, hexane, acetone and aqueous extracts from leaves and fruits of *T. orientalis* plant against *Cx. pipiens* third instar larvae after exposure to extracts for 24 and 48 h. As well, identification of the chemical composition of the tested extracts by the aid of ¹² gas chromatography-mass spectrometry (GC-MS) method of analysis.

2. Materials and Methods

2.1. Plants materials

Thuja orientalis L. (Pinales: Cupressaceae) leaves and fruits were obtained ³⁹ from Arar region, at Northern Region of Saudi Arabia, where growing wild in March 2022 as previously recorded (Elsharkawy and Ali, 2019). The plant was identified by a taxonomist, Prof. ⁷ Dr. Yahya Masrahi, from the Department of Biology, Faculty of Science, Jazan University, Saudi Arabia.

2.2. *Culex pipiens* colony

Mosquitoes (*Cx. pipiens*) were obtained from the Center for Environmental Research and Studies at Jazan University. Rearing was performed ² under controlled conditions (27 ± 2 °C, relative humidity at $70\% \pm 10\%$, and 12:12 h light:dark regime). Mosquito larvae were reared in round enamel plates (25 x 20 x 10 ²² cm) filled with 2 L de-chlorinated water and fed with fish food daily. The third instar *Cx. pipiens* larvae were used for the larvicidal examination.

2.3. Plant extracts

Leaves and fruits were dried in shade for 7 days at laboratory temperatures (27-29 °C). The dried leaves (40 g) and fruits (25 ¹¹ g) were powdered using a commercial electrical stainless-steel blender and extracted for each solvent including methanol, acetone, hexane, and aqueous (in ratio, 1/10 w/v sample/solvent) using Soxhlet apparatus for 6-8 h according to the solvent type. The leaves and fruits extracts were filtered with ³⁷ Whatman number 1 filter paper through a Buchner funnel. The filtrates then ³⁴ dried using a rotary evaporator under vacuum at 40 °C. The plant leaves extract yields 3.7, 3.2, 2.8, and 1.7 g for methanol, acetone, hexane, and aqueous solvents and yields 2.1, 1.5, 2.3 and 1.1 g for plant fruits, respectively.

2.4. Larvicidal assay

The larvicidal activity was determined for the extracts in concentrations that prepared as 25, 50, 100, 200, and 400 ppm based on 1 g/1L (1000 ppm) from each extract stock solution against the 3rd larval instar of *Cx. pipiens* (WHO, 2005). Twenty *Cx. pipiens* third instar larvae were subjected to each extract concentration in 250 mL glass beakers containing 150 mL de-chlorinated water (aqueous suspension) at 27± 2 °C, 70± 10% RH, and a 12:12 h (L/D) photoperiod. Five replicates per concentration per extract and control were conducted. Larval mortalities were recorded after 24 and 48 h of exposure.

2.5. Identification of chemical compounds in extracts by Gas chromatography-mass spectrometer

Extracts from promising plant, *T. orientalis* L. were analyzed to investigate their chemical constituents by GC-MS using the Trace GC-TSQ mass spectrometer (Thermo Fisher Scientific, Waltham, MA, USA) through TG-5MS direct capillary column (30 m × 0.25 mm × 0.25 m thickness of the film) where the column oven temperature initially maintained at 50 °C, then rate was increased 5 °C/min up to 250 °C, for 2 min, and then increased 30 °C/min to 300 °C. The lines for the injector and MS transfer were suspended at 280 °C and 260 °C, respectively, helium was the carrier gas at the rate of 1 mL/min. The solvent delay was 3 min and 2µl samples were injected automatically using Autosampler AS1310 coupled with GC in the splitless mode. In full scanning mode, electrospray ionization (EI) mass spectra were obtained covering the range 50–650 m/s at an ionization voltage of 70 V. The ion source temperature was fixed at 200 °C. The chemical constituents were identified from the Total Ion Chromatogram (TIC), where the chemical compounds were identified by comparison of their retention times and mass spectra with those of WILEY 09 and NIST 11 mass spectral databases.

2.6. Data analysis

The percentage mortalities were determined according to Abbott, (1925). The larval control results did not need correction, as the mortality was less than 5%, according to the WHO guidelines (WHO, 2005) (no larval control mortality recorded throughout the study). Mortality data from all the replicates were performed using one-way analysis of variance (ANOVA) to find the differences among the activity between each plant extract concentrations using the least significant difference (LSD) test. Also, data from all the

replicates⁶ were subjected to analysis to determine the larval LC₅₀, LC₉₀, and LC₉₅ as well as chi-square values within confidence limits at 95% (lower confidence limit (LCL) and upper confidence limit (UCL))²⁸ by using probit analysis and regression between log extract concentration and probit values. Data analysis²⁷ was performed using SPSS software (IBM SPSS Statistics v22 – 64 bit), and p < 0.05 was considered significant.²¹

3. Results

3.1. Larvicidal activity

The data about the larvicidal activities² of the tested *T. orientalis* leaves extracts against the third instar larvae of *Cx. pipiens* after 24 h are summarized in Table (1), and after 48 h are summarized in Table (2). The analysis of responses of the leaves extracts revealed that the larvicidal activities after 24 h showed 100% mortalities at 400 ppm of methanol, acetone and aqueous extracts. While hexane leaves extract at 400 ppm showed 98% larval mortality. Acetone leaves extract showed the highest efficacy by inducing 100% mortality (LC₅₀, 58.04, LC₉₀, 169.27, and LC₉₅, 229.27 ppm), followed by methanol extract by 100% mortality (LC₅₀, 70.20, LC₉₀, 221.66, and LC₉₅, 307.07 ppm), followed by aqueous extract by 100% mortality (LC₅₀, 77.19, LC₉₀, 244.26, and LC₉₅, 338.95 ppm). While, hexane extract showed the least efficacy as compared to others, inducing 98.00% mortality (LC₅₀, 84.25, LC₉₀, 283.47, and LC₉₅, 399.84 ppm) Table (1). The larvicidal activities of the leaves extracts after 48 h showed 100% mortalities at 400 ppm for all tested extracts. Acetone leaves extract showed the highest efficacy (LC₅₀, 43.40, LC₉₀, 102.62, and LC₉₅, 130.98 ppm), followed by methanol extract (LC₅₀, 54.75, LC₉₀, 156.00, and LC₉₅, 209.92 ppm), followed by the aqueous extract (LC₅₀, 63.76, LC₉₀, 198.50, and LC₉₅, 273.89 ppm). Then, hexane showed (LC₅₀, 69.02, LC₉₀, 225.14, and LC₉₅, 314.78 ppm) Table (2).

² The results about the larvicidal activities of the tested *T. orientalis* fruits extracts against the third instar larvae of *Cx. pipiens* after 24 h are summarized in Table (3) and after 48 h are summarized in Table (4). The analysis of fruits extracts data revealed that the larvicidal activities after 24 h showed 100% mortalities at 400 ppm of methanol and hexane extracts. While acetone and aqueous fruits extract at 400 ppm showed 98% and 96% larval mortalities, respectively. Hexane fruits extract showed the highest

efficacy by inducing 100% mortality (LC₅₀, 68.26, LC₉₀, 201.48, and LC₉₅, 273.84 ppm), followed by methanol extract by 100% mortality (LC₅₀, 83.21, LC₉₀, 253.85, and LC₉₅, 348.25 ppm). The acetone extract induced 98% mortality (LC₅₀, 92.81, LC₉₀, 296.34, and LC₉₅, 411.83 ppm). While, the aqueous extract showed the least efficacy as compared to others, inducing 96% mortality (LC₅₀, 102.97, LC₉₀, 333.60, and LC₉₅, 465.54 ppm) Table (3). The larvicidal activities of the fruits extracts after 48 h showed 100% mortalities at 400 ppm for all tested extracts. Hexane extract showed the highest efficacy (LC₅₀, 50.60, LC₉₀, 119.30, and LC₉₅, 152.14 ppm), followed by methanol extract (LC₅₀, 68.62, LC₉₀, 205.88, and LC₉₅, 281.12 ppm), followed by the acetone extract (LC₅₀, 76.43, LC₉₀, 236.40, and LC₉₅, 325.58 ppm). While, aqueous showed (LC₅₀, 81.85, LC₉₀, 253.48, and LC₉₅, 349.23 ppm) Table (4).

3.2. Chemical analysis

The GC-MS analysis revealed that the main constituents of the *T. orientalis* leaves extracts were the sesquiterpenoids and terpenoids Table (5) and (Figure 1). The main % area for sesquiterpenoids compounds commonly detected in all extracts, were, cedrol (33.03, 35.77, 33.99 and 38.25%), caryophyllene (18.49, 25.31, 21.68 and 20.67%) and humulene (10.94, 15.22, 12.28 and 12.35%) for methanol, hexane, acetone and aqueous leaves extracts, respectively. Followed by, germacrene D (2.80, 4.00, 3.60 and 3.35%), allosedrol (2.26, 2.22, 2.16 and 2.43%), α -Eudesmol (1.95, 1.92, 2.91 and 2.58%), cadina-1(10),4-diene (1.29, 1.67, 1.30 and 1.58%), selinene (1.27, 1.18, 1.32 and 1.59%), γ -muurolene (1.17, 1.47, 1.08 and 1.26%), caryophyllene oxide (1.16, 1.25, 1.39 and 1.29%), cedrene (0.39, 2.66, 0.60 and 2.33%) and γ -elemene (0.28, 1.54, 0.62 and 1.52%) for methanol, hexane, acetone and aqueous leaves extracts, respectively. Also, fatty acid methyl ester, cis-4,7,10,13,16,19-docosahexaenoic acid, methyl ester (1.22, 1.31, 0.29 and 0.97%) for the same aforementioned extracts, respectively. Other compounds detected in three extracts only, pimara-7,15-dien-3-ol (6.04, 1.53 and 0.54 %), α -muurolene (0.29, 1.80 and 0.31%), alloaromadendrene (1.11, 2.11 and 0.64%) and butyl 4,7,10,13,16,19-docosahexaenoate (3.52, 1.43 and 3.00%) were, were detected in methanol, acetone and aqueous leaves extracts, respectively. The retinoid compound 4,14-retro-retinol represented by 1.44, 1.25 and 1.25 area % in hexane, acetone and aqueous leaves extracts, respectively. The compounds, Bornyl acetate (0.39 and 0.44%), α -terpinyl acetate (0.48 and

0.36%), β -elemene (0.52 and 1.73%), 1,7-Di-epi- β -cedrene (2.47 and 2.06%), valencen (1.05 and 1.44%), cedarn-8-ol (0.19 and 0.14%) and α -acorenol (1.20 and 1.64%) were recorded in methanol and acetone leaves extracts, respectively. Meanwhile, the terpenoid compound, ehydro-4-epiabietol represented by 0.73 and 1.03 area % and the sesquiterpenoid, γ -gurjunene represented by 0.33 and 1.10 area % in acetone and aqueous leaves extracts, respectively. There are compounds were detected only in the methanol leaves extract as 4-hydroxy-3,3',4-trimethoxystilbene (3.95%), sugiol (1.49%), totarol (0.52%) and copalol (0.51%). β -Chamigrene and linolenic acid, methyl ester were detected in hexane leaves extract and represented by 1.81 and 1.26 area %, respectively.

Elemyl acetate compound detected in acetone leaves extract only and represented by 0.75 area %. Cedrenol and aromandendrene compounds detected in aqueous leaves extract only and represented by 0.20 and 1.65 area %, respectively.

The GC-MS analysis revealed that the main constituents of the fruits extracts were represented in Table (6) and (Figure 2). The main recorded sesquiterpenoids compounds in all extracts were presented by ratios as, cedrol (21.26, 25.11, 18.55 and 36.66%) and caryophyllene (17.06, 19.93, 15.21 and 17.04%) for methanol, hexane, acetone and aqueous fruits extracts, respectively. Followed by the compounds, humuline (9.05, 10.93, 8.62 and 9.06%), germacrene D (4.93, 5.76, 5.16 and 4.44%), elemol (4.20, 4.57, 3.42 and 4.05%), bomyl acetate (3.14, 3.08, 2.48 and 3.35%), terpinolene (1.14, 2.75, 1.77 and 1.62%), α -terpinyl acetate (1.10, 0.93, 1.10 and 1.22%), cedrene (1.45, 1.36, 1.22 and 1.44%), allocedrol (1.48, 1.18, 1.26 and 1.84%), β -elemene (1.15, 1.28, 0.86 and 1.07%), caryophyllene oxide (1.08, 1.14, 1.04 and 1.62%), selinene (0.60, 0.59, 0.56 and 0.65%) for methanol, hexane, acetone and aqueous fruits extracts, respectively. Two compounds were detected in three extracts only, labda-8(20),12,14-triene (4.64, 1.80 and 8.02 %) and trachyloban (0.81, 0.63 and 0.52%), in methanol, hexane and acetone fruits extracts, respectively. Alloaromadendrene recorded in methanol and acetone fruits extracts by 0.48 and 0.53 area%, respectively. While, γ -elemene detected in methanol, hexane and aqueous extracts by 0.51, 0.53 and 0.42 area %, respectively. Isopimaral and olic acid detected in hexane and acetone extracts by 5.64, 5.64 and 2.02 and 0.76 area %, respectively. ⁴ cis-5,8,11,14,17-eicosapentaenoic acid, methyl ester represented by 7.11 and

3.48 area %, i-Propyl 7,10,13,16,19-docosapentaenoate represented by 2.26 and 2.26 area %, arachidonic acid methyl ester represented by 0.58 and 1.13 area %, 6,9,12,15-docosatetraenoic acid, methyl ester represented by 0.63 and 1.89 area %, dodecanoic acid, 2,3-bis(acetyloxy)propyl ester represented by 0.76 and 0.42 area %, and in methanol and aqueous extracts, respectively. β -Eudesmol compound detected in hexane and aqueous fruits extracts and represented by 0.80 and 0.70 area %, respectively.

There are 15 compounds detected only in methanol fruits extract only which were, 13,16-octadecadienoic acid, methyl ester (3.24%), Androst-5-ene, 4,4-dimethyl-, (13 λ)- (1.85), doconexent (1.42%), 2-Carene (1.00%), 1-heptatriacotanol (0.95%), communic acid (0.93%), cis-4,7,10,13,16,19-docosahexaenoic acid (0.84%), δ -elemene (0.68%), α -eudesmol (0.64%), Pseudoarsasapogenin-5,20-dien (0.54%), Cadina-1(10),4-diene (0.47%), 16-hydroxyingenol (0.45%), (-)-Atisirene (0.42%), isogermaene D (0.31%), linolenic acid, 2-hydroxy-1-(hydroxymethyl)ethyl ester (Z,Z,Z)-(0.29%). Six compounds detected only in hexane fruits extract which were, labda-8(20) 12 14-triene-19-oic acid methyl ester (z)-(2.95%), copalol (1.05%), isopimaric acid, methyl ester (1.00%), chamigrene (0.61%), Androstan-17-one, 3-ethyl-3-hydroxy-, (5 λ)-(0.60%) and Dotriacontane (0.45%). Four compounds detected only in acetone fruits extract were, α -linolenic acid, TMS derivative (12.23%), pimaric acid TMS derivative (8.43%), eicosapentaenoic acid, TMS derivative (1.81%) and pimara-7,15-dien-3-ol (1.46%). While, in aqueous fruits extract there are six compounds detected which were, labda-8(20),14-dien-13-ol, (13S)- (1.62%), valencen (1.05%), cubenol (1.48%), 7,10,13,16,19-docosapentaenoic acid, methyl ester (0.81%), 6,9,12-octadecatrienoic acid, methyl ester (0.62%) and 10-heptadecen-8-ynoic acid, methyl ester,(E)- (0.49%) Table (6) and (Figure 2).

4. Discussion

Generally, the study revealed larvicidal activities of *T. orientalis* leaves and fruits extracts in a concentration dependent manner whatever the activities varied between leaves and fruits tested extracts. Leaves extracts ordered according to larval toxicity as acetone > methanol > aqueous > hexane. While, fruits extracts ordered as hexane > methanol > acetone > aqueous. Collectively acetone *T. orientalis* leaves

extract showed throughout the tested extracts, the most effective larvicidal activity against *C. pipiens* larvae (100% mortality) and (LC₅₀, 58.04 ppm LC₉₀, 169.27 ppm LC₅₀, 229.27 ppm) at concentration 400ppm.

Different Plant extracts previously tested for their larvicidal activities against mosquitoes, El-Sheikh et al, (2012) tested the larval toxicity of ethanol, acetone and petroleum ether extracts of *Tribulus terrestris* leaves, against the third instar larvae *Ae. aegypti* and predicted dose dependent larvicidal activities for the tested extracts in addition to variation of larvicidal activities related to solvent used. Where, crude plant extracts showed previously more efficiency in controlling mosquitoes over the purified compounds toxicity, which in line with the present study results (Ghosh et al., 2012). Another study estimated larval toxicity of leaves aqueous extracts in three tested concentrations from *Ricinus communis* L. (0.06, 0.12 and 0.2 g/l), *Daphne gnidium* L (0.09, 0.18 and 0.3 g/l) and *Thymus vulgaris* L. (0.0225, 0.045 and 0.09 g/l), against early instar larvae of *Cx. pipiens* and *Cs. longiareolata* and showed in line with the present study results that the larval mortality increased with exposure times (24, 48 and 72 h), and the LC₅₀ recorded values decreased in the same manner (Dahchar et al., 2016).

The chemical compounds detected in the extracts in line with that detected in previous studies, however, there were differences in the amount or number of the main components (Elsharkawy et al., 2017; Guleria et al., 2008; Nickavar et al., 2003; Ololade et al., 2014; Sanei-Dehkordi et al., 2018).

The common sesquiterpenoids highly represented as % areas in the chromatograms of all tested leaves and fruits extracts were, Cedrol, Caryophyllene, Humulene, Germacrene D and Elemol (common only in fruits extracts). The medium common ones were, γ -Elemene, Cedrene, Selinene, γ -Muurolene, Cadina-1(10),4-diene, Caryophyllene oxide, Allocatedrol, α -Eudesmol and the fatty acid methyl ester, cis-4,7,10,13,16,19-Docosahexaenoic acid, methyl ester in leaves extracts and bornyl acetate, α -Terpinyl acetate, Cedrene, β -Elemene, Selinene and Caryophyllene oxide were medium common in fruits extracts. The observed larvicidal activities of the tested extracts may related to their chemical constituents synergistic actions, either the major or minor ones that may impact on the predicted larvicidal activities (Huong et al., 2020).

The essential oil *P. orientalis* L. (Family Cupressaceae) oil showed insecticide and molluscicidal activity (Hashemi and Safavi 2012; Ju-Hyun et al., 2005; Lei, et al., 2010). *T. orientalis* previously acquired cytotoxic principles and contained terpenoids including pimaric and isopimaric acids, fatty acids, aliphatic compounds like alkanes and bioflavonoids (Mehta et al., 1999). The essential oil extracted from *T. orientalis* leaves predicted larvicidal activity against *Anopheles stephensi* and *Culex pipiens* late third or young 4th instar larvae, where carene and cedrol were from the main constituents of the extract (Sanei-Dehkordi et al., 2018). Also extracts predicted diterpenes and labdane-type diterpenes in line with the previously recorded (Kim et al 2012 and 2013). Cedrol the main constituent in the tested extracts was postulated as alternative to conventional synthetic acaricides for the black-legged ticks, *Ixodes scapularis* Say which is a human vector causing disease (Eller et al., 2014). Caryophyllene oxide and germacrene D individual compounds previously showed potent larvicidal activities against *A. anthropophagus* (Zhu and Tian, 2013). Also β -elemene and α -humulene showed larvicidal activities against *A. subpictus*, *Ae. albopictus*, and *Cx. tritaeniorhynchus* (Govindarajan and Benelli, 2016). Jeon et al. (2005) showed that essential oils extracted from *P. orientalis* had strong activities on mosquito larvae *Cx. pipiens* and *Ae. aegypti* recording 100% mortality at 400 ppm. Furthermore, Sanei-Dehkordi et al. (2018) verified the high efficiency of *P. orientalis* oil against *A. stephensi* (LC₅₀, 11.67 ppm) and *Cx. pipiens* (LC₅₀, 18.60 ppm). Ethanol and acetone leaves extracts of *T. orientalis* previously evaluated larvicidal activities against mosquitoes recording LC₅₀, 13.10 and 200.87 ppm after 24 h and 9.02 and 127.53 ppm after 48 h, against *A. stephensi* third instar larvae, respectively. Besides recording against *C. quinquefasciatus* third instar larvae, LC₅₀, 22.74 and 69.03 ppm after 24h and 16.72 and 51.14 ppm after 48 h, respectively (Sharma et al., 2005).

In addition and in accordance with the present results behavior, hexane plant parts and seeds extracts of *Physalis angulate*, *Peganum harmala*, *Tecrium polium* and *Thymus vulgaris* were evaluated for their larvicidal potentials against the fourth instar larvae of *Culex pipiens molestus*, and showed that mortality increased with exposure times (24 and 48 h) and the LC₅₀ recorded values decreased with exposure times (Mekhlif and Muhammad, 2021). A study tested larvicidal activities of leaves oils extracts against *A.*

aegypti and recorded that the main components in leaves of *G. blepharophylla* was the caryophyllene oxide, while, in *G. friesiana* were α - β - and γ -eudesmols and in *G. hispida*, was (E)-caryophyllene. According to the predicted results oil extracted from *G. friesiana* recorded the best larvicidal effect against *A. aegypti* and hypothesized that sesquiterpenes in oils can reflect more controlling activity as compared to monoterpenes (Aciole et al., 2011). The aforementioned study results may declare the predicted advantage of acetone leaves extract results where the extract acquired highest content of eudesmol as compared to other extracts.

5. Conclusion

The leaves and fruits methanol, hexane, acetone and aqueous extracts of *T. orientalis* revealed high larvicidal potential at 400 ppm concentration ²⁶ against the third instar larvae of *Cx. pipiens* with the advantage of leaves acetone extract that predicted the lowest effective larval toxicity concentration. Further studies about the mode of the larvicidal action of the extracts should be under investigation.

5. References

- Abbott, W.S., 1925. A method of computing the effectiveness of an insecticide. J. econ. Entomol 18(2), 265–267. <https://doi.org/10.1093/jee/18.2.265a>
- Aciole S.D.G., Piccoli C.F., Duque J.E.L., Costa E.V., Navarro-Silva M.A., Marques F.A., Maia B.H.L.N.S., Pinheiro M.L.B., Rebelo M., 2011. Insecticidal activity of three species of Guatteria (Annonaceae) against *Aedes aegypti* (Diptera: Culicidae). Rev. Colomb. Entomol. 37, 262–268. http://www.scielo.org.co/scielo.php?script=sci_arttext&pid=S0120-04882011000200018&lng=en.
- Ahmed, N., Alam, M., Saeed, M., Ullah, H., Iqbal, T., Al-Mutairi, K.A., Shahjeer, K., Ullah, R., Ahmed, S., Ahmed, N.A.A.H., Khater, H.F., Salman, M., 2021. Botanical Insecticides Are a Non-Toxic Alternative to Conventional Pesticides in the Control of Insects and Pests, in: Global Decline of Insects. IntechOpen. <https://doi.org/10.5772/intechopen.100416>

- Azwanida, N.N., 2015. A review on the extraction methods use in medicinal plants, principle, strength and limitation. *Med. Aromat. Plants*, 4: 196 (03): 1-6. <https://doi.org/10.4172/2167-0412.1000196>
- Caruntu, S., Ciceu, A., Olah, N.K., Don, I., Hermenean, A., Cotoraci, C., 2020. *Thuja occidentalis* L.(Cupressaceae): Ethnobotany, phytochemistry and biological activity. *Molecules* 25(22), 5416.. <https://doi.org/10.3390/molecules25225416>
- Choi, H. A., Ahn, S. O., Lim, H. D., Kim, G. J., 2021. Growth Suppression of a Gingivitis and Skin Pathogen *Cutibacterium* (*Propionibacterium*) *acnes* by Medicinal Plant Extracts. *Antibiotics* (Basel, Switzerland), 10(9), 1092. <https://doi.org/10.3390/antibiotics10091092>
- Dahchar, Z., Bendali-Saoudi, F., Soltani, N., 2016. Larvicidal activity of some plant extracts against two mosquito species *Culex pipiens* and *Culiseta longiareolata*. *J. Entomol. Zool. Stud.* 4, 346–350.
- Darwish, R.S., Hammada, H.M., Ghareeb, D.A., Abdelhamid, A.S.A., Harraz, F.M., Shawky, E., 2021. Chemical profiling and identification of anti-inflammatory biomarkers of oriental *Thuja* (*Platyclusus orientalis*) using UPLC/MS/MS and network pharmacology-based analyses. *Nat. Prod. Res.* 1–5. <https://doi.org/10.1080/14786419.2021.2010198>
- Eller, F.J., Vander Meer, R.K., Behle, R.W., Flor-Weiler, L.B., Palmquist, D.E., 2014. Bioactivity of cedarwood oil and cedrol against arthropod pests. *Environ. Entomol.* 43(3), 762–766. <https://doi.org/10.1603/EN13270>
- Elsharkawy, E. R., Ali, A. M. H., 2019. Effect of Drought Condition of North Region of Saudi Arabia on Accumulation of Chemical Compounds, Antimicrobial and Larvicidal Activities of *Thuja Orientalis*. *Oriental Journal of Chemistry*, 35(2), 738–743. <https://doi.org/10.13005/ojc/350233>
- Elsharkawy, E.R., Aljohar, H., Donia, A.M., 2017. Comparative study of antioxidant and anticancer activity of *Thuja orientalis* growing in Egypt and Saudi Arabia. *Br. J. Pharm. Res* 15, 1–9. <https://doi.org/10.9734/bjpr/2017/32387>
- El-Sheikh, T.M.Y., Bosly, H.A.M., Shalaby, N., 2012. Insecticidal and repellent activities of methanolic extract of *Tribulus terrestris* L.(Zygophyllaceae) against the malarial vector *Anopheles arabiensis*

- (Diptera: Culicidae). Egypt. Acad. J. Biol. Sci. A, Entomol. 5, 13–22.
<https://doi.org/10.21608/eajbsa.2012.14784>
- Farajollahi, A., Fonseca, D.M., Kramer, L.D., Kilpatrick, A.M., 2011. “Bird biting” mosquitoes and human disease: a review of the role of *Culex pipiens* complex mosquitoes in epidemiology. *Infect. Genet. Evol.* 11, 1577–1585. <https://doi.org/10.1016/j.meegid.2011.08.013>
- Ghosh, A., Chowdhury, N., Chandra, G., 2012. Plant extracts as potential mosquito larvicides. *Indian J. Med. Res.* 135(5), 581–598. PMID: 22771587
- Govindarajan, M., Benelli, G., 2016. α -Humulene and β -elemene from *Syzygium zeylanicum* (Myrtaceae) essential oil: highly effective and eco-friendly larvicides against *Anopheles subpictus*, *Aedes albopictus*, and *Culex tritaeniorhynchus* (Diptera: Culicidae). *Parasitol. Res.* 115, 2771–2778.
<https://doi.org/10.1007/s00436-016-5025-2>
- Govindarajan, M., Rajeswary, M., Hoti, S.L., Bhattacharyya, A., Benelli, G., 2016. Eugenol, α -pinene and β -caryophyllene from *Plectranthus barbatus* essential oil as eco-friendly larvicides against malaria, dengue and Japanese encephalitis mosquito vectors. *Parasitol. Res.* 115(2), 807–815.
<https://doi.org/10.1007/s00436-015-4809-0>
- Guleria, S., Kumar, A., Tiku, A.K., 2008. Chemical composition and fungitoxic activity of essential oil of *Thuja orientalis* L. grown in the north-western Himalaya. *Zeitschrift für Naturforsch. C* 63, 211–214. <https://doi.org/10.1515/znc-2008-3-409>
- Hashemi, S.M., Safavi, S.A., 2012. Chemical constituents and toxicity of essential oils of oriental arborvitae, *Platycladus orientalis* (L.) Franco, against three stored-product beetles. *Chil. J. Agr. Res.* 72(2), 188–194. <https://doi.org/10.4067/s0718-58392012000200004>
- Huong, L.T., Huong, T.T., Huong, N.T.T., Hung, N.H., Dat, P.T.T., Luong, N.X., Ogunwande, I.A., 2020. Mosquito larvicidal activity of the essential oil of *Zingiber collinsii* against *Aedes albopictus* and *Culex quinquefasciatus*. *J. Oleo Sci.* 69(2), 153–160. <https://doi.org/10.5650/jos.ess19175>
- Jeon, J.H., Lee, S.H., Kim, M.K., Lee, H.S., 2005. Larvicidal activity of *Chamaecyparis obtusa* and *Thuja orientalis* leaf oils against two mosquito species. *J. Appl. Biol. Chem.* 48(1), 26–28.

- Ju-Hyun, J., Sang-Hyun, L., Moo-Key, K., Hoi-Seon, L., 2005. Larvicidal activity of *Chamaecyparis obusta* and *Thuja orientalis* leaf oils against two mosquito species. *Agric. Chem. Biotech.* 48 (1), 26-28.
- Kim, C.S., Choi, S.U., Lee, K.R., 2012. Three new diterpenoids from the leaves of *Thuja orientalis*. *Planta Med.* 78, 485–487. <https://doi.org/10.1055/s-0031-1298215>
- Kim, T.H., Li, H., Wu, Q., Lee, H.J., Ryu, J.H., 2013. A new labdane diterpenoid with anti-inflammatory activity from *Thuja orientalis*. *J. Ethnopharmacol.* 146, 760–767. <https://doi.org/https://doi.org/10.1016/j.jep.2013.02.001>
- Lei, H., Wang, Y., Liang, F., Su, W., Feng, Y., Guo, X., Wang, N., 2010. Composition and variability of essential oils of *Platycladus orientalis* growing in China. *Biochem. Syst. Ecol.* 38(5), 1000-1006. <https://doi.org/10.1016/j.bse.2010.09.018>
- Mehta, B.K., Sharma, M., Shitut, S., 1999. Identification of novel compounds from *Thuja orientalis* (leaves). *Indian J. Chem. - B Org. Med. Chem.* 38(8), 1005–1008.
- Mekhlif, A.F., Muhammad, M.J., 2021. Larvicidal potentials of four medicinal plant extracts on mosquito vector, *Culex pipiens molestus* (Diptera: Culicidae). *Int J Mosq Res.*, 8(4):01-05.
- Mouden, S., Klinkhamer, P.G.L., Choi, Y.H., Leiss, K.A., 2017. Towards eco-friendly crop protection: natural deep eutectic solvents and defensive secondary metabolites. *Phytochem. Rev.* 16, 935–951. <https://doi.org/10.1007/s11101-017-9502-8>
- Nickavar, B., Amin, G., Parhami, S., 2003. Volatile constituents of the fruit and leaf oils of *Thuja orientalis* L. grown in Iran. *Zeitschrift für Naturforsch. C* 58(3–4), 171–172. <https://doi.org/10.1515/znc-2003-3-404>
- Nizam, I., Mushfiq, M., 2007. Antioxidant activity of water and alcohol extracts of *Thuja orientalis* leaves. *Adv. Tradit. Med.* 7, 65–73. <http://dx.doi.org/10.3742/OPEM.2007.7.1.065>
- Ololade, Z.S., Fakankun, O.A., Alao, F.O., Udi, O.U., 2014. Phytochemical and Therapeutic Studies of the Fruit essential oil of *Thuja orientalis* from Nigeria. *Glob. J. Sci. Front. Res. B Chem* 14(7), 14–20.

- Sanei-Dehkordi, A., Gholami, S., Abai, M.R., Sedaghat, M.M., 2018. Essential oil composition and larvicidal evaluation of *Platyclusus orientalis* against two mosquito vectors, *Anopheles stephensi* and *Culex pipiens*. *J. Arthropod. Borne. Dis.* 12(2), 101–107. <https://doi.org/10.18502/jad.v12i2.35>
- Sharma, P., Mohan, L., Srivastava, C.N., 2005. Larvicidal potential of *Nerium indicum* and *Thuja orientalis* extracts against malaria and Japanese encephalitis vector. *J Environ Biol.* 26(4), 657–660.
- Shin, I., Shin, N., Jeon, C., Kwon, O., Hong, J., Kim, H., Oh, S., Ahn, K., 2015. *Thuja orientalis* reduces airway inflammation in ovalbumin-induced allergic asthma. *Mol. Med. Rep.* 12, 4640–4646. <https://doi.org/10.3892/mmr.2015.3910>
- Srivastava, P., Kumar, P., Singh, D.K., Singh, V.K., 2012. Biological properties of *Thuja orientalis* Linn. *Adv Life Sci* 2, 17–20. DOI: 10.5923/j.als.20120202.04
- WHO. 2005. Guidelines for laboratory and field testing of mosquito larvicides. World Health Organization, 1–41. Retrieved from http://whqlibdoc.who.int/hq/2005/WHO_CDS_WHOPES_GCDPP_2005.13.pdf?ua=1
- WHO. 2013. Larval source management: a supplementary malaria vector control measure, 1–2. https://apps.who.int/iris/bitstream/handle/10665/85379/9789241505604_eng.pdf
- Wilson, A.L., Courtenay, O., Kelly-Hope, L.A., Scott, T.W., Takken, W., Torr, S.J., Lindsay, S.W., 2020. The importance of vector control for the control and elimination of vector-borne diseases. *PLoS Negl. Trop. Dis.* 14, e0007831. <https://doi.org/10.1371/journal.pntd.0007831>
- Zhang, N., Park, D.K., Park, H.-J., 2013. Hair growth-promoting activity of hot water extract of *Thuja orientalis*. *BMC Complement. Altern. Med.* 13, 1–12. <https://doi.org/10.1186/1472-6882-13-9>
- Zhu, L., Tian, Y., 2013. Chemical composition and larvicidal activity of essential oil of *Artemisia gilvescens* against *Anopheles anthropophagus*. *Parasitol. Res.* 112(3), 1137–1142. <https://doi.org/10.1007/s00436-012-3243-9>

Chemical characterization and Larvicidal Potential of Thuja orientalis Leaves and Fruits Extracts against Culex pipiens (Diptera: Culicidae)

ORIGINALITY REPORT

13%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|-----------------|
| 1 | Abdel-Shafi, Al-Mohammadi, Hamdi, Moustafa, Enan. "Biological Characterization and Inhibition of Streptococcus pyogenes ZUH1 Causing Chronic Cystitis by Crocus sativus Methanol Extract, Bee Honey Alone or in Combination with Antibiotics: An In Vitro Study", <i>Molecules</i> , 2019
Crossref | 57 words — 1% |
| 2 | www.entomoljournal.com
Internet | 40 words — 1% |
| 3 | bjbas.springeropen.com
Internet | 36 words — 1% |
| 4 | idoc.pub
Internet | 25 words — 1% |
| 5 | www.intechopen.com
Internet | 18 words — < 1% |
| 6 | www.science.gov
Internet | 18 words — < 1% |
| 7 | Hanaa Mahmoud Mohamed, Magdy Sayed Aly. "Evaluation of genotoxicity of Euphorbia | 15 words — < 1% |

triaculeata Forssk. extract on mice bone marrow cells in vivo",
Toxicology Reports, 2018

Crossref

8 Osama A. Al-Mahmood, Nour H. Aboalhaja, 13 words — < 1%
Ismail F. Abaza, Wamidh H. Talib, Fatma U. Afifi.

"Investigations on the spontaneous emitted and hydrodistilled
volatiles of Platycladus orientalis (L.) Franco grown in four
different biogeographic zones of Jordan", Biochemical
Systematics and Ecology, 2020

Crossref

9 Prafull Salvi, Gulshan Kumar, Nishu Gandass, 13 words — < 1%
Kajal, Ashish Verma, Sivasubramanian

Rajarammohan, Nilesh Rai, Vibhav Gautam. "Antimicrobial
Potential of Essential Oils from Aromatic Plant Ocimum sp.; A
Comparative Biochemical Profiling and In-Silico Analysis",
Agronomy, 2022

Crossref

10 Hsiang-Ting Huang, Chien-Chung Lin, Tai-Chih 12 words — < 1%
Kuo, Shiang-Jiuun Chen, Rong-Nan Huang.

"Phytochemical composition and larvicidal activity of essential
oils from herbal plants", Planta, 2019

Crossref

11 Selladurai Subarani, Selvi Sabhanayakam, 12 words — < 1%
Chinnaperumal Kamaraj, Gandhi Elango,

Mohamed Abdul Kadir. "Efficacy of larvicidal and pupicidal
activity of Catharanthus roseus aqueous and solvent extracts
against Anopheles stephensi Liston and Culex quinquefasciatus
Say (Diptera: Culicidae)", Asian Pacific Journal of Tropical
Medicine, 2013

Crossref

12 apjtm.org 12 words — < 1%
Internet

13 Enaam M. AbouZeid, Ahmed H. Afifi, Abeer Salama, Rehab A. Hussein, Fadia S. Youssef, Sherweit H. El-Ahmady, Nagwa Mohamed Ammar. "Comprehensive metabolite profiling of Phoenix rupicola pulp and seeds using UPLC-ESI-MS/MS and evaluation of their estrogenic activity in ovariectomized rat model", Food Research International, 2022

11 words — < 1%

Crossref

14 Ibrahim Taha Radwan, Mohamed M. Baz, Hanem Khater, Abeer Mousa Alkhaibari, Abdelfattah M. Selim. "Mg-LDH Nanoclays Intercalated Fennel and Green Tea Active Ingredient: Field and Laboratory Evaluation of Insecticidal Activities against Culex pipiens and Their Non-Target Organisms", Molecules, 2022

11 words — < 1%

Crossref

15 Lata Batabyal. "Relative toxicity of neem fruit, bitter gourd, and castor seed extracts against the larvae of filaria vector, Culex quinquefasciatus (Say)", Parasitology Research, 06/27/2009

11 words — < 1%

Crossref

16 bioone.org

Internet

11 words — < 1%

17 jddtonline.info

Internet

11 words — < 1%

18 jgeb.springeropen.com

Internet

11 words — < 1%

19 publish.kne-publishing.com

Internet

11 words — < 1%

20 repository.essex.ac.uk

Internet

11 words — < 1%

21	www.scielo.br Internet	11 words — < 1%
22	Abdel-Fattah D. Dawood, Mohamed M. Baz, Moustafa I. Ibrahim. "Influence of Aquatain™, a monomolecular surface film on surface tension for controlling the filarial vector <i>Culex pipiens</i> (Diptera: Culicidae)", Heliyon, 2020 Crossref	10 words — < 1%
23	Sharma, P.. "Phytoextract-induced developmental deformities in malaria vector", Bioresource Technology, 200609 Crossref	10 words — < 1%
24	iresa.agrinet.tn Internet	10 words — < 1%
25	iwaponline.com Internet	10 words — < 1%
26	nanobiofoods.com Internet	10 words — < 1%
27	openaccess.sgul.ac.uk Internet	10 words — < 1%
28	preview-behavioralandbrainfunctions.biomedcentral.com Internet	10 words — < 1%
29	www.chembuyersguide.com Internet	10 words — < 1%
30	www.dipterajournal.com Internet	10 words — < 1%

31 "ION TRAP MASS SPECTRA OF COMPOUNDS", Elsevier BV, 1989 9 words — < 1%
Crossref

32 Giovanni Benelli, Mohan Rajeswary, Marimuthu Govindarajan. "Towards green oviposition deterrents? Effectiveness of *Syzygium lanceolatum* (Myrtaceae) essential oil against six mosquito vectors and impact on four aquatic biological control agents", Environmental Science and Pollution Research, 2016 9 words — < 1%
Crossref

33 K. Dass, P. Mariappan. "Larvicidal Activity of *Colocasia esculenta*, *Eclipta prostrata* and *Wrightia tinctoria* Leaf Extract Against *Culex quinquefasciatus*", Proceedings of the National Academy of Sciences, India Section B: Biological Sciences, 2014 9 words — < 1%
Crossref

34 Khaoula Hellal, Ahmed Mediani, Intan Safinar Ismail, Chin Ping Tan, Faridah Abas. "1H NMR-based metabolomics and UHPLC-ESI-MS/MS for the investigation of bioactive compounds from *Lupinus albus* fractions", Food Research International, 2021 9 words — < 1%
Crossref

35 Rafia Rehman, Muhammad Zubair, Afsar Bano, Peter Hewitson, Svetlana Ignatova. "Isolation of industrially valuable α -Cedrol from essential oil of *Platycladus orientalis* (*Thuja orientalis*) leaves using linear gradient counter current chromatography", Industrial Crops and Products, 2022 9 words — < 1%
Crossref

36 Vinayachandra, R. Shwetha, K. R. Chandrashekar. "Larvicidal activities of *Knema attenuata* (Hook. f. & Thomson) Warb. (Myristicaceae) extracts against *Aedes*

albopictus Skuse and Anopheles stephensi Liston", Parasitology Research, 2011

Crossref

37	ir.uiowa.edu Internet	9 words — < 1%
38	pergamos.lib.uoa.gr Internet	9 words — < 1%
39	www.orientjchem.org Internet	9 words — < 1%
40	www.plantsjournal.com Internet	9 words — < 1%

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES

EXCLUDE MATCHES

OFF

< 9 WORDS