

Research Paper (HM Risk Assessment)

Submission date: 13-Nov-2022 10:44PM (UTC+0500)

Submission ID: 1952500736

File name: Revised_HM_Manuscript.docx (459K)

Word count: 4716

Character count: 27815

5

Human Health Risk Hazards by Heavy Metals Through Consumption of Vegetables Cultivated by Wastewater

Shahla Andleeb^a, Khalil ur Rehman^{a*}, Adeel Mahmood^a, Noor Ul Safa^a,

5

^a Department of Environmental Science Government College Women University Sialkot, Pakistan.

39

*Corresponding author: Adeel Mahmood (adilqau5@gmail.com)


Highlights

32

- Heavy metals in waste- and ground-water irrigated soil and plants were evaluated.
- Daily intake of metal and health risk index of heavy metals was calculated.
- Metal transfer factor from soil to vegetables was determined.
- Wastewater-irrigated mustard leaf and cabbage had highest HRI for Pb and was > 1.
- EF > 1.5 for Zn, Mn, Ni and Pb suggests metals were of anthropogenic origin

15

Graphical Abstract



Abstract

Need for irrigated agriculture is rising daily, but quality water for irrigation is on the decline, necessitating the use of urban wastewater as alternative source, particularly in low or middle income countries. This study assessed the effect of urban wastewater irrigation on the quality of groundwater, soil and vegetables in farms around

Faisalabad, Pakistan. Human health risks of heavy metals were investigated through consumption of wastewater irrigated vegetables. Samples of soil, vegetables and water were obtained and analyzed for heavy metal concentrations (cadmium, Cd; lead, Pb; manganese, Mn; nickel, Ni; cobalt, Co; and zinc, Zn). The groundwater could be declared safe for consumption in the present state, as concentration of heavy metals was well below the WHO (1996) standards. But wastewater-irrigated vegetables had higher Pb, Cd and Mn than the permissible limits. In wastewater-irrigated vegetables, highest HRI-Pb was recorded in mustard leaf and cabbage and was > 1. EF for Zn and Mn in all vegetable plants, Ni in potato and cauliflower, Pb in mustard leaf and cabbage were > 1.5, suggesting that the metals were generated by anthropogenic processes (such as wastewater irrigation). Long-term irrigation of farmlands with the wastewater will result to heavy metal contamination of groundwater, soil and vegetables in the study area. Therefore, strategies to save the groundwater from future contamination are necessary.

Keywords: Contamination; DIM; enrichment factor; Faisalabad; heavy metal concentrations; Health risk index

Introduction

Groundwater plays a significant role as a sustainable source of water for domestic and industrial purposes, particularly in the semi-arid and arid regions. It is estimated that over 4 million people in Faisalabad city, Pakistan rely on groundwater for consumption and other domestic purposes (Ahmed et al., 2019; Rani et al., 2021). However, the quality of groundwater may be altered due to unsustainable anthropogenic activities (Zwolak et al., 2019; Qayoom, 2021). For example, arsenic (As) contamination was found in groundwater in

Hasilpur, Pakistan, with hazard quotient (HQ) and carcinogenic risk up to 58 and 2.3×10^{-3} , respectively (Mahmood et al., 2013; Tabassum et al., 2019; Alam et al., 2021). Higher As concentration to levels that pose health risk has also been found in groundwater-irrigated vegetables in the central region of Italy (Singh and Raj, 2018; Spognardi et al., 2019). Also, in the south of Setif Area of East Algeria, cadmium (Cd) and lead (Pb) concentrations in the groundwater had an HQ above recommended safe limits with likelihood of posing health risk. Irrigation of farmlands with urban wastewater is a known source of groundwater contamination ((Rehman et al., 2019; Alam et al., 2022). Wastewater contains pollutants, resulting in contaminant transfer into irrigated soils leading to groundwater contamination as well as the contamination of irrigated vegetables (Sandeep et al., 2019, Hussain et al., 2019). Consumption of contaminated groundwater or plants could result in serious health problems, such as neurological, genotoxic, metagenomics, carcinogenic and respiratory effects (Mehmood et al., 2019; Mahmood et al., 2021).

In Faisalabad, vegetables sold in the supply market are produced by local farmers who irrigate their fields with groundwater, or largely with wastewater collected from different points along the sewer channels. Therefore, this study, carried out on farmers' fields in the region of Faisalabad, Pakistan, was conducted to assess the heavy metals contamination levels in the (1) groundwater compared to the wastewater and (2) in vegetables irrigated with groundwater compared to wastewater in this region, and (3) to determine the potential risks associated with consuming contaminated vegetables by using the daily intake (DIM) and transfer factor of metals (MTF) to compute health risk index (HRI) and Enrichment Factor (EF).

Materials and methods

Description of study site

Faisalabad is an industrial city in the district with the same name. It is the second largest city in Punjab province and the third biggest city in Pakistan (Fig. 1). The city is situated between the Chenab and Ravi rivers. Irrigation is based on water from these two rivers. To conduct the present study, two sites with different irrigation practices were identified based on a previous survey of Faisalabad. These sites have a long history of wastewater or groundwater irrigation. From each site, three sub-sites were selected, from where samples were collected for analyses. Soil, water, vegetable samples were collected from each selected site according to methods described by the American Public Health Association (APHA, 2005).

Collection of soil, water and plant samples

A quantity of 500 g soil sample was collected in triplicate from a monolith of 10 × 10 × 15 cm³ from each sub-site. Before collecting in plastic bags, materials like rocks, gravels and organic debris were removed manually from the soil.

Water (groundwater and wastewater) samples were filtered before storing in labelled 1 L plastic bottles that had earlier been soaked in 10% HNO₃ for 24 h and afterwards washed with deionized water to remove free soap and contaminants. The water samples were stored on ice before transporting to the laboratory where they were preserved at 4°C for further analyses (APHA, 2005).

List of vegetable crops collected from the sub-sites of each zone is given in Table 1. Five replicates of each vegetable crop were collected per sampling zone. The vegetables were washed and oven dried (65 °C) in the laboratory and preserved for further analyses.

Sample digestion

One gram (1 g) of each oven dried and ground vegetable parts or soil samples was digested at 80 °C in a 15 ml of tri-acid mixture containing HNO₃, H₂SO₄, and HClO₄ in 4.5:1:1 ratio, to

¹⁰ obtain a clear solution (APHA, 2005). The solution was further filtered and the filtrate ⁹ was made up to 50 ml by adding distilled water.

Water (groundwater and wastewater) samples (50 ml) were digested at 80 °C in a 10 ml conc. ¹⁰ HNO₃ to obtain a clear solution (APHA, 2005), that was further ²⁵ filtered and the filtrate made up to 50 ml by adding distilled water.

Analysis of heavy metals

³⁶ Heavy metal concentrations in the soil, plant and water filtrates obtained above were detected through atomic absorption spectrophotometer (UVAS, Lahore). The machine was manually calibrated using standard solutions and a drift blank

Data analyses

Data analyses were carried out as previously described in Rehman et al., (2019). Metal transfer factor (MTF) from soil to plants ²⁸ was calculated as the concentration of heavy metals in dry plant parts divided by that in the soil. ²⁷ Daily Intake of Metal (DIM) was calculated according to Chary et al., (2008). The mean daily vegetable intake (320 mg person⁻¹ day⁻¹) from 200 people with a regular body weight of 52.50 kg was used in the final computation. A dose of oral reference was used to compare with the test crops to calculate the ³⁷ health risk index (HRI) (Cui et al., 2004). Approximated exposure was measured by dividing heavy metals daily intake with their safe limits. HRI < 1 is considered safe for consumption, but may raise health concerns when HRI > 1 (USEPA, 2006). Heavy metal translocation from soil to edible vegetables was determined by calculating the Metal Enrichment Factor (EF) according to Chopra and Pathak, (2012).

Statistical analysis

Data was analyzed using SPSS v. 12. Results were compared with permissible limits by applying descriptive statistics.

Results

Concentration of heavy metals in (ground and waste) water and soil samples

Heavy metal concentrations were greater in wastewater samples than the groundwater samples (Table 2). Metal concentrations in the groundwater samples varied from < 0.01 to 1.50 mg L^{-1} . Zn had the highest ($1.10 \pm 0.23 \text{ mg L}^{-1}$) and Cd ($0.002 \pm 0.001 \text{ mg L}^{-1}$) the lowest mean concentration. All metals sampled in the groundwater were below the permissible limits. In the wastewater samples, heavy metal concentrations varied from 0.01 to 25.50 mg L^{-1} . Mn had the highest ($19.10 \pm 5.10 \text{ mg L}^{-1}$) and Ni ($0.02 \pm 0.01 \text{ mg L}^{-1}$) the lowest mean concentration. Mn concentration in wastewater samples was about 125-fold greater than the WHO limit of 0.2 mg L^{-1} , whereas other metals were well below the permissible limits.

In soil irrigated with wastewater, heavy metal concentration ranged from 0.30 to 39.50 mg kg^{-1} . Pb had the highest ($26.80 \pm 8.00 \text{ mg kg}^{-1}$) and Cd ($0.80 \pm 0.30 \text{ mg kg}^{-1}$) the lowest mean concentration (Table 2). In groundwater-irrigated soil, heavy metals were in the range 0.01 to 1.20 mg kg^{-1} , with Mn having the highest ($1.00 \pm 0.15 \text{ mg kg}^{-1}$) mean concentration and lowest mean concentration was found in Cd ($0.01 \pm 0.01 \text{ mg kg}^{-1}$) (Table 2). In soil irrigated with both water sources, mean metal concentrations were lower than the WHO limits.

Determination of metal concentrations in vegetable parts

The mean metal concentrations were greater in plants grown with wastewater compared to groundwater (Table 3). In plants grown with groundwater, metal concentration was found highest for Mn in mustard leaf ($14.2 \pm 4.10 \text{ mg kg}^{-1}$) and cabbage ($10.6 \pm 1.50 \text{ mg kg}^{-1}$). Among the vegetables, Pb ($0.060 \pm 0.20 \text{ mg kg}^{-1}$), Cd ($0.003 \pm 0.001 \text{ mg kg}^{-1}$), Zn ($0.200 \pm 0.07 \text{ mg kg}^{-1}$)

²³), Mn (14.2 ± 4.10 mg kg⁻¹) and Ni (0.080 ± 0.01 mg kg⁻¹) were highest in mustard leaf, but were lower than the WHO standards. Cobalt was highest in cabbage (0.520 ± 0.28 mg kg⁻¹) and mustard leaf (0.414 ± 0.30 mg kg⁻¹), but ³⁴ was much lower than the WHO standard (40.00 mg kg⁻¹). ⁴⁰ In the wastewater-irrigated vegetables, metal concentration was found highest for Mn in mustard leaf (32.80 ± 5.03 mg kg⁻¹) and cabbage (20.60 ± 3.40 mg kg⁻¹). Among the ¹⁶ vegetables, Pb (24.20 ± 4.30 mg kg⁻¹), Cd (0.10 ± 0.05 mg kg⁻¹), Co (1.70 ± 0.40 mg kg⁻¹), Zn (12.80 ± 1.00 mg kg⁻¹), Mn (32.80 ± 5.03 mg kg⁻¹) and Ni (3.20 ± 0.40 mg kg⁻¹) were highest in ² mustard leaf. In all vegetables, Pb concentration (2.60 ± 0.58 mg kg⁻¹ to 24.20 ± 4.30 mg kg⁻¹) was about 52 to 480 folds greater than the WHO standard (0.05 mg kg⁻¹). In mustard leaf and cabbage, Cd and Mn concentrations were about 5- and 2-fold higher than the standard. ¹

Heavy metals transfer factor (MTF) from irrigated soil to vegetable parts

MTF for Pb, Zn, Mn and Ni was greater in wastewater than groundwater-irrigated vegetables (Table 4). In the groundwater-irrigated vegetables, MTF-Cd was greatest in mustard leaf (14.20) and cabbage (10.60). Among the vegetables, MTF for Pb (0.540), Cd (14.200), Co (0.169), Zn (0.418) and Mn (0.281) were highest in mustard leaf. MTF-Ni was highest in cabbage (0.743) and mustard leaf (0.591). Among the wastewater-irrigated vegetables, highest MTF was observed for Mn (2.482) and Zn in mustard leaf (1.928). Among the vegetables, MTF for Pb (0.903), Cd (1.294), Co (0.167), Zn (1.928), Mn (2.482) and Ni (0.744) were highest in mustard leaf.

Daily Intake Metal (DIM) and Health Risk Index (HRI)

Among groundwater-irrigated vegetables, DIM for Pb ($3.00E-05$), Cd ($7.40E-03$), Co ($2.00E-06$), Zn ($4.00E-05$) and Mn ($1.00E-04$) were highest in mustard leaf (Table 5). DIM for Ni was highest in cabbage ($3.00E-04$) and mustard leaf ($2.00E-04$). Among wastewater-irrigated

vegetables, DIM for Pb (3.00E-05), Cd (7.40E-03), Co (2.00E-06), Zn (4.00E-05), Mn (1.00E-04) and Ni (9.00E-04) were greatest in mustard leaf.

Mustard (2.23E-01) and cabbage (1.66E-01) grown with groundwater showed the greatest HRI-Cd, but was lower than 1 (Table S1). Among the vegetables, HRI for Pb (8.00E-03), Cd (2.23E-01), Co (1.00E-04), Zn (2.00E-03) and Mn (4.00E-04) were highest in mustard leaf. HRI-Ni was highest in cabbage (6.30E-03) and mustard leaf (5.00E-03). In wastewater-irrigated vegetables, the highest HRI was observed for Pb in mustard leaf (3.14E+00) and cabbage (3.11E+00), and were greater than 1. Among the vegetables, HRI for Pb, Cd, Co, Zn, Mn and Ni were greatest in mustard leaf (3.14E+00, 5.16E-01, 2.00E-03, 8.29E-02, 2.20E-02 and 2.10E-03, respectively) followed by in cabbage (3.11E+00, 3.23E-01, 8.00E-04, 6.84E-02, 4.50E-03 and 2.00E-03, respectively).

Heavy Metals Enrichment Factor

EF for Pb was greatest in mustard leaf (1.672) and cabbage (1.658), but was lowest in cauliflower (0.565) (Table S2). EF for Cd and Ni was highest in cauliflower (0.333, 5.145) but lowest in cabbage (0.021, 0.932). EF for Co and Zn was highest in cabbage (1.243, 12.446), but Co was lowest in potato (0.018) and Zn in cauliflower (1.702). EF for Mn was highest in potato (25.962) and cabbage (17.589) but lowest in cauliflower (7.228).

Discussion

Metal concentrations in different water sources and irrigated soil

In this study, and as expected, metal concentrations in the wastewater samples were multiple folds greater than in the groundwater samples. This is because wastewater is generally from diverse sources of industrial and domestic origin and are well known sources of pollutants. In the wastewater samples, heavy metals were in the order Mn > Pb > Cd > Co > Zn > Ni, but in

the groundwater samples, were in the order Zn > Mn > Pb > Co > Ni > Cd. Our results also showed that Cd and Mn concentrations were greater in the wastewater samples than the WHO limits, whereas all metals sampled in the groundwater were below the limits. Results also attributed greater soil heavy metals content to wastewater irrigation, compared to groundwater irrigation; but the soil heavy metal contents were below the WHO standards. Type of soil (amount of clay content), soil depth, length of time of wastewater irrigation are among factors that contribute to heavy metals contamination of groundwater (Abdelwaheb et al., 2019). Abdelwaheb et al., (2019) noted greater groundwater contamination in soils with low clay content, but high clay content minimized contaminant leaching.

Wastewater, if untreated before applying to agricultural fields, could be a potential source of various organic and inorganic contaminants (Rehman et al., 2019), leading to the contamination of groundwater. Faisalabad generates well above $6.45 \text{ m}^3 \text{ s}^{-1}$ wastewater effluents (WWF, 2007), bulk of which is disposed into rivers Ravi and Chenab through the Madhuana and Paharang drains and channels. Seepage from these drains and channels, coupled with the utilization of wastewater for watering agricultural lands, are major routes of soil and groundwater contamination in this region.

According to the WB-SCEA, (2006), only about 8% of urban wastewater generated in Pakistan was treated before disposing into rivers. The most important strategy to preserve the quality of the groundwater is by reducing the contaminant concentrations in the wastewater before releasing into the environment. Sun et al., (2019) described the use of a self-sustained photo-bio electrochemical fuel cell for the removal of oxytetracycline in wastewater. Also, low-cost adsorbents of agricultural sources, such as sawdust, animal manures, eggshell waste and plant residues have been effectively used to reduce metal contaminants and to improve aerobic conditions in wastewater (Zajda and Aleksander-Kwaterczak, 2019). Azadi et al.,

(2018) used extracts from *Persicaria bistoria* plants to synthesis Fe₃O₄ nanoparticles that were used for wastewater treatment. Bacteria isolated from wastewater have also been utilized for heavy metal decontamination of wastewater. Cai et al., (2019) isolated Ni²⁺ resistant strains and Mn²⁺ tolerant strains for removal of respective metals in wastewater. If well treated, wastewater could provide a dependable source of water for irrigation in water-scarce regions.

Heavy metal concentration in vegetable plants

²⁶ Heavy metals (Pb, Zn, Mn, Cd, Co, and Ni) were highest in leaves of mustard and cabbage than other vegetables irrigated with wastewater; ²² concentrations of Pb, Mn and Cd were higher than the WHO permissible standards. ⁷ Bioaccumulation of heavy metal in vegetable plants is an important concern to public health. Here, we found that irrigation with groundwater did not result in high metal concentrations in the plants, because the water quality is still considered good. However, vegetable irrigation with wastewater could lead to increased metal uptake into plant parts. Mahmood and Malik, (2014) reported Pb and Cd above the European Union limits in vegetables grown with wastewater. Khan et al., (2019) also found Cr in grains of wheat irrigated with wastewater exceeded the permissible limit. ³⁵ Bashir et al., (2009) found metals concentrations in edible parts of vegetables irrigated with wastewater exceeded the permissible limit.

⁴⁴ For all heavy metals in groundwater-irrigated vegetables, Cd had the highest metal transfer factors (MTF) and was found in mustard leaf and cabbage; but in wastewater-irrigated vegetables, Mn and Zn had the highest MTF and were found in mustard leaf. The MTF-Cd in the groundwater-irrigated vegetables in this study are well above those in vegetables grown with groundwater in Lahore, Pakistan (Rehman et al., 2019), in vegetables grown with surface water in the vicinity of river Padma, Bangladesh and in food crops grown near

Dabaoshan mine, South China. Although our results indicate that the groundwater did not contribute to increased metal contents in the vegetables beyond permissible limits, a high MTF-Cd in the vegetables could suggest the need to employ rapid remediation strategies so as to save the groundwater from contamination.

³ **Daily intake of metals (DIM), health risk index (HRI) and Enrichment factors (EF)**

In both groundwater and wastewater-irrigated vegetables, highest DIM was observed for Cd and Pb in mustard leaf and cabbage; but, were below those observed in mustard leaf and cabbage grown in open fields in the vicinity of Chinese industrial zones. Compared to the dietary reference intake limits of the Food and Drug Administration, DIM-Pd and Cd were higher in wastewater, but not groundwater, irrigated mustard leaf and cabbage, suggesting the potential health risk to consumers of such ⁴⁶vegetables. In this study, HRI values for groundwater-irrigated vegetables were generally lower than 1, suggesting that consumers are not likely to be exposed to any health problems. Though HRI values for wastewater-irrigated vegetables were mostly lower than 1, mustard leaf and cabbage had HRI < 1 for Pb, suggesting a possibility of potential health risk due to Pb toxicity. Cd showed the potential to increase in HRI, though still lower than 1. The EF values in metals ⁶are used to assess the presence and intensity of anthropogenic contaminant deposition on surface soil. According to research, EF values between 0.5 and 1.5 suggest the metal could be wholly due to materials from ⁷the Earth's crust or natural processes, but EF > 1.5 suggests metals could be from anthropogenic sources. Therefore, our results suggest that Mn, Zn, Pb and Ni were generated by man-made processes (e.g. irrigation with wastewater).

Conclusion

Results revealed that wastewater irrigation is a major source of heavy metal contamination in soil and crops of vegetable gardens in Faisalabad. Heavy metal concentrations in the groundwater and irrigated crops in this region have not exceeded the WHO allowable limit, this could be termed safe for consumption. Health risk index in groundwater-irrigated vegetables was lower than 1; but in wastewater-irrigated mustard leaf and cabbage, HRI was greater than 1. Although the groundwater and irrigated vegetables cultivated in the regions of Faisalabad could be termed safe for consumption, there is a potential of heavy metal contamination over time. Therefore, there is the need to urgently embark on mitigation strategies that will both save the groundwater from being contaminated, while also providing adequate treatment plans for the large volume of wastewater generated in Faisalabad, Pakistan.

Conflicts of interest/Competing interests

The authors confirm there is no conflict of interest among them.

Acknowledgement

References

Abdelwaheb, M., Jebali, K., Dhaouadi, H., & Dridi-Dhaouadi, S. (2019). Adsorption of nitrate, phosphate, nickel and lead on soils: Risk of groundwater contamination. *Ecotoxicology and environmental safety*, 179, 182-187. doi.org/10.1016/j.ecoenv.2019.04.040.

Ahmed, A., Nasir, A., Basheer, S., Arslan, C., & Anwar, S. (2019). Ground water quality assessment by using geographical information system and water quality index: a case study of Chokera, Faisalabad, Pakistan. *Water Conservation and Management*, 3(1), 7-19. DOI : <http://doi.org/10.26480/wcm.01.2019.07.19>.

Alam A., Chaudhry M.N., Ahmad S.R., Batool S.A., Mahmood A., & Al-Ghamdi H.A. (2021). Application of Easewaste Model for Assessing Environmental Impacts from Solid Waste Landfilling. *Archives of Environmental Protection*, 47(4), 84–92. DOI: 10.24425/Aep.2021.139504.

Alam A., Mahmood, A., Chaudhry, M.N., Ahmad S.R., Safa, N., Alghamdi, H.A., Alhamdi, H. W., Ullah, R. (2022). Application of Easewaste Model for Assessing Environmental Impacts from Solid Waste Landfilling. *Archives of Environmental Protection*, 48(3).

APHA. (2005) Standard Methods for the Examination of Water and Wastewater, 21st ed. American Public Health Association, Washington DC., 1220

Azadi, F., Karimi-Jashni, A., & Zerafat, M. M. (2018). Green synthesis and optimization of nano-magnetite using *Persicaria bistorta* root extract and its application for rosewater distillation wastewater treatment. *Ecotoxicology and Environmental Safety*, 165, 467-475. <https://doi.org/10.1016/j.ecoenv.2018.09.032>.

Bashir, F., Kashmiri, M. A., Shafiq, T., & Tariq, M. (2009). Heavy metals uptake by vegetables growing in sewage irrigated soil: relationship with heavy metal fractionation in soil. *Chemical Speciation & Bioavailability*, 21(4), 199-209. <https://doi.org/10.3184/095422909X12471558119088>.

- Cai, X., Zheng, X., Zhang, D., Iqbal, W., Liu, C., Yang, B., ... & Mao, Y. (2019). Microbial characterization of heavy metal resistant bacterial strains isolated from an electroplating wastewater treatment plant. *Ecotoxicology and environmental safety*, 181, 472-480. <https://doi.org/10.1016/j.ecoenv.2019.06.036>.
- Chary, N. S., Kamala, C. T., & Raj, D. S. S. (2008). Assessing risk of heavy metals from consuming food grown on sewage irrigated soils and food chain transfer. *Ecotoxicology and environmental safety*, 69(3), 513-524. <https://doi.org/10.1016/j.ecoenv.2007.04.013>.
- Chopra, A. K., & Pathak, C. (2012). Bioaccumulation and translocation efficiency of heavy metals in vegetables grown on long-term wastewater irrigated soil near Bindal River, Dehradun. *Agricultural Research*, 1(2), 157-164. DOI 10.1007/s40003-012-0016-8.
- Hussain, A., Priyadarshi, M., & Dubey, S. (2019). Experimental study on accumulation of heavy metals in vegetables irrigated with treated wastewater. *Applied Water Science*, 9(5), 1-11. <https://doi.org/10.1007/s13201-019-0999-4>.
- Khan, Z. I., Ahmad, K., Batool, F., Wajid, K., Mehmood, N., Ashfaq, A., ... & Ullah, S. (2019). Evaluation of toxic potential of metals in wheat crop grown in wastewater-contaminated soil in Punjab, Pakistan. *Environmental Science and Pollution Research*, 26(24), 24958-24966. <https://doi.org/10.1007/s11356-019-05715-9>.
- Mahmood, A., Mahmood, A., Malik, R. N., & Shinwari, Z. K. (2013). Indigenous knowledge of medicinal plants from Gujranwala district, Pakistan. *Journal of Ethnopharmacology*, 148(2), 714-723. <https://doi.org/10.1016/j.jep.2013.05.035>.

Mahmood, A., & Malik, R. N. (2014). Human health risk assessment of heavy metals via consumption of contaminated vegetables collected from different irrigation sources in Lahore, Pakistan. *Arabian Journal of Chemistry*, 7(1), 91-99. <https://doi.org/10.1016/j.arabjc.2013.07.002>.

Mahmood, A., Eqan, M., Pervez, S., Alghamdi, H. A., Tabinda, A. B., Yasar, A., ... & Pugazhendhi, A. (2020). COVID-19 and frequent use of hand sanitizers; human health and environmental hazards by exposure pathways. *Science of the Total Environment*, 742, 140561. <https://doi.org/10.1016/j.scitotenv.2020.140561>.

Mehmood, A., Mirza, M.A., Choudhary, M.A., Kim, K.H., Raza, W., Raza, N., Lee, S.S., Zhang, M., Lee, J.H., and Sarfraz, M. (2019). Spatial distribution of heavy metals in crops in a wastewater irrigated zone and health risk assessment. *Environmental Research*, 168, 382-388. <https://doi.org/10.1016/j.envres.2018.09.020>.

Qayoom, U., Bhat, S. U., Ahmad, I., & Kumar, A. (2021). Assessment of potential risks of heavy metals from wastewater treatment plants of Srinagar city, Kashmir. *International Journal of Environmental Science and Technology*, 1-20. <https://doi.org/10.1007/s13762-021-03612-8>.

Rani, M., Ullah, R., Alwahibi, M.S., Elshikh, M.S., Mahmood, A. (2021). Health Risk Assessment by Toxic Metals in Little Egrets (*Egretta garzetta*) and Food Chain Contaminations. *Saudi Journal of Biological Sciences*. <https://doi.org/10.1016/j.sjbs.2021.08.106>.

Singh, A. & Raj, P. (2018). Segregation of waste at source reduces the environmental hazards of municipal solid waste in Patna, India. *Archives of Environmental Protection*, 44(4), pp. 96–110. DOI: 10.24425/aep.2018.122306.

- Spognardi, S., Bravo, I., Beni, C., Menegoni, P., Pietrelli, L., & Papetti, P. (2019). Arsenic accumulation in edible vegetables and health risk reduction by groundwater treatment using an adsorption process. *Environmental Science and Pollution Research*, 26(31), 32505-32516. <https://doi.org/10.1007/s11356-019-06396-0>.
- Sun, J., Xu, W., Yang, P., Li, N., Yuan, Y., Zhang, H., ... & Chen, K. (2019). Enhanced oxytetracycline removal coupling with increased power generation using a self-sustained photo-bioelectrochemical fuel cell. *Chemosphere*, 221, 21-29. <https://doi.org/10.1016/j.chemosphere.2018.12.152>.
- Tabassum, R. A., Shahid, M., Dumat, C., Niazi, N. K., Khalid, S., Shah, N. S., ... & Khalid, S. (2019). Health risk assessment of drinking arsenic-containing groundwater in Hasilpur, Pakistan: effect of sampling area, depth, and source. *Environmental Science and Pollution Research*, 26(20), 20018-20029. <https://doi.org/10.1007/s11356-018-1276-z>.
- Rehman, K., Bukhari, S. M., Andleeb, S., Mahmood, A., Erinle, K. O., Naeem, M. M., & Imran, Q. (2019). Ecological risk assessment of heavy metals in vegetables irrigated with groundwater and wastewater: the particular case of Sahiwal district in Pakistan. *Agricultural Water Management*, 226, 105816. <https://doi.org/10.1016/j.agwat.2019.105816>
- USEPA. (2006). United States, Environmental Protection Agency, Integrated Risk Information System. <<http://www.epa.gov/iris/substS>>.
- WB-SCEA. (2006). Pakistan Strategic Country Environmental Assessment, Main report., Report no. 36946-PK, Volume I, World Bank. 1-66.

WHO. (1996). Trace elements in human nutrition and health. World Health Organization. Geneva.

WWF. (2007). Pakistan's Water at Risk: Water & Health Related Issues in Pakistan & Key Recommendations. Lahore: WWF Pakistan. From <http://www.wwfpak.org/pdf/water-report>. Pdf.

Zajda, M., & Aleksander-Kwaterczak, U. (2019). Wastewater treatment methods for effluents from the confectionery industry—an overview. *Journal of Ecological Engineering*, 20(9). DOI 10.12911/22998993/112557.

Zwolak, A., Sarzyńska, M., Szyrka, E., & Stawarczyk, K. (2019). Sources of soil pollution by heavy metals and their accumulation in vegetables: A review. *Water, air, & soil pollution*, 230(7), 1-9. <https://doi.org/10.1007/s11270-019-4221-y>.

Graphical Abstract




Figure 1. Sampling site map, Faisalabad, Pakistan




Table 1. Vegetables collected for this study

English name	Local name	Family	Plant species	Part sampled
Potato	Aloo	Solanaceae	<i>Solanum tuberosum L.</i>	Tubers
Mustard Leaf	Saag	Amaranthaceae	<i>Spinacia oleracea L.</i>	Leaves
Cabbage	Band Gobi	Brassicaceae	<i>Brassica oleracea</i>	Leaves
Carrot	Gajr	Apiaceae	<i>Daucus carota L.</i>	Underground stem
Cauliflower	Phool Gobi	Brassicaceae	<i>Brassica oleracea</i>	Fruiting flower

Table 2. Heavy metals concentrations in water (mg L⁻¹) and soil (mg kg⁻¹) irrigated with different water sources

Sample	Value	Pb	Mn	Cd	Ni	Zn	Cu
<i>Water</i>							
Wastewater	Mean ± S.D	2.30±.84.00	19.10±5.10	0.11±0.02	0.02±0.01	0.06±0.02	0.09±0.01
	Range	1.30-3.50	12.70-25.50	0.09-0.15	0.01-0.04	0.03-0.08	0.07-0.10
Groundwater	Mean ± S.D	0.04±0.03	0.09±0.01	0.002±0.001	0.004±0.002	1.10±0.23	0.02±0.01
	Range	0.02-	0.07-0.10	< 0.01	< 0.01	0.90-1.50	0.01-0.03
<i>WHO Standard</i>		5	0.2	0.1	0.2	2	Not Listed
<i>Soil</i>							
Wastewater	Mean ± S.D	26.80±8.00	25.40±4.60	0.80±0.30	1.70±0.50	5.14±0.38	2.30±0.60
	Range	19.00-39.50	19.00-31.00	0.30-1.10	1.10-2.10	4.70-5.70	1.75-3.35
Groundwater	Mean ± S.D	0.11±0.04	1.00±0.15	0.01±0.01	0.20±0.08	0.74±0.12	0.70±0.20
	Range	0.08-0.09	0.80-1.20	0.01-0.03	0.10-0.30	0.60-0.92	0.30-0.90
<i>WHO standard</i>		84	80	3	75	300	Not Listed

Table 3. Heavy metals concentrations (mg kg⁻¹) in vegetables grown with groundwater and wastewater

Vegetables	Value	Pb	Cd	Cu	Zn	Mn	Ni
<i>Groundwater</i>							
Potato	Mean ± S.D	0.010±0.002	0.003±0.003	0.046±0.033	0.010±0.003	2±0.8	0.020±0.005
	Range	0.007-0.013	0.001-0.009	0.010-0.090	0.009-0.016	0.96-3	0.016-0.027
Mustard Leaf	Mean ± S.D	0.060±0.200	0.003±0.001	0.414±0.300	0.200±0.070	14.2±4.1	0.080±0.010
	Range	0.034-0.083	0.002-0.005	0.170-0.800	0.120-0.290	9-19	0.059-0.090
Cabbage	Mean ± S.D	0.010±0.004	0.001±0.000	0.520±0.280	0.020±0.010	10.6±1.5	0.020±0.010
	Range	0.009-0.019	< 0.001	0.200-0.900	0.010-0.031	9-13	0.015-0.031
Carrot	Mean ± S.D	0.010±0.002	0.001±0.000	0.032±0.010	0.020±0.010	2.1±0.8	0.020±0.002
	Range	0.010-0.017	0.001-0.002	0.010-0.050	0.011-0.029	1.1-3	0.012-0.019
Cauliflower	Mean ± S.D	0.030±0.020	0.002±0.005	0.031±0.020	0.150±0.030	2.4±1	0.050±0.010
	Range	0.010-0.053	0.002-0.003	0.012-0.070	0.110-0.190	1.5-4	0.039-0.070
<i>Wastewater</i>							
Potato	Mean ± S.D	2.60±0.58	0.002±0.001	0.25±0.05	2.20±1.30	7.26±1.62	0.70±0.40
	Range	1.90-3.20	0.001-0.003	0.19-0.30	1.00-4.00	5.20-9.20	0.29-1.15
Mustard Leaf	Mean ± S.D	24.20±4.30	0.10±0.05	1.70±0.40	12.80±1.00	32.80±5.03	3.20±0.40
	Range	19.00-30.00	0.10-0.20	1.10-2.30	11.50-14.00	28.50-41.2	2.70-3.80
Cabbage	Mean ± S.D	24.00±2.80	0.05±0.02	1.60±0.30	2.60±0.70	20.60±3.40	2.60±0.50
	Range	21.00-27.00	0.03-0.09	1.10-1.90	1.75-3.50	17.00-25.20	2.10-3.40
Carrot	Mean ± S.D	3.20±1.00	0.002±0.001	0.09±0.01	1.90±0.70	7.00±2.70	1.20±0.40
	Range	2.10-4.70	0.001-0.003	0.09-0.12	1.00-2.90	3.50-10.50	0.70-1.70
Cauliflower	Mean ± S.D	4.50±1.50	0.004±0.002	0.50±0.20	7.70±1.90	5.50±1.70	0.90±0.30
	Range	2.70-6.50	0.002-0.009	0.20-0.80	5.50-10.30	3.50-7.70	0.50-1.30
Standard	WHO	0.05	0.02	40	50	16.61	10

Table 4. Heavy metal transfer factor (MTF) in vegetables grown with groundwater and wastewater

Vegetables	Pb	Cd	Cu	Zn	Mn	Ni
<i>Groundwater</i>						
Potato	0.091	1.950	0.135	0.119	0.016	0.066
Mustard Leaf	0.540	14.200	0.169	0.418	0.281	0.591
Cabbage	0.120	10.600	0.048	0.128	0.029	0.743
Carrot	0.113	2.100	0.056	0.092	0.028	0.046
Cauliflower	0.295	2.430	0.116	0.304	0.208	0.045
<i>Wastewater</i>						
Potato	0.096	0.286	0.002	0.439	0.428	0.109
Mustard Leaf	0.903	1.294	0.167	1.928	2.482	0.744
Cabbage	0.896	0.809	0.060	1.590	0.504	0.692
Carrot	0.121	0.276	0.002	0.699	0.377	0.043
Cauliflower	0.166	0.217	0.006	0.518	1.504	0.231

Table 5. Daily intake of metals (DIM, mg person⁻¹ day⁻¹) for vegetables grown with groundwater and wastewater

Vegetables	Pb	Cd	Cu	Zn	Mn	Ni
<i>Groundwater</i>						
Potato	1.00E-05	1.00E-03	1.00E-06	1.00E-05	1.00E-05	2.00E-05
Mustard Leaf	3.00E-05	7.40E-03	2.00E-06	4.00E-05	1.00E-04	2.00E-04
Cabbage	1.00E-05	5.50E-03	0.00E+00	1.00E-05	1.00E-05	3.00E-04
Carrot	1.00E-05	1.10E-03	1.00E-06	1.00E-05	1.00E-05	2.00E-05
Cauliflower	2.00E-05	1.30E-03	1.00E-06	3.00E-05	8.00E-05	2.00E-05
<i>Wastewater</i>						
Potato	1.00E-03	3.80E-03	1.00E-06	4.00E-04	1.10E-03	1.00E-04
Mustard Leaf	1.30E-02	1.70E-02	1.00E-04	1.70E-03	6.60E-03	9.00E-04
Cabbage	1.20E-02	1.07E-02	3.00E-05	1.40E-03	1.30E-03	8.00E-04
Carrot	2.00E-03	3.60E-03	1.00E-06	6.00E-04	1.00E-03	1.00E-04
Cauliflower	2.00E-03	2.90E-03	2.00E-06	4.00E-04	4.00E-03	3.00E-04

Research Paper (HM Risk Assessment)

ORIGINALITY REPORT

13%

SIMILARITY INDEX

7%

INTERNET SOURCES

11%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | Xingming Wang, Chuncai Zhou, Guijian Liu, Zhongbing Dong. "Transfer of Metals from Soil to Crops in an Area near a Coal Gangue Pile in the Guqiao Coal Mine, China", <i>Analytical Letters</i> , 2013
Publication | 1% |
| 2 | Yanglong Li, Chaoqun Wang, Chaowu Yan, Shaowen Liu, Xiangteng Chen, Mansheng Zeng, Yuhong Dong, Ruzhen Jiao. "Heavy metals concentrations and accumulation characteristics of dominant woody plants in iron and lead-zinc mining areas in Jiangxi, Southeast China", <i>Research Square Platform LLC</i> , 2022
Publication | 1% |
| 3 | assets.researchsquare.com
Internet Source | 1% |
| 4 | acikerisim.ozal.edu.tr
Internet Source | <1% |
| 5 | researchportal.northumbria.ac.uk
Internet Source | <1% |
-

6

www.researchgate.net

Internet Source

<1 %

7

Soil Biology, 2015.

Publication

<1 %

8

Razieh Khalilzadeh, Alireza Pirzad, Ebrahim Sepehr, Shahbaz Khan, Sumera Anwar. "The *Salicornia europaea* potential for phytoremediation of heavy metals in the soils under different times of wastewater irrigation in northwestern Iran", *Environmental Science and Pollution Research*, 2021

Publication

<1 %

9

Said Akbar, Zeshan Ali, Sadam Hussain, Ashiq Mohammad et al. "Metal accumulation potential, human health risks, and yield attributes of hundred bread wheat genotypes on irrigation with municipal and remediated wastewater", *Environmental Science and Pollution Research*, 2021

Publication

<1 %

10

Zeshan Ali, Abdul Mujeeb-Kazi, Umar Masood Quraishi, Riffat Naseem Malik. "Deciphering adverse effects of heavy metals on diverse wheat germplasm on irrigation with urban wastewater of mixed municipal-industrial origin", *Environmental Science and Pollution Research*, 2018

Publication

<1 %

11	Submitted to Lincoln University Student Paper	<1 %
12	academic.oup.com Internet Source	<1 %
13	Submitted to Rutgers University, New Brunswick Student Paper	<1 %
14	www.watconman.org Internet Source	<1 %
15	ojs.gjournals.org Internet Source	<1 %
16	Dalia A. Ahmed, Dalia F. Slima. "Heavy metal accumulation by Corchorus olitorius L. irrigated with wastewater", Environmental Science and Pollution Research, 2018 Publication	<1 %
17	Hadi Eslami, Abbas Esmaeili, Mohsen Razaeeian, Mahnaz Salari, Abdolreza Nassab Hosseini, Mohammad Mobini, Ali Barani. "Potentially toxic metal concentration, spatial distribution, and health risk assessment in drinking groundwater resources of southeast Iran", Geoscience Frontiers, 2022 Publication	<1 %
18	Henry A. Cordoba-Novoa, Jeimmy Cáceres-Zambrano, Esperanza Torres-Rojas.	<1 %

"Assessment of native cadmium-tolerant bacteria in cacao (*Theobroma cacao* L.) - cultivated soils in Cundinamarca-Colombia",
Research Square Platform LLC, 2022

Publication

19

Meena Sehgal, Ankur Garg, R. Suresh, Priya Dagar. "Heavy metal contamination in the Delhi segment of Yamuna basin",
Environmental Monitoring and Assessment, 2011

Publication

20

Ouafae El Hamiani, Hicham El Khalil, Catherine Sirguy, Ahmed Ouhammou, Gabriel Bitton, Christophe Schwartz, Ali Boularbah. "Metal Concentrations in Plants from Mining Areas in South Morocco: Health Risks Assessment of Consumption of Edible and Aromatic Plants",
CLEAN - Soil, Air, Water, 2015

Publication

21

Zhongxiao Fu, Taitea Dykstra, Morgan Wall, Huiping Li et al. "GABAergic neuronal IL-4R mediates T cell effect on memory",
Cold Spring Harbor Laboratory, 2021

Publication

22

jest.srbiau.ac.ir

Internet Source

23

pr.hec.gov.pk

Internet Source

<1 %

<1 %

<1 %

<1 %

<1 %

- | | | |
|----|--|------|
| 24 | www.aesacademy.org
Internet Source | <1 % |
| 25 | www.journalcjast.com
Internet Source | <1 % |
| 26 | www.nexusacademicpublishers.com
Internet Source | <1 % |
| 27 | www.thepab.org
Internet Source | <1 % |
| 28 | www.ujecology.com
Internet Source | <1 % |
| 29 | J. R. V. PILS, A. D. KARATHANASIS, T. G. MUELLER. "Concentration and Distribution of Six Trace Metals in Northern Kentucky Soils", Soil and Sediment Contamination: An International Journal, 2010
Publication | <1 % |
| 30 | Kifayatullah Khan, Yonglong Lu, Hizbullah Khan, Muhammad Ishtiaq, Sardar Khan, Muhammad Waqas, Luo Wei, Tieyu Wang. "Heavy metals in agricultural soils and crops and their health risks in Swat District, northern Pakistan", Food and Chemical Toxicology, 2013
Publication | <1 % |
| 31 | Md. Monirul Islam, Nusrat Jahan Avha, Shamim Ahmed, Md. Ahedul Akbor et al. | <1 % |

"Trace Metals and Organochlorine Pesticide Residues in Imported Fishes in Bangladesh and Human Health Risk Implications",
Research Square Platform LLC, 2021

Publication

32

Muhammad Usman Khan, Adeel Mahmood, Riffat Naseem Malik. "Assessment of Heavy Metals in Pharmacological Important Medicinal Plants Consumed in the Bannu District, Pakistan", Human and Ecological Risk Assessment: An International Journal, 2014

Publication

<1 %

33

Shumaila Majeed, Sajid Rashid, Abdul Qadir, Colin Mackay, Faisal Hayat. "Spatial patterns of pollutants in water of metropolitan drain in Lahore, Pakistan, using multivariate statistical techniques", Environmental Monitoring and Assessment, 2018

Publication

<1 %

34

Xiuying Zhang, Dongmei Chen, Taiyang Zhong, Xiaomin Zhang, Min Cheng, Xinhui Li. "Assessment of cadmium (Cd) concentration in arable soil in China", Environmental Science and Pollution Research, 2014

Publication

<1 %

35

full.escipub.org
Internet Source

<1 %

jreh.mums.ac.ir

36

Internet Source

<1 %

37

www.cjees.ro

Internet Source

<1 %

38

www.epa.gov

Internet Source

<1 %

39

Adeel Mahmood, Maryam Eqan, Saher Pervez, Amtul Bari Tabinda, Abdullah Yasar, Kathirvel Brindhadevi, Arivalagan Pugazhendhi. "COVID-19 and frequent use of hand sanitizers; human health and environmental hazards by exposure pathways", *Science of The Total Environment*, 2020

Publication

<1 %

40

Sher Ali Khan, Xiaoyu Liu, Bakht Ramin Shah, Wenting Fan, Hong Li, Sher Bahadhar Khan, Zahoor Ahmad. "Metals Uptake by Wastewater Irrigated Vegetables and their Daily Dietary Intake in Peshawar, Pakistan / Pobieranie Metali Przez Warzywa Nawadniane Ściekami I Ich Dienne Stężenie W Diecie Ludności Peszawaru, Pakistan", *Ecological Chemistry and Engineering S*, 2015

Publication

<1 %

41

A. Baysal, H. Saygin, G. S. Ustabasi. "Risks of graphene nanomaterial contamination in the

<1 %

soil: evaluation of major ions", Environmental Monitoring and Assessment, 2020

Publication

42

Bin Liang, Hao-Yi Cheng, De-Yong Kong, Shu-Hong Gao et al. "Accelerated Reduction of Chlorinated Nitroaromatic Antibiotic Chloramphenicol by Biocathode", Environmental Science & Technology, 2013

Publication

<1 %

43

Minhaz Ahmed, Masaru Matsumoto, Akinori Ozaki, Nguyen Thinh, Kiyoshi Kurosawa. "Heavy Metal Contamination of Irrigation Water, Soil, and Vegetables and the Difference between Dry and Wet Seasons Near a Multi-Industry Zone in Bangladesh", Water, 2019

Publication

<1 %

44

Muhammad Haroon, Abdulaziz A. Al-Saadi, Muhammad Adnan Iqbal. "Comparative Exposure Assessment of Potential Health Risks through the Consumption of Vegetables Irrigated by Freshwater/Wastewater: Gujranwala, Pakistan", Chemical Research in Toxicology, 2021

Publication

<1 %

45

S. Mounicou, J. Szpunar, D. Andrey, C. Blake, R. Lobinski. "Concentrations and bioavailability of cadmium and lead in cocoa

<1 %

powder and related products", Food Additives and Contaminants, 2003

Publication

46

Shammi, Mashura, Mohammad Abul Kashem, Md. Mostafizur Rahman, Md. Delwar Hossain, Rashadur Rahman, and M. Khabir Uddin. "Health risk assessment of textile effluent reuses as irrigation water in leafy vegetable *Basella alba*", International Journal Of Recycling of Organic Waste in Agriculture, 2016.

Publication

<1 %

47

Zafar Iqbal Khan, Kafeel Ahmad, Fauzia Batool, Kinza Wajid et al. "Evaluation of toxic potential of metals in wheat crop grown in wastewater-contaminated soil in Punjab, Pakistan", Environmental Science and Pollution Research, 2019

Publication

<1 %

Exclude quotes On

Exclude matches < 4 words

Exclude bibliography On

Research Paper (HM Risk Assessment)

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14

PAGE 15

PAGE 16

PAGE 17

PAGE 18

PAGE 19

PAGE 20

