

The Effect of Aloin in Blood Glucose and Antioxidants in Male Albino Rats with Streptozotocin-induced Diabetic

By Noorah Al-Sowayan

The Effect of Aloin in Blood Glucose and Antioxidants in Male Albino Rats with Streptozotocin-induced Diabetic

Reema Mohammad AL-Sallali¹, Noorah Saleh Al-Sowayan^{2*}

3

¹ Master student and Biology department, College of Science, Qassim University.

^{2*} Department of Biology, College of Science, Qassim University, P.O. Box 30230, Buraydah (51477), Saudi Arabia,

nsaoiean@qu.edu.sa, knaaj1@yahoo.com

* Corresponding authors

Abstract

The anti-diabetic effect of aloin was examined in vivo in the current study, similar to the study, which for four weeks examined how aloin affected the blood sugar, insulin, and antioxidant levels in both healthy and diabetic rats generated by streptozotocin. Adult male albino rats were experimentally given a single intraperitoneal injection of 60 mg/kg of streptozotocin to induced diabetes. The in vitro experiment utilized blood samples, and adult male albino rats' kidneys, pancreas and liver were separated. In comparison to control diabetic rats, aloin isolates (30 mg / kg) body weight for 30 days considerably decreased serum glucose and effectively boosted serum insulin levels. Liver and kidney congeners of malondialdehyde (MDA) were substantially decreased while, Catalase (CAT), superoxide dismutase (SOD) and Congeners Glutathione (GSH) were significantly boosted as contrasted to diabetic rats after aloin therapy. Moreover, a histological assessment of the pancreas was performed.

16

These results show the promise of aloin at a dose of 30 mg/kg body weight for 30 days as an antidiabetic and antioxidant agent.

Keyword

Diabetes mellitus (DM); *Aloe vera L.*; *Liliaceae*; Aloin; Glutathione (GSH); Malondialdehyde (MDA); Catalase (CAT); Superoxide Dismutase (SOD)

6

1. Introduction

Diabetes mellitus (DM), a group of metabolic disorders with several different etiologies that are characterized by elevated serum glucose levels or persistent hyperglycemia and abnormalities of insulin hormone secretion, insulin hormone resistance, or both, is one of the biggest threats to human health in the world today (Khan, 2021). Diabetes-related metabolic disorders increase the formation of mitochondrial superoxide in the heart

muscle and endothelial cells of either small and large blood arteries (Kelkar, Aditya, Jai, Mehta, & Amoaku, 2018).

Corrected :Commented [J1]

21

Consequently, oxidative stress is a crucial factor in the emergence of microvascular and cardiovascular problems associated with diabetes mellitus. (Tabatabaei S. R., Ghaderi, Tapehebur, Farbood, & Rashno, 2017; Sha, Hu, & Bu, 2020). Studies also show available treatments for controlling diabetes and its complications have various side effects with limited benefits; therefore, the desire for herbal products with anti-diabetic properties and fewer side effects that are more cost-effective furthermore is increasing (Association, 2018). Aloin (*Aloe vera L., Liliaceae family*), extracted from the outer bark of the *aloe vera* plant, has diverse applications in medicinal products and cosmetics, while research on animals has shown that aloin can be toxic, there is little indication of this in humans. If consumed excessively, especially over a lengthy period, aloin can be toxic, which signifies that it began having adverse effects on the body that may even be life-threatening. (Zhang, Liu, Liu, Zhao, & Tian, 2016; Association, 2018). Recent approaches suggest that the treatment of diabetes should not only focus on insulin secretion but also antioxidant protection of the β -cell in the pancreas, this may facilitate the repair of β -cells undergoing damage by oxidative stress secondary to hyperglycemia (Tangvarasittichai, 2015; Xu, Li, & Li, 2017) the objective of the study was to investigate how aloin affected blood sugar, insulin, and antioxidants to healthy rats, as well as rats with streptozotocin-induced diabetes, for four weeks.

Corrected :Commented [J2]

2. Material and Methods

I. Chemicals

Streptozotocin was supplied by Alfa Aesar USA and Aloin was supplied by Sigma Aldrich Co USA., Catalase (CAT), Superoxide dismutase activity (SOD), Kit Malondialdehyde (MDA) and Glutathione (GSH), and were also purchased from the NWLSS™.

22 Experimental animals

The College of Pharmacy at King Saud University provided 26 male Wistar rats for this investigation, each weighing 200–250 g. The rats were kept in polyacrylate cages with constant conditions and typical light/dark cycles. Animals were left for one month before experimentation to adapt to laboratory conditions. Throughout the experiment's acclimation phase and duration, a commercial balanced diet (General Organization for Grain Silos and Flour Mills, Riyadh) and water were provided ad libitum.

III. Induction of DM and study design

Rats were given intraperitoneal injections of streptozotocin after being starved for the previous night (Alfa Aesar, USA) (60 mg/kg). Throughout the first twenty-four hours following the administration of streptozotocin, mice treated with STZ were given a 5% glucose solution rather than water to prevent hypoglycemia shock-related mortality. Fasting blood glucose was measured seventy-two hours later utilizing a One Touch

glucometer (Icare TD-4279); diabetic rats were defined as those with fasting blood glucose levels greater than 250 mg/dL. (Mestry, Dhodi, Kumbhar, & Juvekar, 2017; Elekofehinti, Onunkun, & Olaleye, 2020)

Six rats each were given to the four groups,

Group 1: was normal control rats given an aqueous solution,

Group 2: The diabetic control rats were given an aqueous solution daily for 30 days,

Group 3: The normal control rats were given an aqueous solution daily for 30 days, and

Group 4: The diabetic rats were administered Aloin (30 mg/kg) in an aqueous solution daily for 30 days.

IV. Sampling

In the first, second, third, and fourth weeks, a blood glucose meter (Icare TD-4279) was utilized to check the blood sugar level in the tail vein.

By the completion of the experiment phase (day 30th), The retro-orbital plexus was used to collect blood samples, and serum samples subsequently separated using centrifugation. (Thermo Scientific, USA) and used immediately for Insulin hormone analyses (qing Su, diWang, & yanChi, 2017).

Post-sacrificed by anesthesia, the Liver, kidneys, and pancreas were harvested and rinsed in ice-cold saline, and calculation of liver, and kidney oxidative stress concentrations (MAD, GSH, SOD, CAT) (Sun, Wang, & Lu, 2017).

1. Determination of glucose

In this experience, we estimate blood glucose based on the use of the enzyme's method (Glucose oxidase) by the blood glucose meter (Icare TD-4279) (Rahmati, Keshvari, Mimasouri, & Chehelcheraghi, 2021)

2. Measurement of serum insulin concentration

Commercial quantitative enzyme-linked immunosorbent assay [(insulin ELISA kit) (DRG International, USA) (Jeffrey, Flier, Kahn, & Roth, 1979) was employed in a microplate reader to evaluate the optical density at an absorbance of 450 nm in order to evaluate the serum insulin levels of STZ-induced diabetic rats and also normal rats (Alborno, 2016).

V. Oxidative Stress Indicators

1. Sample Preparation

Animals were anaesthetized with chloroform and slaughtered at the conclusion of the procedure right after blood was collected. Livers and kidney tissues were quickly collected, collected organs were cleaned from fatty tissues and washed with Phosphate-buffered Heparin 0.16 mg/mL is introduced to saline (PBS), pH 7.4, to eliminate any red blood cells and clots. The samples were then homogenized in cold phosphate buffer using an ultrasonic homogenizer., the homogenate was centrifuged at 4000 rpm for 10 min at 4

°C in a refrigerated centrifuge and the supernatant was collected and kept at -20 °C and used as a sample for antioxidant assays (Al-musa & Al-hashem, 2013).

8 2. Determination of tissue samples

malondialdehyde (MDA), glutathione (GSH), superoxide dismutase (SOD) and catalase (CAT)

Tissue samples MDA was estimated by the method of Janero (1990). GSH was estimated by the method of Paglia DE, Valentine WN. (1967). Depending on the enzyme's capacity to prevent autoxidation using the pyrogallol methodology of Martin JP Jr, Dailey, and Sugarman, SOD activity was determined. Catalase was estimated by the method of BE S RF Jr, SIZER IW. (1952).

Kit Malondialdehyde (MDA), Glutathione (GSH), Superoxide dismutase activity (SOD), and Catalase (CAT) was used by the NWLSS™.

VI. Hematoxylin and eosin staining

Procedure: The pancreas tissues were dried in ethanol at increasing concentrations (70–100%) before being embedded in paraffin and fixed in a 10% neutral buffered formalin solution. Using a microtome, tissues were divided into sections that were 4 μm thick, mounted on glass slides, and afterwards dried slides. Following that, routine hematoxylin and eosin staining was performed on all samples. Thereafter, sections were examined at Olympus Optical in Tokyo, Japan, using a light microscope. All groups' slides were examined and captured on camera. (Alkudhayri, Osman, Alshammari, Al Maiman, & Yahya, 2021; Sarfraz, Khaliq, Hafizur, Raza, & Ullah, 2021)

19

VII. Statistical analysis

Data were expressed as the mean ± SEM. statistically analyzed using the SPSS program (version 23) and With the use of ANOVA and the Tukey multiple comparisons test, the various groups were compared. Significance was acceptable at P < 0.05.

* Significant variance unlike the standard control

^a significant variance unlike the standard control

3. Result

3.1. Effect of 30 days daily dose administration of Aloin on serum glucose and insulin levels.

Figures 1 and 2 depict the levels of serum-glucose as well as insulin in healthy and diabetic rats caused by streptozotocin, respectively. When contrasted to the matching control group, the serum glucose level was significantly elevated during diabetes, and the serum insulin level was significantly reduced. The current study demonstrated a significantly gradual decline in the serum glucose level. In the second week in the diabetic and Aloin group compared to the diabetic control group and was significantly lower versus normal control group in the 4th week, and the values of insulin levels and serum glucose tended to be close to normal in the aloin group.




Figure 1: Effect of Aloin on glucose level

* Significant different unlike the standard control
 a significant different unlike the standard control

ns $P < 0.05$




Figure 2: Effect of Aloin on insulin level

* Significantly variance unlike the standard control

a significantly variance unlike the diabetic control

ns $P < 0.05$

3.2. Effect of 30 days daily dose administration of aloin on liver and kidney tissue Malondialdehyde (MDA) levels

MDA level was significantly increased in the diabetic control group compared with the normal controls group, in the liver and kidney tissue re- respectively (Figure 3). With, MDA levels were decreased in the Diabetic + Aloin group contrast with the Diabetic control group. However, no significant ($P > 0.05$) difference in malondialdehyde level in the normal + Aloin group contrast to the normal control group.




Figure 3: Effect of Aloin on MDA

* Significant variance unlike the standard control

^a Significant variance unlike the diabetic control

^{ns} $P < 0.05$

3.3. Effect of 30 days daily dose administration of aloin on liver and kidney tissue Glutathione (GSH) levels

Liver and kidney tissue Glutathione activity were considerably lower in the Diabetic control group contrasted with the normal control group, respectively. GSH activity was significantly increased in the Diabetic + Aloin group contrast with the Diabetic control group. However no significant ($P > 0.05$) difference in GSH level in the normal + Aloin contrast to the usual control group (Figure 4).




Figure 4: Effect of Aloin on GSH

* Significant variance unlike the standard control

a Significant variance unlike the diabetic control

ns P < 0.05

3.4. Effect of 30 days daily dose administration of aloin on liver and kidney tissue Superoxide dismutase (SOD) levels

Liver and kidney tissue Superoxide dismutase activity was considerably increased in the Diabetic + Aloin group contrasted with the Diabetic control group, respectively. SOD activity was significantly lower in the Diabetic control group contrast with the normal control group. However no significant (P>0.05) difference in SOD level in the normal + Aloin group contrast to the usual control group (Figure 5).




Figure 5: Effect of Aloin ²³ SOD

* Significantly variance from the normal control.

a significantly variance from the diabetic control

ns $P < 0.05$

3.5. Effect of 30 days daily dose administration of aloin on liver and kidney tissue Catalase (CAT) levels ¹²

Liver and kidney tissue Catalase level was significantly lower in the Diabetic control group compared with the normal control group, ⁷ respectively. CAT level was significantly increased in the Diabetic+ Aloin group contrast with the Diabetic control group.

However ²⁵ significant ($P > 0.05$) difference in CAT level in the normal + Aloin group contrast to the usual control group (Figure 6).




Figure 6: Effect of Aloin on CAT

* Significantly variance unlike the standard control.

a significantly variance unlike the diabetic control

^{ns} $P < 0.05$

6. Histological results




Figure 7: Photomicrographs of the pancreases from the control (A&B) and control + Aloiin (30 mg/kg)-treated rats (C&D).

Both groups are showing normally sized circular islets of Langerhans intact α -cells (peripheral small cells with dark nuclei (small arrow) and β -cells (central larger with light nuclei (arrowhead)).




Figure 8: Photomicrographs of the pancreases from the diabetic rats (A and B) and diabetic + Aloin (30 mg/kg)-treated rats. In C and D:

There was a severe reduction in the size of the islets of the Langerhans, with obvious cytoplasmic denegation and a reduction in the number of both α and β cell (short arrow). Note the increase in inflammatory cell infiltration (long arrow). In C and D: there was a considerably increase in the size of the islets of the Langer-hans (long arrow) with an increase in the number of both α (short arrow) and β -cell (Arrowhead).

4. DISCUSSION

The current investigation's findings overwhelmingly point to aloin's potential as an anti-diabetic. When compared to diabetic controls, aloin-treated rats' in-vivo blood glucose and insulin levels were higher depicted in Figure 1 and Figure 2. (Abo-Youssef & Messiha 2013; Wexler, 2014). In conjunction, earlier writers stated that aloin had such an anti-diabetic effect in laboratory animals. (Patel & Patel, 2013)

There are many theories as to why aloin has an anti-diabetic effect.

The first theory is that Aloin's capacity to boost insulin sensitivity and encourage insulin production from the remaining β -cells may be the cause of these effects. (abatabaei, Ghaderi, Tapehebur, & Farbood, 2017).

The strong antioxidant properties of aloin extract provide the second justification. It has long been understood that aloin has antioxidant potential by reducing the production of free radicals and improving cellular thiol status. (Lee, Jeong, Chang Baek, Kwang Ku, & Sup Bae, 2019; Cichocka, Wietchy, Nabrdalik, & Gumprecht, 2016). Additionally, it is said to increase the activity of the enzyme glutathione-S-transferase. (Baruah, Bordoloi, & Baruah, 2016)

The quantitative determination of their metabolic end-product malondialdehyde (MDA), which is quantified in nmol/g utilizing the thio-barbituric acid test, is a straightforward and sensitive approach for estimating lipid peroxide concentration (TBA-test).

Malondialdehyde is a very hazardous consequence of lipid oxidation that is produced in part by free radicals. Its concentration is dramatically elevated in diabetes mellitus, according to numerous research. MDA combines with proteins and phospholipids in both irreversible and reversible ways, and it can have serious consequences [ManalKamal, MonaSalem, NaglaaKholousi, & KhadegaAshmawy, 2009]. MDA levels in the kidney and liver of all groups were measured. When comparing the control group of diabetics to the ordinary control group, we found a substantial rise in malondialdehyde (MDA). These results are consistent with those achieved by (Gina, Aulani, & Mahdi, 2016).

Our observation significantly confirmed aloin's ability to act as an antioxidant, since it was discovered to reduce excessive levels of malondialdehyde in tissue samples and to strengthen kidney and liver tissue. Glutathione, Superoxide dismutase, and Catalase levels (Gina, Aulani, & Mahdi, 2016). Recent strategies emphasize the significance of oxidative stress in pancreatic β -cells damage.

(Bajaj & Khan, 2012; Sumi, et al., 2019; ManalKamal, MonaSalem, NaglaaKholousi, & KhadegaAshmawy, 2009). The aetiology of diabetes mellitus involves oxidative stress, however antioxidants including aloin could have a real antidiabetic impact due to their antioxidant activity (Aldayel, Grace, & Lila, 2020; Ariani, Gofur, Listyorini, & Susanto, 2015).

Corrected :Commented [J3]

Corrected :Commented [J4]

Corrected :Commented [J5]

Conclusion

Antioxidant GSH, SOD, and CAT levels in the kidney and liver tissues of cured rats were significantly greater, whereas MDA levels were substantially lower after the program comparison with control rats consequently these data suggest that aloin may have promise as diabetes preventative and/or therapeutic agent, moreover, toxicological and clinical experiments are required to evaluate this plant extract and its possible applications in the medical and pharmaceutical fields.

5. References

Association, A. D. (2018, Mar 22). Economic Costs of Diabetes in the U.S. in 2017.

Retrieved from diabetesjournals.org ; 41(5):917–928: <https://doi.org/10.2337/dci180007>

Corrected :Commented [J6]

abatabaei, S. R., Ghaderi, S., Tapehebur, M. B., & Farbood, Y. (2017). Aloe vera gel improves behavioral deficits and oxidative status in streptozotocin-induced diabetic rats. *Biomedicine & Pharmacotherapy*, 96, 279-290. doi:<https://doi.org/10.1016/j.biopha.2017.09.146>

Abo-Youssef, A. M., & Messiha, B. A. (2013). Beneficial effects of Aloe vera in treatment of diabetes: Comparative in vivo and in vitro studies. *Bulletin of Faculty of Pharmacy, Cairo University*, 51(1), 7-11. Retrieved from <https://doi.org/10.1016/j.bfopcu.2012.03.002>

Alborno, N. T. (2016). *erum vaspin level in type 2 diabetic patients from gaza strip*. doi:<http://hdl.handle.net/20.500.12358/21587>

Aldayel, T. S., Grace, M. H., & Lila, M. A. (2020). LC-MS characterization of bioactive metabolites from two Yemeni Aloe spp. with antioxidant and antidiabetic properties. *Arabian Journal of Chemistry*, 13, 5040-5049. Retrieved from <https://doi.org/10.1016/j.arabjc.2020.02.003>

Khan, I. A. (2021, Aug 27). Do second generation sequencing techniques identify documented genetic markers for neonatal diabetes mellitus? Retrieved from cell.com; VOLUME 7, ISSUE 9: DOI:<https://doi.org/10.1016/j.heliyon.2021.e07903>

Corrected :Commented [J7]

Alkhudhayri, D. A., Osman, M. A., Alshammari, G. M., Al Maiman, S. A., & Yahya, M. A. (2021). Moringa peregrina leaf extracts produce anti-obesity, hypoglycemic, anti-hyperlipidemic, and hepatoprotective effects on high-fat diet fed rats. doi:<https://doi.org/10.1016/j.sjbs.2021.02.078>

Al-musa, h., & Al-hashem, f. (2013). hypoglycemic hepato renal and antioxidant potential effects of chamomile recutita flowers ethanolic extract in streptozotocin-diabetic rats. *American journal of pharmacology and toxicology*, 9(1), 1-12. doi:<http://thescipub.com/pdf/ajptsp.2014.1.12.pdf>

Ariani, N. L., Gofur, A., Listyorini, D., & Susanto, H. (2015, Sep). EFFECT OF Aloe vera GEL ON SUPEROXIDE DISMUTASE (SOD) LEVEL IN STREPTOZOTOCIN (STZ)-INDUCED DIABETIC

WISTAR Rattus novergicus LIVER. Retrieved from researchgate.net:
DOI:10.18502/klis.v2i1.116

Corrected :Commented [J8]

Bajaj , S., & Khan, A. (2012). Antioxidants and diabetes. *Indian J Endocrinol Metab*, 16(2), s267-s271. doi:10.4103/2230-8210.104057

Baruah, A., Bordoloi, M., & Baruah, H. (2016). Aloe vera: A multipurpose industrial crop. *Industrial Crops and Products*, 94, 951-963. doi:https://doi.org/10.1016/j.indcrop.2016.08.034

BEERS , R. J., & SIZER, I. A. (1952). spectrophotometric method for measuring the breakdown of hydrogen peroxide by catalase. *J Biol Chem*, 195(1), 133-40.

Cichočka, E., Wietchy, A., Nabrdalik, K., & Gumprecht, J. (2016). Insulin therapy - new directions of research. Retrieved from pubmed.ncbi.nlm.nih.gov; 67(3):314-24; DOI: 10.5603/EP.2016.0044

Corrected :Commented [J9]

Elekofehinti, O. O., Onunkun, A. T., & Olaleye, T. M. (2020). Cymbopogon citratus (DC.) Stapf mitigates ER-stress induced by streptozotocin in rats via down-regulation of GRP78 and up-regulation of Nrf2 signaling. 262. doi:https://doi.org/10.1016/j.jep.2020.113130Get rights and content

Gina, L. P., Aulani, A., & Mahdi, C. (2016, April). MDA and Histologic Profile of Pancreatic Diabetic-Rats Model Administered With Extract of Glycine max (L.) Merr. Retrieved from researchgate.net: DOI:10.21776/ub.jpacr.2016.005.01.226

Corrected :Commented [J10]

Hirano, T. (2018, July 12). Pathophysiology of Diabetic Dyslipidemia. Retrieved from jstage.jst.go.jp: https://doi.org/10.5551/jat.RV17023

Corrected :Commented [J11]

Janero, D. R. (1990). Malondialdehyde and thiobarbituric acid-reactivity as diagnostic indices of lipid peroxidation and peroxidative tissue injury. *Free Radical Biology and Medicine*, 9(6), 515-540. doi:https://doi.org/10.1016/0891-5849(90)90131-2

Jeffrey , S., Flier, C., Kahn, R., & Roth, J. (1979). Receptors, Antireceptor Antibodies and Mechanisms of Insulin Resistance. *The New England Journal of Medicine*. doi:10.1056/NEJM197902223000808

ManalKamal, MonaSalem, NaglaaKholousi, & KhadegaAshmawy. (2009, December). Evaluation of trace elements and Malondialdehyde levels in type II diabetes mellitus. Retrieved from sciencedirect.com; Volume 3, Issue 4, pp 214-218: <https://doi.org/10.1016/j.dsx.2009.07.007>

Corrected :Commented [J12]

Kelkar, Aditya, Jai, Mehta, & Amoaku. (2018, Oct). Cataract surgery in diabetes mellitus. Retrieved from journals.lww.com; 66(10):p 1401-1410: DOI: 10.4103/ijo.IJO_1158_17

Corrected :Commented [J13]

Lee, W., Jeong, G.-S., Chang Baek, M., Kwang Ku, S., & Sup Bae, J. (2019). Renal protective effects of aloin in a mouse model of sepsis. *Food and Chemical Toxicology*, 132. doi:<https://doi.org/10.1016/j.fct.2019.110651>

Martin Jr, J. P., Dailey, M., & Sugarman, E. (1987). Negative and positive assays of superoxide dismutase based on hematoxylin autoxidation. 255(2), 329-336. doi:[https://doi.org/10.1016/0003-9861\(87\)90400-0](https://doi.org/10.1016/0003-9861(87)90400-0)

Mestry, S. N., Dhodi, J. B., Kumbhar, S. B., & Juvekar, A. R. (2017). Attenuation of diabetic nephropathy in streptozotocin-induced diabetic rats by Punica granatum Linn. leaves extract. *Journal of Traditional and Complementary Medicine*, 7, 273-280. Retrieved from <https://doi.org/10.1016/j.jtcme.2016.06.008>

Paglia, D., & Valentine, W. (1967). Studies on the quantitative and qualitative characterization of erythrocyte glutathione peroxidase. doi:<https://pubmed.ncbi.nlm.nih.gov/6066618/>

Patel, K., & Patel, D. K. (2013). Medicinal importance, pharmacological activities, and analytical aspects of aloin: A concise report. *Journal of Acute Disease*, 2, 262-269. Retrieved from [https://doi.org/10.1016/S2221-6189\(13\)60141-9](https://doi.org/10.1016/S2221-6189(13)60141-9)

qing Su, L., diWang, Y., & yanChi, H. (2017). Effect of curcumin on glucose and lipid metabolism, FFAs and TNF- α in serum of type 2 diabetes mellitus rat models. *Saudi Journal of Biological Sciences*, 24(8), 1776-1780. doi:<https://doi.org/10.1016/j.sjbs.2017.11.011>

Rahmati, M., Keshvari, M., Mirasouri, R., & Chehelcheraghi, F. (2021). Exercise and *Urtica dioica* extract ameliorate hippocampal insulin signaling, oxidative stress, neuroinflammation, and cognitive function in STZ-induced diabetic rats. *Biomedicine & Pharmacotherapy*, 139. doi:<https://doi.org/10.1016/j.biopha.2021.111577>

Sarfraz, M., Khaliq, T., Hafizur, R., Raza, S., & Ullah, H. (2021). Effect of black pepper, turmeric and ajwa date on the endocrine pancreas of the experimentally induced diabetes in wister albino rats: A histological and immunohistochemical study. *Endocrine and Metabolic Science*, 4. doi:<https://doi.org/10.1016/j.endmts.2021.100098>

Sha, W., Hu, F., & Bu, S. (2020). Mitochondrial dysfunction and pancreatic islet β -cell failure (Review). *Exp ther med*, 20(6). doi:10.3892/etm.2020.9396

Sumi, F. A., Sikder, B., Rahman, M., Lubna, S. R., Ulla, A., Hossain, M., . . . Subhan, N. (2019). Phenolic Content Analysis of Aloe vera Gel and Evaluation of the Effect of Aloe Gel Supplementation on Oxidative Stress and Fibrosis in Isoprenaline-Administered Cardiac Damage in Rats. *Prev Nutr Food Sci*, 24(3), 254–264. doi:10.3746/pnf.2019.24.3.254

Sun, H., Wang, N., & Lu, Y. (2017). Lead Exposure Induces Weight Gain in Adult Rats, Accompanied by DNA Hypermethylation. *plos one*, 12(7). doi: 10.1371/journal.pone.0169958

Tabatabaei, S. R., Ghaderi, S., Tapehebur, M. B., Farbood, Y., & Rashno, M. (2017). Aloe vera gel improves behavioral deficits and oxidative status in streptozotocin-induced diabetic rats. *Biomedicine & Pharmacotherapy*, 96, 279-290. Retrieved from <https://doi.org/10.1016/j.biopha.2017.09.146>

Tangvarasittichai, S. (2015). Oxidative stress, insulin resistance, dyslipidemia and type 2 diabetes mellitus. *World journal of diabetes*, 6(3), 456-480. doi:10.4239/wjd.v6.i3.456

Wexler, P. (2014, Mar 11). *Encyclopedia of Toxicology*. Retrieved from [elsevier.com: https://www.elsevier.com/books/encyclopedia-of-toxicology/wexler/978-0-12-386454-3?original=true](https://www.elsevier.com/books/encyclopedia-of-toxicology/wexler/978-0-12-386454-3?original=true)

Xu, D.-P., Li, Y., & Li, H.-B. (2017). Natural Antioxidants in Foods and Medicinal Plants: Extraction, Assessment and Resources. *int J MOI Sci*, 18(1). doi: 10.3390/ijms18010096

Zhang, Y., Liu, W., Liu, D., Zhao, T., & Tian, H. (2016). Efficacy of Aloe Vera Supplementation on Prediabetes and Early Non-Treated Diabetic Patients: A

Corrected :Commented [J14]

Systematic Review and Meta-Analysis of Randomized Controlled Trials. 8(7), 388.
Retrieved from <https://doi.org/10.3390/nu8070388>

The Effect of Aloin in Blood Glucose and Antioxidants in Male Albino Rats with Streptozotocin-induced Diabetic

ORIGINALITY REPORT

18%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** Seyed Reza Fatemi Tabatabaei, Shahab Ghaderi, Mohammad Bahrami-Tapehebur, Yaghoob Farbood, Masome Rashno. "Aloe vera gel improves behavioral deficits and oxidative status in streptozotocin-induced diabetic rats", *Biomedicine & Pharmacotherapy*, 2017
Crossref 49 words — 2%
- 2** www.saepub.com
Internet 38 words — 1%
- 3** "Comparative study on The effect of two sweeteners stevia and sucrose on blood glucose levels in healthy studies", *Egyptian Journal of Chemistry and Environmental Health*, 2021
Crossref 33 words — 1%
- 4** scirp.org
Internet 33 words — 1%
- 5** worldwidescience.org
Internet 32 words — 1%
- 6** Imran Ali Khan. "Do second generation sequencing techniques identify documented genetic markers for neonatal diabetes mellitus?", *Heliyon*, 2021
Crossref 29 words — 1%

7	www.i-scholar.in Internet	26 words — 1%
8	www.researchgate.net Internet	25 words — 1%
9	www.medsci.org Internet	23 words — 1%
10	ir.ymlib.yonsei.ac.kr Internet	22 words — 1%
11	jeb.co.in Internet	20 words — 1%
12	www.mdpi.com Internet	15 words — 1%
13	Nurettin Aydogdu. "PROTECTIVE EFFECTS OF I-CARNITINE ON MYOGLOBINURIC ACUTE RENAL FAILURE IN RATS", <i>Clinical and Experimental Pharmacology and Physiology</i> , 1/2006 Crossref	14 words — 1%
14	Khaled B. Alharbi, H.M. Mousa, Z.H. Ibrahim, I.M. El-Ashmawy. "Hepatoprotective Effect of Methanolic Extracts of <i>Prosopis farcta</i> and <i>Lycium shawii</i> Against Carbon Tetrachloride-induced Hepatotoxicity in Rats", <i>Journal of Biological Sciences</i> , 2016 Crossref	13 words — < 1%
15	Mayank Choubey, Ashutosh Ranjan, Puran S. Bora, Amitabh Krishna. "Protective role of adiponectin against testicular impairment in high-fat diet/streptozotocin-induced type 2 diabetic mice", <i>Biochimie</i> , 2019 Crossref	12 words — < 1%

-
- 16 dspace.uzhnu.edu.ua 12 words — < 1%
Internet
-
- 17 Nageswara Rao Thalapaneni, Mohana Lakshmi Sabapathi, Faraz Raza Ansari, Subhash C. Mandal. "Antidiabetic and antioxidant effect of methanol extract of edible plant *Talinum portulacifolium* (Forssk) in Streptozotocin induced diabetic rats", *Oriental Pharmacy and Experimental Medicine*, 2011 11 words — < 1%
Crossref
-
- 18 eprints.mums.ac.ir 11 words — < 1%
Internet
-
- 19 journals.abu.edu.ng 11 words — < 1%
Internet
-
- 20 Eman Fawzy El Azab, Heba Sayed Mostafa. "Geraniol ameliorates the progression of high fat - diet/streptozotocin - induced type 2 diabetes mellitus in rats via regulation of caspase - 3, Bcl - 2, and Bax expression", *Journal of Food Biochemistry*, 2022 10 words — < 1%
Crossref
-
- 21 Manuela Cabiati, Agnese Sgalippa, Giovanni Federico, Silvia Del Ry. "C-type natriuretic peptide in childhood obesity", *Peptides*, 2021 10 words — < 1%
Crossref
-
- 22 Noorah Saleh Al-Sowayan. "Possible modulation of nervous tension-induced oxidative stress by vitamin E", *Saudi Journal of Biological Sciences*, 2020 10 words — < 1%
Crossref
-
- 23 cyberleninka.org 10 words — < 1%
Internet
-

24 spandidos-publications.com
Internet

10 words — < 1%

25 www.frontiersin.org
Internet

10 words — < 1%

EXCLUDE QUOTES OFF

EXCLUDE SOURCES < 5 WORDS

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 10 WORDS