

# JKSUS

*by Javed Iqbal*

---

**Submission date:** 24-Sep-2023 03:32AM (UTC-0700)

**Submission ID:** 2175003244

**File name:** d\_Investigation\_of\_Different\_In\_Vitro\_Biological\_Activities.docx (135.19K)

**Word count:** 4140

**Character count:** 22311

**Evaluation and Chemical Profiling of Different *Centaurea iberica* Extracts and  
Investigation of Different *In Vitro* Biological Activities**

**Haleema Bibi<sup>1</sup>, Javed Iqbal<sup>2\*</sup>, Banzeer Ahsan Abbasi<sup>3</sup>, Sobia Kanwal<sup>4\*</sup>, Maryam Tavafoghi<sup>5</sup>, Mohammad Z. Ahmed<sup>6</sup>, Tariq Mahmood<sup>1</sup>**

<sup>13</sup>  
1Department of Plant Sciences, Faculty of Biological Sciences, Quaid-i-Azam University Islamabad, 45320, Pakistan; Email: [hriaz@bs.qau.edu.pk](mailto:hriaz@bs.qau.edu.pk); [tmahmood@qau.edu.pk](mailto:tmahmood@qau.edu.pk)

<sup>3</sup>  
2Department of Botany, Bacha Khan University, Charsadda, 24420, Khyber Pakhtunkhwa, Pakistan; Email: [javed89qau@gmail.com](mailto:javed89qau@gmail.com)

3Department of Botany, Rawalpindi Women University, 6th Road, Satellite Town, Rawalpindi 46300, Pakistan; [banzeer.abbasi@f.rwu.edu.pk](mailto:banzeer.abbasi@f.rwu.edu.pk)

<sup>5</sup>  
4Department of Biology and Environmental Sciences, Allama Iqbal Open University, Islamabad; Email: [sobia.kanwal@aiou.edu.pk](mailto:sobia.kanwal@aiou.edu.pk)

<sup>3</sup>  
5Department of Bioengineering, University of California, Los Angeles, 410 Westwood Plaza, Los Angeles, CA 90095, USA; [Mary.tavafoghi@gmail.com](mailto:Mary.tavafoghi@gmail.com)

<sup>5</sup>  
6Department of Pharmacognosy, College of Pharmacy, King Saud University, Riyadh, Saudi Arabia; [mahmed4@ksu.edu.sa](mailto:mahmed4@ksu.edu.sa)

\*Correspondence: [javed89qau@gmail.com](mailto:javed89qau@gmail.com) (J.I); [Sobia.kanwal@aiou.edu.pk](mailto:Sobia.kanwal@aiou.edu.pk) (S.K)

**SHORT TITLE:** Evaluation and Chemical Profiling of Different *Centaurea iberica* Extracts

<sup>1</sup>  
**Declarations**

**Conflicts of interest:** The authors declare that they have no conflict of interest.

**Ethics approval:** Not Applicable

**Consent to participate:** All authors consent to participate in this manuscript

**Consent for publication:** All authors consent to publish this manuscript in Saudi Journal of Biological Science

**Data Availability Statement:** Data will be available on request to corresponding or first author

**Code availability:** Not Applicable

**AUTHOR CONTRIBUTIONS**

HB., and JI<sup>4</sup> generated the idea and performed research work. Supervision was provided by T.M. Experimental facilities were also provided by T.M. H.B., J.I. wrote the manuscript. B.A.A., M.T., MZA., and, S.K.<sup>8</sup> helped with software, manuscript editing and revision. All authors have read and agreed to the published version of the manuscript.

**Abstract:** In almost all cultures medicinal plants are used as a rich source of phytochemicals and nutraceuticals components and have played promising role in maintaining and improving human health. The current study was aimed to unveil biological evaluation of *Centaurea iberica*. Plant extracts were prepared in five different solvents viz. methanol, ethanol, n-hexane, ethyl acetate and chloroform. The antioxidant potentials (2,2-diphenyl, 1-picrylhydrazine scavenging activity, total reducing power) of the plant revealed that the methanolic extract showed significant DPPH (2,2-diphenyl, 1-picrylhydrazine) scavenging activity (IC<sub>50</sub> value 77.20 ± 8.81), reducing power (20.45 ± 0.54 AAE mg/g). In case of antibacterial assay, highest zone of inhibition (20 ± 1.5 mm) was recorded in methanolic extract against *Klebsiella pneumonia*. Likewise, for antifungal activity<sup>26</sup> the highest zone of inhibition was recorded for methanolic extract (18 ± 2.4 mm) against *Aspergillus niger*. Moreover, 50 % inhibition was observed for methanolic extract against targated PC3 (prostate cancer cell line) cell lines and 3T3 cell lines. In the light of present study, it is recommended that further<sup>27</sup> *in vitro* and *in vivo* studies should be performed using different animal models and characterization should be done to isolate different compounds that could be used to treat different types of ailments.

**Keywords:** Antibacterial, antifungal, anticancer, antioxidant, *Centaurea iberica*.

## 1. INTRODUCTION

Medicinal plants have strong therapeutic purposes and have been used since centuries for the preparation of functional drugs. Plants are in use as medicine for human beings at least 60,000 years back to the middle Paleolithic period (Zahra et al., 2022). The 21<sup>st</sup> century is considered as the century of biology, drove and powered by technological advances and scientific knowledge (Lenzner et al., 2019). Medicinal plant cultivation has started about 5000 years ago

in Egypt, India and China and approximately 2500 years ago in Greece and Central Asia (Jamshidi et al., 2018). Humans in the initial era slowly realized the importance of plants for different purposes including, food, shelter, clothing, fertilizers, flavors, fragrances (Anand et al., 2019). Medicinal plants are being used for the service of humanity and are very important, irrespective of the region and era throughout the globe. Plants were, are, and will remain beneficial ever for various purposes including, nutritional, cultural, social, religious, environmental, and human well-being especially for the people living in low-income countries like Pakistan, where people are facing poverty, poor health, malnutrition and unemployment (Ullah, 2010). Following the ancient time when plant-based medicines were used for the treatment, demand for herbal medicine and herbal products has increased gradually. For primary healthcare, approximately 80 % of the Asian and African population is dependent upon traditional medicine as herbal medicines are economical and are without side effects (Sahoo et al., 2010). Chinese, Egyptians and Greeks are considered the earliest people who used plants as medicine since more than centuries. The knowledge regarding medicinal plants has been transferred gradually from one generation to other and has been updated gradually with modern technologies and facilities (Schippmann et al., 2006). Because of quality, safety and effectiveness, herbal medicines became an important part of therapeutics in developing countries (Jamshidi et al., 2018). Medicinal plants have very important role in treating various fatal diseases. All parts of the plants contain various active compounds which can be used as a remedy for different diseases (Shinwari et al., 2017). Phytochemicals also known as bioactive nutrients or compounds which naturally found in plants which are released or produced by the plants for health benefits (Zahra *et al.*, 2021, Zahra *et al.*, 2022).

To check the medicinal importance of the plants, <sup>14</sup> present research study was conducted to analyze the phytochemical composition and biological potential of *C. iberica* (*Asteraceae*). This plant is native to District Chakwal, Pakistan and is frequently found in the northern areas

of Pakistan (Nasir and Sikander, 2006). Different chemical substances such as steroid, volatile constituents, flavones, terpenoids, fatty acids, lactones have been reported. *Centaurea* species had widely been part of traditional therapeutics for the treatment of various health issues including, digestive, gynecological and dermatological problems (Khammar and Djeddi, 2012). It is an important part of Turkish traditional medicine and have shown significant biological activities such as wound healing, ease the inflammation and pain in patients having rheumatoid arthritis, headache, and high fever and for quick wound healing (Koca et al., 2009, Khan et al., 2011). *Centaurea iberica* is a rich source of different phytochemicals (lignans anthocyanins, flavonoids, phenolic acids, and stilbenes), xanthophylls and carotenes and exhibited several medicinal properties including anti-cancer, cytotoxicity, anti-inflammatory, and antioxidant (Alper et al., 2021). The *Centaurea iberica* has also shown antifungal and anti-microbial potential due to the presence of high amount of sesquiterpene lactones (Aktumsek et al., 2013). <sup>9</sup> The current study was conducted to explore the phytochemical composition and biological potentials of asynthesized *C. iberica* mediated extracts.

<sup>20</sup>

## **2 Materials and Methods**

### *2.1. Plant collection and extract preparation*

The medicinal plant *Centaurea iberica* Trevir. ex Spreng. was collected in its flowering season from the mountainous region of Murree (33.9070° N, 73.3943° E) Pakistan with dominant surrounding flora *Acacia modesta*, *Diosypros lotus* etc) and was taxonomically identified consulting literature review and flora of Pakistan. The study was conducted at Plant Biochemistry and Molecular Biology lab, QAU Islamabad. After collection and identification, whole plant was thoroughly washed by running tap water and was shade dried. Next, the plant extract was grounded into fine powder. Further, five different solvents including two polar (ethanol and methanol) and three non-polar (ethyl acetate, n-hexane and chloroform) were used to prepare extracts. The extracts were prepared using 250 mL of each solvent. Further, extracts

were filtered and the filtrate was placed in fume hood for drying. After drying, the extracts were scratched and were shifted into eppendorf's tube and finally placed at 4°C. The mixtures were used for further quantitative and qualitative analysis.

### 2.2. DPPH free radical scavenging assay

A method which was outlined by Clarke et al. (2013) was used for DPPH determination with minor changes. Plant extract of 10  $\mu$ L and 190  $\mu$ L of DPPH was poured in microtiter plate to make a volume of 200  $\mu$ L. Ascorbic acid was used as positive and DMSO as negative control. Afterwards, the incubation of reaction mixture was done at room temperature in darkness for half an hour. Changes in the color of mixture indicates oxidation potential of plants. Finally, absorbance of the experimental sample was recorded at 517 nm using formula below.

$$\text{Percent scavenging of sample} = \frac{\text{Absorbance of control} - \text{Absorbance of sample} \times 100}{\text{Absorbance of control}}$$

### 2.3. Reducing power assay

Protocol by Ravisankar et al. (2014) was used to analyze the reducing power of different plant extracts. To achieve this purpose, 100  $\mu$ L of stock solution of each plant sample was taken and was added with 1 % solution of potassium ferricyanide (250  $\mu$ L) solution and 0.2 M phosphate buffer (200  $\mu$ L) solution. Incubation of these mixtures were done for 20 minutes at 50°C. For the acidification of solution, 200  $\mu$ L trichloroacetic acid was added. Centrifugation of the resultant mixtures was performed for 10 minutes at 300 rpm. Supernatant (150  $\mu$ L) was taken and 50  $\mu$ L (0.1 %) ferric chloride was added in it. Later on, the reaction mixtures of 200  $\mu$ L was shifted into 96 wells plate and absorbance was recorded at 630 nm. To determine the reducing power of the experimental sample Gallic acid (GA) was taken as positive control and reducing potential was recorded as GA equivalents mg/g of extract.

#### *2.4. Antibacterial activity*

Disc diffusion method performed by Razmavar <sup>30</sup> et al. (2014) was used to evaluate the antibacterial potential of the plant samples. For antibacterial assay, 100  $\mu$ L of each bacterial inoculum was smoothly spread on media using sterile cotton swabs. Different concentrations were pre-prepared as stock dilutions. Paper discs of 6 mm diameters were soaked with 25  $\mu$ L of the stock solution from each extract and were placed on petri plates containing bacterial inoculum. For positive control Oxytetracycline was used followed by incubation at 24 hr at 37 °C. Antibacterial activity of selected plant extracts was checked by calculating their respective MIC values.

#### *2.4. Antifungal activity*

For the assessment of antifungal potential of each plant sample disc diffusion method was done as described by Sharma et al. (2012). Sabouraud dextrose agar media (65 g) per 1000 mL of distilled H<sub>2</sub>O was used for the growth of fungal cultures and its pH was maintained to 6.5. Filter paper Whatman # 1 was used for the preparation of round discs of about 6 mm in size. The experiment was performed in sterilized conditions. Media was first autoclave and it was poured and solidified. Each sample solution of about 5  $\mu$ L along with 2  $\mu$ L of standard drug was loaded on filter discs and it was kept on the plated media (SDA). Parafilm was used to cover <sup>28</sup> the plates and were incubated for 24 hr at 28°C. The next day, clear zones of inhibition were recorded and measured by mean of Vernier caliper. Minimum inhibitory concentration (MIC) was observed.

#### *2.6. Anticancer activity on PC3 and 3T3 cell lines*

Prostate cancer (PC3) activity of the plant was detected by protocol outlined by Asadi-Samani et al. (2018) and for 3T3 cell lines method outlined by Mossam et al. (1983) was used. For evaluating cells viability, the anticancer assay was performed in 96 wells micro plates in which

MTT was used as a standard. The PC3 (Prostrate cancer) and 3T3 (Mouse fibroblast) cells were cultured using DMEM media, 5 % FBS, 100 IU/mL penicillin, 100 µg/mL of streptomycin was added in flask, and incubated in 5 % CO<sub>2</sub> incubator at 37 °C. After 24 hr of incubation, old media was removed and cells were replaced with fresh media (200 µL) containing different concentrations of plant extracts (ranges from 1-30 µM). After 48 hr, 200 µL of medium contains MTT (0.5 mg/ml) was loaded into each well and were incubated for 4 hr. Absorbance of the plates were recorded at 570 nm using 96 well plate to determine the (IC<sub>50</sub>) value of the experimental values against for PC3 cell lines and 3T3 cell lines.

### 2.7. Statistics

Each experiment was carried out in triplicate. The descriptive statistics were applied using MS excel (2016). Results of biological evaluation and antioxidant were subjected to Analysis of variance (ANOVA) and least significant difference (LSD), by using Statistics version 8.1. Graph pad prism version 5.01 was used for calculating IC<sub>50</sub> values.

## 3. RESULTS AND DISCUSSION

### 3.1. Antioxidant Assays.


It is one of the important biological activity which shows capacity of the extract to scavenge the free radical species which may cause harmful effects on body, if they are present in high concentrations inside the body. Oxidative stress caused by ROS is one of the main cause of various diseases. Plants contains natural antioxidant potential which may help in scavenging the ROS and thus can be used in therapeutic preparations. For the detection of antioxidant potentials, various assays were performed which includes DPPH, and total reducing power (TRP).

### 3.2. DPPH free radical scavenging assay.

Five different concentrations of plant extracts were used in free radical scavenging assay. It was noticed that CIE exhibited the lowest IC<sub>50</sub> value (82.01 ± 8.9 µg/mL). Ascorbic acid was taken as standard with IC<sub>50</sub> value (9.187 ± 0.473 µg/mL). Whereas, CIM showed value of 77.20 ± 8.81 µg /mL as shown in Table 1. The maximum IC<sub>50</sub> values (127.6 ± 11.6 µg /mL, 179.4 ± 2.27 µg /mL, 294.8 ± 2.97 µg/mL) were noted for CIA, CIC and CIH, respectively. The CIE exhibited dose dependent activity at 1000 µg/mL (75.10 %) and (49.01 %) at 62.5 µg/mL (Figure 1). For CIA the maximum scavenging activity were recorded at 1000 µg/mL (72.30 %) minimal activity (43.55 %) at 62.5µg/mL. The CIC scavenged 65.52 % DPPH free radical at 1000 µg/mL and 32.87 % at 62.5 µg/mL. CIH showed maximum scavenging effect (66.20 %) at 1000 µg/mL and minimum scavenging effect at 62.5 µg/mL (33.81 %). CIM (74.21 %) showed maximum % inhibition at 1000 µg/mL respectively and minimum % inhibition was recorded at 62.5 µg/mL (48.53 %).

**Table 1:** DPPH IC<sub>50</sub> (µg/mL) values of various extracts of *C. iberica*.


Sr. #	Plant Extract	IC <sub>50</sub> (µg/mL)
1	CIE	82.01 ± 8.9
2	CIA	127.6 ± 11.6
3	CIH	294.8 ± 2.97
4	CIC	179.4 ± 2.27
5	CIM	77.20 ± 8.81
6	Ascorbic Acid	9.187 ± 0.473



**Figure 1:** DPPH scavenging potential of *C. iberica* mediated extracts at various concentrations. **Key:** **CIE:** *C. iberica* ethanolic extract; **CIA:** *C. iberica* ethyl acetate extract; **CIH:** *C. iberica* n-hexane extract; **CIC:** *C. iberica* chloroform extract; **CIM:** *C. iberica* methanolic extract.

### 3.2. Total reducing power (TRP)

The TRP is also known as phospho-molybdenum assay. TRP was evaluated using equation obtained from Gallic acid calibration curve ( $y = 0.0004x + 0.2492$ ), and results were expressed as Gallic acid equivalent (GAE mg/g). The highest value was shown by CIM ( $20.45 \pm 1.6$  mg/g) followed by CIA ( $17.13 \pm 0.8$  mg/g), CIE ( $16.40 \pm 3.0$  mg/g), CIC ( $12.6 \pm 2.4$  mg/g) and the minimum value was recorded by CIH ( $8.9 \pm 2.4$  mg/g) as shown in the Figure 2.



**Figure 2:** Total reducing power of various extracts of *C. iberica*. **Key:** **CIE:** *C. iberica* ethanolic extract; **CIA:** *C. iberica* ethyl acetate extract; **CIH:** *C. iberica* n-hexane extract; **CIC:** *C. iberica* chloroform extract; **CIM:** *C. iberica* methanolic extract. Error bars = Standard error.

### 3.3. Antibacterial activity.

Antibacterial activity was performed on two extracts of *C. iberica* which includes one polar CIM (*C. iberica* methanolic extract) and one non polar CIH (*C. iberica* n-hexane extract) by using disc diffusion assay. For this, *S. aureus* was used as a gram-positive bacteria and *E. coli* and *K. pneumonia* were used as gram negative bacteria. Oxytetracycline was taken as a standard and for negative control DMSO was used. All plant extracts were examined at four different concentrations (250, 100, 33.33, 11.11  $\mu\text{g}/\text{mL}$ ) and then MIC was recorded. Results revealed that the selected plant extracts showed resistance against three bacterial strains (Table 2). CIM showed maximum zone of inhibition against *K. pneumonia* (ATCC 1705) ( $20 \pm 1.5$  mm) followed by *S. aureus* (ATCC 6538) ( $19 \pm 1.24$  mm) and *E. coli* (ATCC 33456) ( $14 \pm 1.16$  mm). In case of CIH, maximum antibacterial activity was observed against *K. pneumonia* ( $12 \pm 1.94$  mm) followed by *E. coli* ( $6 \pm 1.04$  mm). However, CIH did not showed any zone of inhibition against *S. aureus*. Hence, CIM plant extract revealed significant antibacterial activity as compared to n-hexane extracts when tested against three bacterial strains. Results are expressed as mean  $\pm$  SD (n=3).

#### 3.4. Antifungal activity.

Antifungal assay of methanol and n-hexane extracts of *C. iberica* was done using agar disc diffusion method. Selected plant extracts were examined using four different concentrations (250, 100, 33.33, 11.11  $\mu\text{g}/\text{mL}$ ). Total three fungal strains (*C. albicans*, *M. racemosus*, and *A. niger*) were used to examined antifungal activity of these extracts (Table 3). Chloramphenicol was taken as a standard and for negative control DMSO was used. Maximum zone of inhibition was produced by CIM extract of *C. iberica* against *A. niger* (FCBP 0918) ( $18 \pm 2.4$  mm) followed by *M. racemosus* (FCBP 0300) ( $16 \pm 2.5$  mm) and *C. albicans* (FCBP 478) ( $15 \pm 2.6$  mm). However, minimum zone of inhibition was determined by CIH against *C. albicans* ( $14 \pm 2.6$  mm) and *A. niger* ( $12 \pm 1.5$  mm). No zone of inhibition was formed by CIH against *M. racemosus*.

**Table 2:** Anti-bacterial activity of *C. Iberica* (CIM and CIH).

Microorganisms		<i>E. coli</i> ATCC33456		<i>K. pneumonia</i> ATCC 1705		<i>S. aureus</i> ATCC6538	
Plant Extracts		<sup>15</sup> ZOI (mm)	MIC ( $\mu\text{g/ml}$ )	ZOI (mm)	MIC ( $\mu\text{g/ml}$ )	ZOI (mm)	MIC ( $\mu\text{g/ml}$ )
CIM		14 $\pm$ 1.16	100	20 $\pm$ 1.5	33.33	19 $\pm$ 1.24	33.33
CIH		6 $\pm$ 1.04		12 $\pm$ 1.94	100	NI	
DMS (Negative Control)		NI		NI		NI	
Oxytetracycline (Standard)		30 $\pm$ 1.5		28 $\pm$ 3.5		32 $\pm$ 2.8	

**Key:** CIM: *C. iberica* methanolic extract; CIH: *C. iberica* n-hexane extract;

**Table 3:** Antifungal activity of *C. iberica* (CIM and CIH).

Fungal strains	<sup>22</sup> <i>M. racemosus</i>	FCBP 0300	<i>C. albicans</i>	FCBP 478	<i>A. niger</i>	FCBP 0918
Plant Extracts	<sup>7</sup> ZOI (mm)	MIC ( $\mu\text{g/ml}$ )	ZOI (mm)	MIC ( $\mu\text{g/ml}$ )	ZOI (mm)	MIC ( $\mu\text{g/ml}$ )
CIM	16 $\pm$ 2.5	33.33	15 $\pm$ 2.6	100	18 $\pm$ 2.4	100
CIH	NI	-	14 $\pm$ 2.6	100	12 $\pm$ 1.5	33.33
DMSO (Negative Control)	NI	-	NI	-	NI	-
Chloramphenicol (Standard)	28 $\pm$ 2.5	-	32 $\pm$ 3.5	-	31 $\pm$ 1.8	-

**Key:** CIM: *Centaurea iberica* methanolic extract; CIH: *Centaurea iberica* n-hexane extract;

### 3.5. Anticancer assay.

Anticancer effect of the methanolic extract of *C. iberica* was tested against 3T3 cell line and PC3 cell line using MTT assay. For this, MTT was taken as a standard against 3T3 cell lines and PC3 cell lines. The results revealed significant anticancer activity of the extract towards PC3 cell lines (5.0 % inhibition) compared to the 3T3 cell lines (30.4 % inhibition) respectively. IC<sub>50</sub> values also confirmed highest activity against PC3 cell lines (89.9 %) while it was found to be lowest in case of 3T3 cell lines i.e < than 50 % (Table 4).

**Table 4:** Anticancer activity 3T3 cell line and PC3 cell line of *Centaurea iberica*.

Anticancer assay	3T3 cell line		PC3 cell line	
	% inhibition	IC <sub>50</sub>	% inhibition	IC <sub>50</sub>
Plant extract	30.4 %	< 50	5.0%	89.9 %
Standard	89.9 %	0.8 ± 0.14	Inactive	1.9 ± 0.4

**Key** CIM: *C. iberica* methanolic extract; **standard:** MTT

against 3T3 cell line and PC3 cell line.

#### 4. DISCUSSION

<sup>14</sup> Medicinal plants play a very promising role in the treatment of a variety of diseases (Naseer et al., 2022, Ijaz et al., 2023). The nature has blessed medicinal plants with a wide range of bioactive phytochemicals (Abbasi et al., 2022, Zahra et al., 2023). <sup>24</sup> The present study was designed to explore the biological potential of *C. iberica*. <sup>9</sup> The current research, study was designed to explore the phytochemical composition and biological potentials of *C. iberica*. Besides beneficial microorganisms, various pathogenic organisms are also widespread in nature that lead to serious health issues. Hence, antimicrobial activity of two extracts of *C. iberica* (CIM and CIH) were tested <sup>17</sup> against one-gram positive bacteria (*S. aureus*), two gram negative bacteria (*E. coli* and *K. pneumonia*) and three <sup>35</sup> fungal strains (*C. albicans*, *M. racemosus* and *A. niger*) using disc diffusion assay. Selected plant extracts were examined at four different concentrations (250, 100, 33.33, 11.11 μg/mL) and oxytetracycline and chloramphenicol were used as positive control (standard). Antibacterial results revealed that the methanolic plant extract showed maximum zone of inhibition against *K. pneumonia* (ATCC

1705)  $20 \pm 1.5$  mm, *S. aureus* (ATCC 6538)  $19 \pm 1.24$  mm and *E. coli* (ATCC 33456)  $14 \pm 1.16$  mm. However, n-hexane extracts formed less than 14 mm zone of inhibition against all bacterial strains. Similarly, in case of antifungal activity, maximum zones were revealed by CIM against all fungal strains as compared to n-hexane extracts which revealed less than 14 mm zones of inhibition. Altogether, the methanol extracts of *C. iberica* were found to be very effective against wide range bacterial and fungal strains than n-hexane extracts. Previously, Albayrak et al. (2017) reported significant antibacterial activity ( $7.00 \pm 0.00$  mm) of *C. aksoyi* against *E. coli* and 7.00 to 11.00 mm inhibition zones were detected by *C. amaena* against various gram positive and gram-negative bacterial strains. Similarly, according to Barbour et al. (2004) *C. erengoides* and *C. aintensis* exhibits 66.66 % and 88.8% antibacterial activity respectively. From the obtained results, it can be suggested that antimicrobial activity in the examined species might be due to the presence of bioactive compounds present in *Centaurea* species and hence can be utilized in various industries (Barbour et al., 2004). To the best of our knowledge and literature surveys, this is the first study reporting the antimicrobial activity of *C. iberica* extracts.

Medicinal plants are considered as the rich source of antioxidants and reduces the oxidative stress to great extent due to the presence of different phenolic compounds. These species can be used to cure various human ailments such as diabetes, cancer and heart ailments (Ali et al., 2022, Gul et al., 2022). In present studies, antioxidant activity of *C. iberica* extracts was examined via two assays viz. DPPH and TRP assays. Results revealed that CIM extracts exhibited maximum % inhibition (97.14 %) at 1000  $\mu\text{g/mL}$  and  $\text{IC}_{50}$  value of 1.54  $\mu\text{g/mL}$  followed by CIH extract which displayed 94.65 % inhibition and  $\text{IC}_{50}$  value of 36.61  $\mu\text{g/mL}$  at 1000  $\mu\text{g/mL}$ . Highest TRP ( $20.45 \pm 1.6$  GAE mg/g) was also noted in CIM extracts as compared to other extracts of *C. iberica*. Current research confirmed that the and TRP values are greatly influenced by the use of different solvents (Kumar and Jain, 2015; Mehwish et al.,

2019). Previously, Zengin et al. (2011) reported 44.0 % DPPH inhibition (137.06 IC<sub>50</sub>) and 39.70 mg AAE/g TAC value in the methanol extract of *C. urvillei*. Similarly, Escher et al. (2018) observed that *C. cyanus* possess 51-62 % DPPH (2,2-diphenyl, 1-picrylhydrazine) inhibition, 487 to 846 mg GAE/100g reducing capacity and 1134 to 1359 mg AAE/100 g TAC activity.

31 Cancer is one of the major causes of death in the worldwide and has been reported for centuries that plants have anticancer properties and they are important source of anticancer agents (Cragg and Newman, 2013). The *Centaurea* genus contains anticancer properties due to the availability of phytochemicals (Alper and Gunes, 2019). Hence, present 34 study was carried out to observe the anticancer activity of CIM extracts against 3T3 and PC3 cell lines. Results revealed that CIM extract exhibited 30.4 % inhibition against 3T3 cell lines and 5.0 % inhibition against PC3 cell lines. Previously similar experiment was conducted on various species of *Centaurea* including *C. solstitialis*, *C. kilaea*, *C. cuneifolia*, *C. salicifolia* and *C. 25 stenolepis* (Sekerler et al., 2018). According to their results *Centaurea* genus have good potential as an anti-cancer agent. The plant demand is increasing day by day due to their less side effects on normal. Numerous species were studied from Asia and Africa, the herbal therapies against cancer are common in undeveloped countries (Azmi et al., 2006). Chemotherapy is not as safe as plant based medicines, therefore the mandate for substitute care with naturally derived anticancer active agents with plants being the favorite cause (Uddin et al., 2021). According to the results CIM is not a potential candidate for insecticidal. However, on the basis of overall results it can be suggested that *C. iberica* is highly enriched with secondary metabolites and thus is effective against ailments including cancer and inflammation (Koca et al., 2009 and Ochwang et al. (2009).

## 5. CONCLUSION

In current study, antioxidant, antifungal, anti-bacterial, anticancer and antioxidant activities were carried out using different extracts of *C. iberica*. Among all the plant extracts, antioxidant activity was found to be maximum in the methanolic extract. Further, methanolic extract has shown promising antimicrobial activities as compared to other extracts. Therefore, it can be assumed from the biological activities of selected plant exhibits significant phyto-compounds that can be utilized effectively against various pathogens for defensive purposes. In future, different other *in vitro* and *in vivo* studies should be carried out to further confirm the bio-efficacy of selected plant extract and compound characterization will help in isolating various novel compounds that could be used to treat different types of ailments.

**Acknowledgments:**

16

This work was funded by Researchers Supporting Project number (RSPD2023R728), King Saud University, Riyadh, Saudi Arabia.

15%

SIMILARITY INDEX

13%

INTERNET SOURCES

11%

PUBLICATIONS

5%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** Ishan Saini, Prashant Kaushik, Asma A. Al-Huqail, Faheema Khan, Manzer H. Siddiqui. "Effect of the diverse combinations of useful microbes and chemical fertilizers on important traits of potato", Saudi Journal of Biological Sciences, 2021  
Publication 1%
- 2** Sana Naseer, Javed Iqbal, Abeel Naseer, Sobia Kanwal et al. "Deciphering Chemical Profiling, Pharmacological Responses and Potential Bioactive Constituents of Saussurea lappa Decne. extracts through In Vitro Approaches", Saudi Journal of Biological Sciences, 2022  
Publication 1%
- 3** [pubmed.ncbi.nlm.nih.gov](https://pubmed.ncbi.nlm.nih.gov)  
Internet Source 1%
- 4** [mdpi-res.com](https://mdpi-res.com)  
Internet Source 1%
- 5** Zakir Ullah, Javed Iqbal, Banzeer Ahsan Abbasi, Wasim Akhtar et al. "Assessment of Gus Expression Induced by Anti-Sense OsPPO 1%

# Gene Promoter and Antioxidant Enzymatic Assays in Response to Drought and Heavy Metal Stress in Transgenic *Arabidopsis thaliana*", Sustainability, 2023

Publication

6

Submitted to Higher Education Commission  
Pakistan

Student Paper

1 %

7

[pr.hec.gov.pk](http://pr.hec.gov.pk)

Internet Source

1 %

8

[www.mdpi.com](http://www.mdpi.com)

Internet Source

1 %

9

Syeda Anber Zahra, Javed Iqbal, Banzeer Ahsan Abbasi, Tabassum Yaseen et al. " Scanning electron microscopy of L. seeds and their cytotoxic, antimicrobial, antioxidant, and enzyme inhibition potentials ", Microscopy Research and Technique, 2021

Publication

1 %

10

Syeda Anber Zahra, Javed Iqbal, Banzeer Ahsan Abbasi, Amir Shahbaz, Sobia Kanwal, Syed Lal Shah, Parvaiz Ahmad, Tariq Mahmood. " Antimicrobial, cytotoxic, antioxidants, enzyme inhibition activities, and scanning electron microscopy of (Boiss.) Boiss. seeds ", Microscopy Research and Technique, 2021

Publication

1 %

11	<a href="https://link.springer.com">link.springer.com</a> Internet Source	1 %
12	Ayesha Hameed, Muhammad Zafar, Mushtaq Ahmad, Shazia Sultana et al. " Chemotaxonomic and biological potential of highly therapeutic plant Retz. using multiple microscopic techniques ", Microscopy Research and Technique, 2021 Publication	<1 %
13	<a href="http://www.rombio.eu">www.rombio.eu</a> Internet Source	<1 %
14	<a href="http://www.science.gov">www.science.gov</a> Internet Source	<1 %
15	<a href="http://www.hindawi.com">www.hindawi.com</a> Internet Source	<1 %
16	Submitted to Sophia University Student Paper	<1 %
17	<a href="http://ijmpronline.com">ijmpronline.com</a> Internet Source	<1 %
18	<a href="http://www.journalijar.com">www.journalijar.com</a> Internet Source	<1 %
19	Javed Iqbal, Banzeer Ahsan Abbasi, Tabassum Yaseen, Syeda Anber Zahra et al. "Green synthesis of zinc oxide nanoparticles using <i>Elaeagnus angustifolia</i> L. leaf extracts and	<1 %

their multiple in vitro biological applications",  
Scientific Reports, 2021

Publication

20

[www.informaticsjournals.com](http://www.informaticsjournals.com)

Internet Source

<1 %

21

[starc.stthomas.ac.in:8080](http://starc.stthomas.ac.in:8080)

Internet Source

<1 %

22

[www.thejaps.org.pk](http://www.thejaps.org.pk)

Internet Source

<1 %

23

[baadalsg.inflibnet.ac.in](http://baadalsg.inflibnet.ac.in)

Internet Source

<1 %

24

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1 %

25

Hanifa Fatullayev, Leyla Paşayeva, Ismail Celik, Ufuk İnce, Osman Tugay. "

Phytochemical Composition, Antimicrobial, Antioxidant, and Enzyme Inhibition Activities, and Molecular Docking and Dynamics

Simulations of : A Computational and Experimental Approach ", ACS Omega, 2023

Publication

<1 %

26

Syed Shah, Yaseen Rehman, Arshad Iqbal, Zia Rahman, Bangwei Zhou, Mu Peng, Zhijian Li.

"Phytochemical Screening and Antimicrobial Activities of Stem, Leaves and Fruit Extracts of

<1 %

# Viscum album L.", Journal of Pure and Applied Microbiology, 2017

Publication

27

[cmjournal.biomedcentral.com](http://cmjournal.biomedcentral.com)

Internet Source

<1 %

28

[core.ac.uk](http://core.ac.uk)

Internet Source

<1 %

29

[dx.doi.org](http://dx.doi.org)

Internet Source

<1 %

30

[ijarbs.com](http://ijarbs.com)

Internet Source

<1 %

31

[www.prolekare.cz](http://www.prolekare.cz)

Internet Source

<1 %

32

Ajila, C.M.. "Bioactive compounds and antioxidant potential of mango peel extract",  
Food Chemistry, 2007

Publication

<1 %

33

[journaljpri.com](http://journaljpri.com)

Internet Source

<1 %

34

[psasir.upm.edu.my](http://psasir.upm.edu.my)

Internet Source

<1 %

35

[www.jbpr.in](http://www.jbpr.in)

Internet Source

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off