

Asad Final 23-05-24

by Saadullah Malik

Submission date: 22-May-2024 09:47PM (UTC-0700)

Submission ID: 2386155552

File name: asad_final_23-05-24.docx (4.86M)

Word count: 4347

Character count: 24852

Appraisal of the anti-arthritic potential of *Strelitzia reginae* in Wistar rats via modulating molecular inflammatory cascades IL-6, IL-17a, IL-1 β , TNF- α , IFN- γ , RANKL, OPG, and IL-4 and attenuating oxidative stress biomarkers

Abstract

Methanolic aerial extract of the plant *Strelitzia reginae* (MSRA) was the subject of the present investigation because of its potential anti-arthritic effects. The research utilised Wistar albino rats, and all groups except the control group had arthritis in their right hind paws caused by 1 ml of CFA (Complete Freund's adjuvant) emulsion. For 21 days, all rats (except those in the normal and arthritic groups) were administered the extract at doses of 600, 300, and 150 mg/kg/day in conjunction with 10 mg/kg of piroxicam. By enhancing biochemical markers, reducing paw oedema, and reestablishing immunological organs and body weight, the extract shown substantial effects. Polymerase chain reaction (PCR) results demonstrated a significant decrease in receptor activator of nuclear factor kappa-beta ligand (RANKL), interleukin-17a (IL-17a), tumour necrosis factor alpha (TNF- α), interleukin-1 β (IL-1 β), and interferon- γ (IFN- γ). The study also found that when polyarthritic groups were treated with dosages of MSRA, the expression of osteoprotegerin (OPG) and IL-4 was upregulated. Around 600 mg/kg/day, the highest levels of OPG (1.24 \pm 0.025 ng/ml) and IL-4 (80.31 \pm 0.23-fold) were seen. The levels of catalase (CAT) and superoxide dismutase (SOD) increased ($p < 0.05$) in the groups treated with 600mg/kg MSRA and piroxicam, while malondialdehyde (MDA) decreased. Arthritic rats showed less inflammation and pannus development in their ankle joints when examined histologically.

Key words: rheumatoid arthritis, *Strelitzia reginae*, oxidative stress biomarkers

Introduction

In its most basic form, rheumatoid arthritis is a kind of symmetric polyarthritis that affects both the big and tiny diarthroid joints in the body. The synovium becomes invaginated by CD4+ T lymphocytes, B cells, and macrophages in rheumatoid arthritis, causing distinct lymphoid aggregates to form with a germinal core (Firestein 2003). Approximately 0.5% to 1% of the global population is impacted by rheumatoid arthritis. There are three distinct stages to the course of a disease. Fluid retention and synovial hypertrophy occur during the initial, asymptomatic phase of the process, which is characterised by the activation of inflammatory

cells and cytokines. The activation of T cells in the second phase (the amplification phase) leads to the infiltration and destruction of subchondral bone and articular cartilage; this process is gradual, permanent, and symptomatic. In the end, the last stage leads to ankylosis of the joints and long-term abnormalities (Scionti et al., 2016). In rheumatoid arthritis, the metabolic enzymes lipo-oxygenase and cyclo-oxygenases (COX) and the proinflammatory cytokines interleukin-1 β , IL-6, and tumour necrosis factor-alpha (TNF- α) find a balance. The breakdown of proteoglycans and collagen is aided by pro-inflammatory cytokines, which hinder their formation. In addition, prostaglandins—most notably prostaglandin E2—are produced by cyclo-oxygenase COX 2, which in turn causes inflammation, pain, and impairment (El-Miedany and El Rasheed 2002). Therefore, rofecoxib and celecoxib are selective COX-2 inhibitors used to treat rheumatoid arthritis (RA) because they lessen inflammation and discomfort without causing the gastrointestinal side effects sometimes seen with nonsteroidal anti-inflammatory drugs (NSAIDs) (Farzaei et al., 2016, Saleem et al., 2020). An herbaceous perennial with evergreen foliage, *Strelitzia reginae* is a monocotyledon plant. Subtropical temperate climates are ideal for its growth (Yassen and Badawy ◌ Abou-Sreea 2016).

When it came to chemical bioprospecting of *Strelitzia reginae*, there was basically very little information available in the scientific literature. Therefore, this study set out to assess the anti-arthritic potential of *Strelitzia reginae*'s methanolic aerial extract. In this investigation, the dosages of 600, 300, and 150 mg/kg/day were chosen by considering the rats' body weight. Furthermore, by employing the CFA-induced arthritic model, the molecular and cellular mechanisms that alleviate arthritic and inflammatory activity were ascertained. By reducing oxidative stress indicators, regulating inflammatory cytokines, and improving histological implications, the study demonstrated substantial anti-arthritic potential.

Materials and methods

Collection of plant and preparation of extraction

The *Strelitzia reginae* plant was bought from a nursery in Patoki, Pakistan. *Strelitzia reginae* was recognised by renowned taxonomist Mansoor Hammed of the University of Agriculture Faisalabad and given the voucher number 1-173-2022. It was sent to the herbarium after that. To remove any dirt or insects, the gathered plant was washed with distilled water. The next step was to let the plant dry in the shade for ten days. There were 20 kilogrammes of dried plant material. The aerial and root portions of the plant were separated. The parts that were used for the extraction process were ground into a coarse powder separately, and then the triple maceration was applied. Using 50 cc of 70% methanol using the

same method as before, the residue was again extracted. A volumetric flask was used to collect the supernatant, which was then evaporated using a rotary evaporator (Stuart®, UK) at 40°C under reduced pressure (Oroian et al., 2020). The methanolic aerial parts of extract were named MSRA (methanolic extract of aerial parts of *Strelitzia reginae*).

Experimental animals

The study was carried out by choosing previously acclimatized healthy Wistar rats with a weight between 150 and 220g. There were six different groups made with a random allocation of rats. The rats were confined in the animal house of the faculty of pharmaceutical sciences at GC University Faisalabad.

CFA-induced arthritis (Complete Freund's adjuvant-induced)

We divided the rats into six groups of two each. Separate from Group II, which was dubbed AC (arthritic control), Group I was called NC (normal control). Likewise, Group (III) was given piroxicam (10 mg/kg/day) as the usual control group, while Groups (IV), (V), and (VI) were given MSRA doses of 150, 300, and 600 mg/kg/day, respectively. On the initial day, the right subplantar area of the hind paw was targeted to induce arthritis in all mice that were not administered NCG. The procedure to induce arthritis was injecting 1 millilitre of CFA emulsion intradermally. (Sigma-Aldrich, Washington, DC, USA). Over the course of seven days, the animals did not receive any therapeutic dosages. Oral gavage was then initiated and maintained until day 28 of therapy.

Phytochemical assessment by High performance liquid chromatography (HPLC)

Using HPLC in accordance with the protocol laid forth in a research by Saleem et al., 2019 allowed us to identify the different chemicals included in MSRA.

Evaluation of polyarthritis

Throughout the course of the research, we noticed a few of the following physical characteristics: At 1, 7, 12, 16, 20, and 28 days, an electronic vernier calliper was used to assess paw diameters, body weight, and arthritis score in order to diagnose polyarthritic edoema (Zhang et al., 2017).

Hematological and biochemical investigation

After 29 days of starvation and a light ether anaesthesia, the animals were prepared for sacrifice. For haematological and biochemical tests, blood samples were taken, correspondingly. In order to measure C-reactive protein (CRP), a commercially available kit from Antec Diagnostic Products®, UK, was also utilised. The levels of creatinine, urea, alkaline phosphatase (ALP), alanine transaminase (ALT), rheumatoid factor (RF), and aspartate aminotransferase (AST) were also measured using an automated chemistry analyzer (Microlab 300®) and protocols developed by Analyticon Biotechnologies AG of Germany. Similarly, a complete blood count (CBC) was also determined using an automated hemocytometer (Sysmex®, Roche, Germany).

Immune organ weight

The sacrificed rats were cut open through the abdomen. The thymus and spleen were taken out, cleaned with distilled water, and weighed.

In vivo estimation of oxidative stress biomarkers

Oxidative stress biomarkers such as malondialdehyde (MDA), catalase (CAT), superoxide dismutase (SOD) levels were calculated from CFA-induced arthritis-developed rats (Jurczuk et al., 2004).

qRT-PCR

The following inflammatory biomarkers were evaluated in cases of arthritis: IL-17a, TNF- α , IL-4, IL-6, interferon- γ , OPG, and RANKL. Blood was drawn using a heart puncture and deposited in EDTA tubes for analysis. Geno All® of Korea supplied the RiboEx™LS kit that the TRizol technique utilised to extract RNA. In addition, the amount of RNA was measured using a Nano drop spectrophotometer. Additionally, the WizScript® cDNA synthesis kit (maker's protocol) was used to synthesise complementary DNA (Cui et al., 2022).

Statistical analysis

The results were presented as the average plus or minus the standard deviation after a statistical analysis. The data was analysed using two-way analyses of variances (ANOVA). This included percentage inhibition, paw diameter, body weight fluctuation in arthritis-induced rats, and all other relevant data. A similar analysis was conducted using GraphPad Prism® 6.01 to assess the

impact of plant extracts and the gold standard medicine, piroxicam, on the arthritic score of rats throughout many study days.

Results

High performance liquid chromatography (HPLC) Evaluation

Figure 1 shows that the aerial sections of the methanolic extract included the most essential components identified by HPLC evaluation: p-coumaric acid, kempherol, and gallic acid.




Figure 1 The HPLC spectrum shows phytochemicals detected in methanolic roots extract of *Strelitzia reginae*

Paw swelling assessment

After administering CFA subplantarly on the eighth day of the research, the most significant edoema was noted. The arthritic control (AC) group showed an increase in paw diameter from day eight to day twenty-one. Paw diameter was shown to be reduced by treatment with a methanolic extract of MSRA; the most significant impact was observed at 600 mg/kg when compared to AC, as indicated. At 600 mg/kg, the MSRA extract had the greatest impact compared to the arthritic control group, according to the x-ray examination of the hind paws (figure 2).




Fig.2 x-ray evaluation showed significant improvement in CFA-induced arthritic hind paws by methanolic aerial extract of *Strelitzia reginae* especially at 600 mg/kg dose. Here, (a) shows (NC), (b) represents (AC), (c) exhibited piroxicam-treated, (d) 150 mg/kg (e) 300 mg/kg and (f) 600 mg/kg represents effects of different doses

Effect on body weight, paw swelling and arthritic score

Throughout the course of the research, participants' weight, paw edoema, and arthritis score were measured at various intervals. Body weight, paw edoema, and arthritic score were all positively affected by the 16–28 day treatment of piroxicam and MSRA dosages. Figure 3b shows that the piroxicam-treated group had changes in body weight, paw edoema, and arthritic score at the 300 mg and 150 mg dosing levels, respectively.




Fig.3a-c The effect on body weight in CFA-induced arthritic rats by treatment with methanolic extract of aerial parts of *Strelitzia reginae* (MSRA). Here, a shows effect on paw diameter, b exhibited effect on arthritic score and c represents effect on body weight.

Effect on hematological parameters, kidney and liver function parameters

Rats given FCA had elevated levels of rheumatoid factor (RF), alkaline phosphatase (ALP), C-reactive protein (CRP), and alanine transferase (ALT). In rats treated with conventional medication and plant extracts, a significant decrease in ALP, ALT, CRP, and RF was seen ($p < 0.0001$). The most effective dose of MSRA in alleviating arthritis in rats was 600 mg/kg, which restored HB, platelets, RBCs, and WBCs significantly ($p < 0.05$) when compared to AC without HB, platelets, RBCs, and WBCs. MSRA extract treatment resulted in an increase of HB and RBCs in arthritic rats. Figure 4 shows that white blood cell and platelet counts dropped compared to the normal control group (NC). Control groups without arthritis exhibited significantly higher levels of the systemic biomarkers C-reactive protein and RF. Additionally, as seen in Figure 5, there were no statistically significant changes in creatinine and urea levels between the groups treated with piroxicam and plant extract.

The arthritic control group also had significantly higher levels of ALP and ALT. Figure 6 shows that compared to the piroxicam-treated groups, the ALP and ALT levels reduced at 600mg dosages.




Fig.4 Effect of aerial methanolic extract of *Strelitzia reginae* (MSRA) on hematology of FC-induced arthritic rats. One-way ANOVA followed by Tukey's test was performed and values were expressed as mean \pm S.D (n = 6). Where ***, +++, ... exhibited significant difference (p < 0.05) in respect to (NC) normal control, (AC) arthritic control and piroxicam-treated group respectively. Here, FCA= Freund's complete adjuvant, MSRA=Methanolic extract of aerial parts of *Strelitzia reginae*.




Fig.5 In rats with hind paw-induced arthritis, the effect of an aerial methanolic extract of *Strelitzia reginae* (MSRA) on C-reactive protein (CRP), rheumatoid factor (RF), and parameters measuring renal function was investigated. A one-way analysis of variance (ANOVA) and Tukey's test were used for statistical analysis. The mean \pm S.D. values were calculated from a sample size of 6. Except for the normal control, arthritic control, and Piroxicam, there was a significant difference ($p < 0.05$) between ***, +++, and... Specifically, "NC" stands for "normal control," "AC" for "arthritic control," "FCA" for Freund's complete adjuvant, and "MSRA" for a methanolic extract of *Strelitzia reginae* aerial parts.




Fig.6 In a study of rats with hind paw-induced arthritis, the researchers looked at how the aerial methanolic extract of *Strelitzia reginae* (MSRA) affected liver function tests. The results were evaluated using one-way ANOVA followed by Tukey's test, and the average plus or minus the standard deviation (n = 6) were used as the values. In comparison to the normal control, the arthritic control, and piroxicam, ***, +++, and... showed a significant difference ($p < 0.05$) accordingly. Specifically, "NC" stands for "normal control," "AC" for "arthritic control," "FCA" for Freund's complete adjuvant, and "MSRA" for a methanolic extract of *Strelitzia reginae* aerial parts.

Effect on immune organ weight

The weights of thymus and spleen were significantly higher in arthritic control group ($p < 0.05$) as compare to normal control. It was noted the significant weight restoration of the thymus and spleen by treating the arthritic rats with plant extracts and maximum action was exhibited at the dose of 600 mg/kg.




Figure 7 Immune system mass in arthritic rats as a function of *Strelitzia reginae* methanolic extract. We used one-way ANOVA and Tukey's test to evaluate the data, and we calculated the values as the mean \pm S.D. (n = 6). There was a significant difference ($p < 0.05$) between the normal control, the arthritic control, and the group treated with piroxicam when the letters ***, +++, and... were used. *Strelitzia reginae* aerial parts methanolic extract (MSRA)

Effect on oxidative stress markers

The AC group showed a significant decrease in SOD and CAT activity ($p < 0.0001$) when compared to the normal control group. However, as seen in figure 8, the SOD and CAT activities were considerably enhanced by the combination of 600 mg of MSRA and piroxicam. When comparing the AC group to the usual control group, a significant increase in MDA levels ($p < 0.0001$) was also noted. Fig.8 shows that compared to the arthritic control group, MSRA at 600 mg plus piroxicam significantly reduced MDA levels.




Figure 8 a-c Reaction between oxidative stress indicators in arthritic rats and a methanolic extract of *Strelitzia reginae* aerial parts. Biomarkers for oxidative stress were markedly enhanced by the MSRA extract. In contrast to the groups treated with NC, AC, and piroxicam, respectively, ***, +++, and... showed a significant difference ($p < 0.05$). The data is examined using one-way ANOVA, then Tukey's test, and the results are shown as the mean \pm S.D. ($n = 6$). Specifically, "FCA" stands for Freund's complete adjuvant, "MSRA" for methanolic extract of aerial portions of *Strelitzia reginae*, "NC" for normal control, and "AC" for arthritic control.

Effect on gene expression

During the course of the investigation, genetic markers for inflammation were expressed in Wistar rats. Arthritic control group levels significantly increased compared to normal control group levels of IL-6, TNF-Alpha, IL-17a, IFN- γ , IL-1 β , and RANKL expression ($p < 0.0001$). Nonetheless, at doses of 600 and 300 mg/kg, MSRA extract reduced this increase in arthritic rats. Figure 11a shows that rats treated with 600 mg/kg of extract outperformed those treated with piroxicam.

The arthritic control group showed a considerably lower level of mRNA expression of osteoprotegerin (OPG) and IL-4 compared to the normal control group ($p < 0.0001$). Nevertheless, as shown in figure 12a, the arthritic control group showed no increase in OPG levels when treated with piroxicam at a dosage of 600 mg/kg.




Fig.9a-f

Modulation of inflammatory cytokine expression in CFA-induced polyarthritic rats by a methanolic extract of *Strelitzia reginae* aerial parts. At 600 mg/kg, MSRA had the greatest

impact in reducing the expression of IL-6, TNF-ALPHA, IL-17a, IFN- γ , IL-1 β , and RANKL mRNA. In comparison to the normal control, the arthritic control, and the piroxicam-treated groups, ***, +++, and... showed significant differences, respectively. The data are analysed using one-way ANOVA followed by Tukey's test, and the results are shown as the mean \pm S.D. (n = 6). Specifically, "NC" stands for "normal control," "AC" for "arthritic control," and "MSRA" for "methanolic extract of aerial parts of *Strelitzia reginae*."




Fig. 10: Anti-inflammatory cytokine expression in CFA-induced polyarthritic rats and the effect of methanolic aerial extract of *Strelitzia reginae* (MSRA). At 600 mg/kg, MSRA therapy leads to the greatest upregulation of OPG and IL-4 expression levels. In relation to the NC, AC, and piroxicam-treated groups, ***, +++, and... showed a significant difference ($p < 0.05$) accordingly. The data are shown as the mean \pm S.D. (n = 6), and they were analysed using one-way ANOVA followed by Tukey's test. We have "FCA" (Friend's complete adjuvant), "MSRA" (Methanolic extract of aerial portions of *Strelitzia reginae*), "NC" (normal control), and "AC" (arthritic control).

Discussion

The study's overarching goal was to identify the probable cause of polyarthritis in Wistar rats and

assess the anti-arthritic capabilities of the *Strelitzia reginae* plant. The investigation was conducted using a CFA-induced arthritis model. The use of HPLC allowed for the qualitative evaluation of the phytochemical ingredients of MSRA. The measurements of the paws and the differences in body weight were also documented. Additionally, the effects of MSRA were revealed by recording the biochemical and haematological markers. Hydraulic acid, gallic acid, and kempherol were among the phytochemicals identified by high-performance liquid chromatography (HPLC) as having medicinal, antioxidant, and anti-inflammatory properties. The anti-inflammatory properties of gallic acid have also been deduced from many research (Kroes et al., 1992). Among the flavonoids present in fruits and vegetables is the well-known antioxidant kaempferol. The source is Silva dos Santos et al. (2021). The phenolic hydroxycinnamic acid (P-coumaric acid) is present in many foods and drinks, including fruits, vegetables, and beverages. According to Abramovič (2015), many studies have found that HCs have antioxidant properties and shield crucial components from oxidation. One of the most used methods for assessing the possibility of anti-arthritic effects is the CFA-induced arthritic model. Animal studies have shown that it can imitate rheumatoid arthritis, and its similarities to the processes of human arthritis have led to its widespread acceptance. The stated source is Liu et al. (2022). There are two potential pathways that might culminate in the onset of arthritis following CFA treatment. The acute stage, which lasts for the first 10 days, is characterised by the development of primary lesions. During the subsequent 11–28 days, known as the chronic stage, synovitis and lesions on the opposite side of the joint will appear (Saleem et al., 2020). The results from the polyarthritic rats show that thrombocytopenia and total leukocyte count (TLC) are higher in arthritic patients. The correlation between thrombocytosis and rheumatoid arthritis is substantial. Hutchinson et al. (1976) found that active intravascular coagulation causes an increase in platelet formation in this condition. The results of this investigation demonstrate that piroxicam and plant extracts can restore TLC and thrombocytes. Thirty to sixty percent of arthritis patients reportedly experience anaemia symptoms (Perumal et al., 2017). In rheumatoid arthritis, anaemia manifests as a low red blood cell count. Shabbir et al. (2018) found that polyarthritic is associated with elevated IL-6 expression, another factor that contributes to anaemia. The results showed that the groups treated with the plant extract were able to restore red blood cell count and anaemia.

One marker of disease activity in RA patients is the blood level of C-reactive protein (Deyo et al., 1980). According to Scott (2000), RF positive is just as significant as other markers for the development of synovitis, the arthritic condition that causes joint destruction and functional impairment. In the arthritis control group, CRP and RF levels were found to be higher. The levels of ALP and ALT are increased by bone erosion and inflammatory chemicals in the body, but they are decreased in the groups treated with plant extract and piroxicam. The levels were significantly decreased after being treated with piroxicam and plant extracts (Comar et al., 2013).

Arthritic rats have heavier immune organs, as shown in earlier studies (Zhang and An 2007). Similarly, the present study found that administering plant extract to arthritic rats resulted in a return to normalcy of immune organ weights and systemic indicators. Along with antioxidant enzymes superoxide dismutase (SOD) and catalase (CAT), malondialdehyde (MDA) is now acknowledged to be a significant oxidative stress indicator. Lipid per oxidation produces malondialdehyde (MDA), a clinically characterised end product of oxidative stress. By transforming radicals of superoxide anion into oxygen and hydrogen peroxide, superoxide dismutase (SOD) is an essential enzyme. As a result, it plays a crucial role in regulating ROS levels within cells. In a similar vein, CAT helps cells avoid oxidative stress by breaking down hydrogen peroxide into oxygen and water. The stress indicators in the present investigation showed an increase in rats that were induced arthritis, but they were effectively recovered with exercise treatment with a combination of plant extract and piroxicam (Camkurt et al., 2017).

A typical symptom of rheumatoid arthritis is cachexia. The overproduction of TNF- α and interleukin-1 β is likely the primary cause of this. Packed together, tumour necrosis factor- α and interleukin-1 β produce cachexia. Changes in the ratio of protein synthesis to breakdown in muscles are brought about by these causes. According to Walsmith and Roubenoff (2002), when there is an imbalance like this, muscle atrophy occurs. A research found that inflammation driven biochemical processes are increased when activated macrophages spread pro-inflammatory cytokines. The research indicates that cytokines such as TNF- α , IL-1, and IL-6 are vital in the development of pain. The present investigation found that administering a plant extract to arthritic rats reduced levels of pro-inflammatory cytokines (Zhang and An 2007).

Evidence suggests that RA patients have elevated COX-2 expression in their synovial tissues. This increased expression is caused by pro-inflammatory cytokines including IL-1 and TNF- α . Using a plant extract to treat arthritic rats significantly restores pro-inflammatory factors (Crofford 1999). A cytokine with pleiotropic effects is tumour necrosis factor alpha (TNF- α). It is a homotrimer protein of 157 amino acids that is produced by T-lymphocytes, activated macrophages, and natural killer cells. Evidence suggests it controls the development of some inflammatory autoimmune illnesses. Initiating biological programmes like inflammation and cell death is its job, and it does it via producing molecular signals. The protein processes a death domain (DD) after binding to TNFR1 and TNFR2 (Jang et al., 2021). This study's arthritic groups exhibited an increase in TNF-alpha levels, which is indicative of the production of inflammatory cytokines. Rats with arthritis showed a significant decrease in this factor after receiving dosages of plant extract. Based on what we know about the biology of type II inflammatory response, interleukin-4 is a cytokine that protects against certain illnesses. Basophils, eosinophils, NKT cells, and CD4 T cells stimulate ILC2, leading to IL-4 synthesis in the body, according to the cellular study based on IL-4 sources. IL-4 is well-known for its role in stabilising lymphocyte activities, which it regulates. A recent research shown that a dosage of 600 mg/kg of the anti-inflammatory cytokine IL-4 significantly increased its levels in rats with arthritis (Raphael et al., 2015). The control of bone metabolism is greatly influenced by osteoprotegerin (OPG). In addition to being an inhibitor of bone resorption, it is a member of the TNF-receptor family. It prevents RANKL from binding to its receptor RANK by binding to its ligand RANKL. The immune system is modulated by the OPG/RANKL system. The hormones control the system by changing the amount of calcium in the cellular environment. The production of RANKL mRNA is triggered by several cytokines, including as IL-1, TNF-alpha, and IL-17, which all contribute to the activation of T cells. When RANKL is secreted, it triggers inflammatory responses, bone resorption, and osteoclastogenesis. In contrast, IL-4, OPG, and IL-10 all work to reduce inflammation, which in turn helps to stop bone resorption and osteoclast development (Boyce and Xing 2007). According to the results of this study, OPG levels are increased following MSRA therapy, and RANKL expression is suppressed as a result of OPG. At the initial stage of inflammation, IL-6 is generated in a local lesion. As it travels through the circulation, it moves from the area of inflammation to the liver. Acute phase proteins such haemoglobin, fibrinogen, α 1-antichymotrypsin, serum amyloid A (SAA), and CRP are induced

by it. Nevertheless, IL-6 plays a role in decreasing albumin, fibronectin, and transferrin serum levels. Amyloidosis and other chronic inflammatory illnesses can develop when SAA levels remain high for an extended period of time. Hypozincemia, caused by the overexpression of the zinc importer ZIP14, is also IL-6's fault. In addition to stimulating the release of platelets from the bone marrow, it aids in the maturation of megakaryocyte cells. Inflammation intensity is indicated by changes in red blood cell and platelet counts as well as acute phase protein levels. The present investigation found that arthritic groups had elevated IL-6 expression. The goal of the piroxicam and plant extract therapy is to reduce tumour necrosis factor alpha levels (Tanaka et al., 2014). A key role of immunoregulatory molecules known as anti-inflammatory cytokines is to modulate the response to pro-inflammatory cytokines. Cytokines IL-4, IL-10, IL-1 receptor antagonist, IL-11, and IL-13 have a significant role in reducing inflammation. Because it inhibits the production of inflammatory cytokines including IL-1, TNF- α , and IL-6, IL-10 is thought of as a very effective anti-inflammatory cytokine. Beyond that, IL-10 regulates the overexpression of endogenous cytokines and the downregulation of pro-inflammatory cytokine receptors on many levels. Further, IL-10 is implicated in the alleviation and potential reversal of neuropathic pain. Patients suffering from chronic pain may have low concentrations of anti-inflammatory cytokines such as IL-4 and IL-10. The results of this study clearly show that administering plant extract to arthritis rats raises their IL-4 levels (Zhang and An 2007).

4

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper

Acknowledgement

Researchers Supporting Project number (RSP2024R397), King Saud University, Riyadh, Saudi Arabia.

Asad Final 23-05-24

ORIGINALITY REPORT

16%

SIMILARITY INDEX

13%

INTERNET SOURCES

9%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

| | | |
|---|---|-----|
| 1 | link.springer.com Internet Source | 4% |
| 2 | www.science.gov Internet Source | 2% |
| 3 | www.mdpi.com Internet Source | 1% |
| 4 | digitallibrary.aau.ac.ae Internet Source | 1% |
| 5 | Submitted to Higher Education Commission Pakistan Student Paper | 1% |
| 6 | escholarship.org Internet Source | 1% |
| 7 | www.frontiersin.org Internet Source | 1% |
| 8 | immunityageing.biomedcentral.com Internet Source | 1% |
| 9 | Submitted to University of Huddersfield Student Paper | <1% |

10

Ammara Saleem, Mohammad Saleem, Muhammad Furqan Akhtar. "Antioxidant, anti-inflammatory and antiarthritic potential of *Moringa oleifera* Lam: An ethnomedicinal plant of Moringaceae family", *South African Journal of Botany*, 2020

Publication

<1 %

11

Ammara Saleem, Mohammad Saleem, Muhammad Furqan Akhtar, Muhammad Shahzad, Shah Jahan. "Polystichum braunii extracts inhibit Complete Freund's adjuvant-induced arthritis via upregulation of I- κ B, IL-4, and IL-10, downregulation of COX-2, PGE2, IL-1 β , IL-6, NF- κ B, and TNF- α , and subsiding oxidative stress", *Inflammopharmacology*, 2020

Publication

<1 %

12

www.researchgate.net

Internet Source

<1 %

13

Chen, Zhejie. "Development of Bilirubin-Based Drug Delivery Systems for the Treatment of Inflammatory Bowel Disease", *University of Macau*, 2023

Publication

<1 %

14

e-century.us

Internet Source

<1 %

15

www.ingentaconnect.com

Internet Source

<1 %

| | | |
|----|--|------|
| 16 | pubmed.ncbi.nlm.nih.gov Internet Source | <1 % |
| 17 | Acevedo, M.. "C-reactive protein and atrial fibrillation: "Evidence for the presence of inflammation in the perpetuation of the arrhythmia"", International Journal of Cardiology, 20060414 Publication | <1 % |
| 18 | Submitted to Hong Kong Baptist University Student Paper | <1 % |
| 19 | research-information.bris.ac.uk Internet Source | <1 % |
| 20 | www.exeley.com Internet Source | <1 % |
| 21 | acamedicine.org Internet Source | <1 % |
| 22 | pdfs.semanticscholar.org Internet Source | <1 % |
| 23 | www.thieme-connect.com Internet Source | <1 % |
| 24 | oak.chosun.ac.kr Internet Source | <1 % |
| 25 | www.mdainternet.com Internet Source | <1 % |

26

topsecretapiaccess.dovepress.com

Internet Source

<1 %

27

www.ajol.info

Internet Source

<1 %

28

Submitted to Massey University

Student Paper

<1 %

29

discovery.ucl.ac.uk

Internet Source

<1 %

30

Ammara Saleem, Mohammad Saleem, Muhammad Furqan Akhtar, Mirza Muhammad Faran Ashraf Baig, Azhar Rasul. " HPLC analysis, cytotoxicity, and safety study of Lam. (wild type) leaf extract ", Journal of Food Biochemistry, 2020

Publication

<1 %

31

Ammara Saleem, Mohammad Saleem, Muhammad Furqan Akhtar, Muhammad Shahzad, Shah Jahan. "Moringa rivae leaf extracts attenuate Complete Freund's adjuvant-induced arthritis in Wistar rats via modulation of inflammatory and oxidative stress biomarkers", Inflammopharmacology, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off