

HY_JKSUS

by Humaira Yasmin

Submission date: 09-Jun-2024 01:02AM (UTC-0400)

Submission ID: 2398516038

File name: Plagiarism_report.docx (1.71M)

Word count: 5673

Character count: 32028

Potential of *Lactobacillus agilis*, *Lactobacillus plantarum*, and *Lactobacillus acidophilus* to enhance wheat growth under drought and heat stress

Allah Nawaz Khan^{1,2}, Muhammad Nadeem Hassan², Rumana Keyani², Zaineb Amir², Mohammad Raish³, Rattandeep Singh⁴, Humaira Yasmin^{2*}

¹Health Services Academy (HSA) Park Road, Chak Shahzad, Islamabad-44000

²Department of Biosciences, COMSATS University Islamabad (CUI), Islamabad, Park road, Chak shazad, 44000 Islamabad, Pakistan.

³Department of Pharmaceutics, College of Pharmacy, King Saud University, 11451 Riyadh, Saudi Arabia

⁴School of Bioengineering and Biosciences, Lovely Professional University, Phagwara, Punjab-144411, India

*Corresponding author: humaira.yasmin@comsats.edu.pk

Declarations

41

Conflicts of interest/Competing interests

The authors have no conflicts of interest to declare.

Ethics approval

Not applicable

Consent to participate

All authors consent to participate in the manuscript publication

55

Consent for publication

All authors approved the manuscript to be published

Availability of data and material

40

The data supporting the conclusions of this article are included within the article. Any queries regarding these data may be directed to the corresponding author.

Funding:

²⁴
This project was funded by Researchers Supporting Project Number (RSPD2024R957) at King Saud University, Riyadh, Saudi Arabia.

This research received no external funding

Acknowledgements:

¹⁵
The authors are grateful to the Researchers Supporting Project Number (RSPD2024R957) at King Saud University, Riyadh, Saudi Arabia.

Author Contributions:

A.Khan: Problem identification and construction of experimental design, M.⁴⁹H.: Provision of resources, R.K: Data retrieval, Z.A.: Data Collection¹⁶ and processing, H.Y.: Data analysis, data interpretation and manuscript preparation. MR and RS analyzed the data and revised the language of the manuscript. All authors have read and agreed to publish the version of manuscript.

Abstract

Drought and heat stress affect plant functioning and turgor pressure by reducing water content and affecting cell growth, and together, they can cause severe damage. *Lactobacillus agilis*, *Lactobacillus acidophilus*, and *Lactobacillus plantarum* strains were identified and evaluated for growth-promoting properties via Petri plate test and pot experiment. Wheat seeds were exposed to heat and drought separately and in combination with the bacterial isolates primed for the seeds. In drought-stressed plants, after treatment with *L. agilis*, there was an elevation in carotenoids by 37% and chlorophyll a and b levels by 55% and 10%, respectively. Similarly, heat-stressed plants, after treatment with *L. agilis*, showed an elevation in carotenoids by 55% and chlorophyll a and b levels by 70% and 82%, respectively. A further 5% increase in the SOD concentration was noted after the treatment of *L. agilis* in plants with drought stress. A significant increase (24%) in the APX level was also observed after the treatment of *L. agilis* in plants with heat stress. In plants with drought stress, a rise in CAT concentration was observed after the treatments of *L. agilis* and *L. plantarum* with 33% and 11%, respectively. In plants with drought and heat stresses, the level of POD in combined treatments increased by 39% and 41%, respectively. The study reveals that rhizospheres can serve as reservoirs of plant growth-promoting bacteria, potentially replacing chemical fertilizers and promoting plant growth in stressful conditions.

Key Words: Medicinal plants, Rhizosphere, PGPR, Drought stress, Heat stress

1. Introduction

Aside from the increased interest in food, the scarcity of natural resources and high agricultural production are posing significant challenges in both developing and established countries. However, in today's world, crop yield has not shown significant improvement despite increasing compost input (Pii *et al.*, 2015). By enhancing plant nutrition and shielding them against microbes, beneficial bacteria found in the microbiome of plant roots provide essential benefits to the plant. PGPR, a type of plant growth-promoting rhizobacteria, colonize plant roots, promoting plant growth through the production or modification of growth regulators and a symbiotic N₂ fixation process. Plant growth-promoting rhizobacteria additionally produce siderophores, cyanide, and anti-microbials to provide defense against phytopathogenic microorganisms (Bulgarelli *et al.*, 2013). A diverse number of PGPRs activate plant growth and advancement by inhibiting the action and growth of plant pathogens. Different PGPRs restrain infection indirectly by triggering induced systematic resistance (ISR), which is a technique that gets plant-borne protection systems ready to respond quickly and stronger against abiotic stress (Conrath *et al.*, 2015), as well as facilitating the uptake of essential nutrients from the soil. Rhizobium species are restoring the development of root nodules by settling air nitrogen in those nodules and making it accessible to plants.

Important nutrients generated by PGPR are related with soil phosphate, which promotes iron and nitrate uptake by plant systems (Wintermans *et al.*, 2016). Sustainable agriculture has a global challenge due to the water shortage, as around 215 million hectares (MHA) of wheat are farmed worldwide using irrigation water (12%) available (Aaboud *et al.*, 2018). Global climate change and uneven rainfall patterns resulted in a significant global rise in drought stress conditions. Water scarcity and heat stresses minimize crop yields, ultimately directing farmers' poor earnings (Daryanto *et al.*, 2016; Zhao *et al.*, 2017). The average worldwide temperature is anticipated to rise approximately 2 °C in the following 50 years (Ahmed *et al.*, 2017). High temperatures retard growth, cell division, and cell elongation in plants, in addition to reducing root growth, number, and diameter.

As harsh and extreme environmental conditions prevail in different areas, it is necessary to cope with conditions such as heat and drought. These stressful conditions affect the yield of crops. So, the key goal of the current research remained to examine the potential of *L. agilis*, *L. plantarum*, and *L. acidophilus* for wheat germination and biomass growth. Secondly, individual and collective strains were explored for their involvement in mitigating the impacts of stressors (drought and heat) on wheat plants. The morphological, physiological, and

biochemical responses of stressed wheat by applying *L. agilis*, *L. plantarum*, and *L. acidophilus* were also analyzed in the current study.

2. Methodology

The study used three repetitions and a randomized block design to be conducted in a greenhouse conditions with five treatments in thirteen Petri plates and soil-filled Pots. The treatments are given in table 01.

Table 1: Treatments involved in the Petri plates and pots experiment

Abbreviations	Treatments Abbreviations
²⁵ C	Control
H	Heat Stress
D	Drought Stress
D+ H	Drought Stress + Heat Stress
<i>L. ag</i> + D	<i>Lactobacillus agilis</i> + Drought
<i>L. ag</i> + H	<i>Lactobacillus agilis</i> + Heat
<i>L. ag</i> + D + H	<i>Lactobacillus agilis</i> + Drought + Heat
<i>L.pla</i> +D	<i>Lactobacillus plantarum</i> + Drought
<i>L.pla</i> +H	<i>Lactobacillus plantarum</i> + Heat
<i>L.pla</i> +D + H	<i>Lactobacillus plantarum</i> + Drought + Heat
<i>L. ag</i> + <i>L.pla</i> + D	<i>Lactobacillus agilis</i> + <i>Lactobacillus plantarum</i> + Drought
<i>L. ag</i> + <i>L.pla</i> + H	<i>Lactobacillus agilis</i> + <i>Lactobacillus plantarum</i> + Heat
<i>L. ag</i> + <i>L.pla</i> + D + H	<i>Lactobacillus agilis</i> + <i>Lactobacillus plantarum</i> +Drought + Heat

2.1 Isolation and morphological examination of *Lactobacillus* isolates

⁴⁸ *Lactobacillus agilis* (*L. agilis*) strain NMCC-15 (MK614016) was used, as already reported in our previous study (Khan *et al.*, 2021). Each (1 g) rhizosphere soil sample collected from *Ficus carica* and *Azadirachta indica* was serially diluted to 10^{-1} , 10^{-2} , 10^{-3} , 10^{-4} , and 10^{-5} according to the protocol of Khan *et al.* (2021) for the identification of *Lactobacillus plantarum* (*L. plantarum*) and *Lactobacillus acidophilus* (*L. acidophilus*) strains. To inoculate and spread each dilution on Petri plates, MRS agar media (selective for *Lactobacillus* isolates) at pH 5.4 was prepared (De Man *et al.*, 1960) and pure cultures were obtained by sub-culturing on MRS agar Petri plates and incubated the cultured plates at 32°C for 24 h. Bergey's handbook was used to define the morphological characteristics, which include colony color, form, margin, elevation, texture, and size (Cossart, 1987). In order to identify the strains of *L. plantarum* and *L. acidophilus*, several conventional and biochemical tests, such as gas generation, estimation of catalase, oxidase, urease, methyl red, citrate, triple sugar iron, and fermentation were applied. Phase-contrast microscope (Phase Contrast 2, Nikon, Tokyo, Japan), and Gram staining of the bacterial isolates was carried out to analyze the specific type of bacteria. The interpretation of the data was based on "Bergey's Manual of Determinative Bacteriology" (Holt *et al.*, 1984).

2.2 Collection of seeds and germination

Johar-16 wheat (*Triticum aestivum* L.) seeds were provided by the National Agriculture Research Centre (NARC), Islamabad. The seeds were decontaminated for one minute via a 70% ethanol solution, and then five times washed with distilled water. Before conducting the studies, the seeds were given an overnight period to get vernalized.

2.3 Preparation of the bacterial suspension

The bacterial strains of *L. agilis*, *L. plantarum*, and *L. acidophilus* were used to evaluate their potential as growth promoters under heat and drought stress. Luria broth (LB) media of 25 mL was prepared aseptically and added to sterilized flacon tubes in a culture hood for each bacterial inoculum. After the sterilized medium reached room temperature, it was transferred into falcon tubes within the biosafety cabinet. The isolates of *L. agilis*, *L. plantarum*, and *L. acidophilus* were then individually inoculated in falcon tubes with 100 µL of each and labeled according to the strain names. Tubes were sealed and placed in a shaking incubator at 30°C for 24 h. After 24 h of incubation, falcon tubes were picked and centrifuged at 8000 rpm for 15 min. The pellet formed on the bottom of the tubes was stored, and the supernatant was removed.

Pallets of each *L. agilis*, *L. plantarum*, and *L. acidophilus* were suspended in 10 µL of autoclaved water and shaken properly.

2.4 Bacterial isolates for plant growth enhancement

The bacterial isolates were evaluated in the laboratory of applied microbiology and biotechnology at COMSATS University Islamabad for their potential to promote plant growth. Before being planted in pots, seeds were first cultured in glass Petri plates with aluminum and sterile cotton to retain water. Seeds were primed for three to four hours prior to sprouting with strains of *L. agilis*, *L. plantarum*, and *L. acidophilus* and were then left to grow for two to three days at room temperature and in the dark (Khalid *et al.*, 2004). Each day, distilled water was sprayed on the seedlings. After 15 days, wheat plants were collected, and analyzed for promptness index (PI), seedling vigor index (SVI), germination percentage (GP), and germination index (GI) as part of a biomass study.

2.4.1 Petri plates experiment under drought stress condition

Seeds were primed with solutions containing *L. agilis*, *L. plantarum*, and *L. acidophilus* strains for 3–4 h. A single Petri plate contained about 10–15 seeds for each treatment. Average room temperature and dark conditions were provided for an initial 2-3 days until the seeds germinated. Distilled water was regularly provided for 15 days. The wheat harvesting process was performed according to the protocol of Xiang *et al.* (2006) after 15 days, and a biomass study was done, which involved the analysis of lateral root (LR), fresh weight (FW), root length (RL), dry weight (DW), shoot length (SL), and leaf area (LA). The best PGPRs were used in the final pot experiment that were selected in the Petri plates experiment on the basis of their potential growth-promoting properties.

2.4.2 Pot assays

Pot assays were carried out at an average temperature of 15 °C and 64% humidity in the greenhouse at COMSATS University in Islamabad (33.7294 °N, 73.0931 °E). In order to execute the experiment, natural light was ensured. Each 10x10 cm pot included 1.5 kg of autoclaved dried soil (3:1, soil:sand), along with holes for drainage in the bottom. Each pot had a plastic plate underneath it to catch the leachate, which was then watered on the surface of soil. Eight (08) seeds primed in triplets with bacterial solutions were planted in each pot containing autoclaved soil. Using distilled water, the soil's moisture content was kept at about 60%. Plants at the three-leaf stage were exposed to drought and heat stress and were harvested

after 15 days of treatment, in accordance with Zhang *et al.* (2009) established protocol. Table 1 lists the specifics of the randomized trial design. Biomass analysis of RL, SL, LA, F.W., D.W., and GP was performed.

2.5 Biomass study of wheat

A biomass study included the evaluation of characteristics such as root length, fresh weight, shoot length, dry weight, relative water content, germination percentage, germination index, seedling vigor index, and promptness index. A measuring scale and weighing balance were used to measure the harvested plants for RL, SL, LA, F.W, D.W, and GP (Tian *et al.*, 2015). Given below were the formulas used for these measurements:

Germination percentage = (Number of germinated seeds / numbers of total seeds) x 100

Germination index (Gt/Tt) = where, Gt is referred to the number of germinated seeds on Tth day and Tt refers to the total number of days.

Seedling vigor index (SVI) = (seedling length x germination percent) / 100

Promptness index (PI) = $nd_2 (1.00) + nd_4 (0.75) + nd_6 (0.5) + nd_8 (0.25)$ where, n refers to the germinated seeds at the respective days like $d_2 = 2^{nd}$ day.

2.6 Assessing photosynthetic pigments

Fresh leaves samples of the wheat plants weighing 0.5 g were crushed and placed in test tubes to measure the concentration of carotenoids and chlorophyll *a* and *b*. For this purpose, ten (10) milliliters of dimethyl sulfoxide (DMSO) were added to each test tube, and was left at room temperature/ in water bath for 4 h at 65°C. Following this time, the absorbance of the extracted supernatant was measured at 480, 648, and 665 nm respectively for carotenoids and chlorophyll *a* and *b*. Every computation was completed in compliance with the Chappelle *et al.* (1992) protocol.

2.7 Estimation of antioxidative enzymes

2.7.1 Preparation of crude enzyme extracts

The plant's leaves were homogenized in mortar and pestle using liquid nitrogen in an ice-cold sodium phosphate buffer (50-mM, 7.2 pH) and polyvinylpyrrolidone (1% , w/v), then centrifuged at 6,708 g for 20 min. The supernatant was used for enzymatic assays.

2.7.2 Catalase (CAT) assessment

El-Shabrawi *et al.* (2010) protocols were modified in order to measure the catalase activity. A reaction mixture consisting of 0.1 mL of 300 mM hydrogen peroxide, 2.8 mL of

100% diluted 50 mM phosphate buffer, and crude enzyme solution of 0.1 mL was prepared to carry out the catalase test. The absorbance of the combination was measured via spectrophotometry at 240 nm on intervals of 30 seconds (Kodydková *et al.*, 2014).

2.7.3 Superoxide dismutase (SOD) assessment

Superoxide dismutase activity was assessed after adding 100 µL of crude enzyme extract to 50 mM sodium phosphate, a buffer solution of 7.6 pH, 50 mM sodium carbonate, 0.1 mM EDTA, 50 mM nitro blue tetrazolium, and 10 µM riboflavin to make a total volume of 3.0 mL. This method confirmed the SOD activity after bringing halt in the nitro blue tetrazolium photo-reduction. The reaction mixture was placed at room temperature for 15 min under white light. Following incubation period absorbance was measured at 560 nm via spectrophotometer (Ighodaro and Akinloye, 2018). The values were measured and compared with a crude-extract-free control group.

2.7.4 Peroxidases (POD) assessment

A reaction mixture was prepared that comprised of 100 mM sodium phosphate buffer (pH 7-8), 4-methyl catechol (5 mM), hydrogen peroxide (5 mM), and 500 µL of crude enzyme extract for the determination of POD. The total volume of the mixture was raised to 3.0 mL, and via spectrophotometer, absorbance was measured at 420 nm (Onsa *et al.*, 2004).

2.7.5 Ascorbate-peroxidase (APX) assessment

The oxidation of guaiacol was measured at 470 nm for the confirmation of peroxidase activity (Rao, 1996). A volume of 100 µL of diluted crude enzyme extract, potassium phosphate buffer of 0.9 mL (100 mM), 1 mM EDTA, and 0.1 mL of 10 mM guaiacol were mixed to carry out the experiment. A volume of 100 µL containing 10 mM hydrogen peroxidase was utilized to start the reaction. A 470 m absorbance check was performed using the spectrophotometer at 15-second intervals for up to two minutes. The enzyme's activity was expressed in mg/min.

2.8 Statistical examination

Statistix (version 8.1) software was used to perform statistical study on the data set. ANOVA (analysis of variance) was the method used to examine the variation in the data set in triplicates. Between each treatment's three replicates, the standard error was computed. Using the least significant difference (LSD) at $p = 0.05$, the treatments were compared.

3. Results

3.1 Isolation

Bacterial isolates were obtained after subculturing the samples on MRS agar media, that were all biochemically and morphologically distinct. The color (creamy/white), form (circular/rods), and surface texture (smooth/rough) of all the isolates varied considerably. Bacteria with Gram-positive, catalase-negative, oxidase negative and positive for citrate utilization were found in all the samples which are the typical properties of lactic acid bacteria. *L. plantarum* and *L. acidophilus* strains were identified by using [Bergey's manual](#) and [Bergey's Manual of Determinative Bacteriology](#).

3.2 Bacterial isolates for plant growth enhancement

Wheat showed a 40–65% increase in rate of germination with PGPR priming. The maximum germination rate (65%) was observed in the *L. agilis* treatment by comparison with the control (without PGPR treatment). In comparison with the control, *L. plantarum* and *L. acidophilus*, also showed a prominent increase in seedling vigor index (SVI), germination index (GI), promptness index (PI), and germination percentage (GP). Maximum effectiveness was seen in the treatment of *L. agilis* such as GP 20%, GI 21%, PI 20%, and SVI 60% when compared to the controls seedling (table 2).

Table 2: Screening of *Lactobacillus agilis*, *Lactobacillus plantarum* and *Lactobacillus acidophilus*

Treatments	GP	GI	PI	SVI
Control	^C 83.33 ±0.12	^B 2.08 ±0.11	^B 6.25 ±0.13	^C 7.12 ±0.14
<i>L. agilis</i>	^A 99 ±0.12	^A 2.5 ±0.11	^A 7.5 ±0.13	^A 11.46 ±0.14
<i>L. acido</i>	^B 96.67 ±0.12	^A 2.41 ±0.11	^A 7.25 ±0.13	^B 9.99 ±0.14
<i>L. pla</i>	^A 99 ±0.12	^A 2.5 ±0.11	^A 7.5 ±0.13	^B 10.97 ±0.14

Alpha 0.05. Between the means, there is significant statistical difference

3.3 Screening of PGPR against drought and heat stress

Comparing the stressed plants to the control, the stressed plants displayed substantial reductions in leaf area (LA), shoot length (SL), lateral root (LR), dry weight (DW), fresh weight (FW), and shoot length (RL). In the absence of PGPR priming, only 35% of the wheat seeds were germinated. In contrast to the control (without PGPR treatment), plants treated with *L. agilis* showed 50% and 60% higher germination rates in stressed and non-stressed plants, respectively. While *L. acidophilus* showed decreased activity in stressful conditions, the other strain of bacteria, *L. plantarum*, demonstrated a significant growth rate with better seed germination potential. Based on the plant growth promotion abilities, *L. agilis* and *L. plantarum* were selected for the pot experiment conducted on the plants (table 3; figure 1).

Table 3: Screening of *Lactobacillus agilis*, *Lactobacillus plantarum* and *Lactobacillus acidophilus* under stress condition

Treatments	SL (cm)	RL (cm)	FW (g)	DW (g)	LR (cm)	LA (cm ²)
Control	8.63 ^B ±0.12	3.40 ^E ±0.11	0.37 ^C ±0.07	0.10 ^C ±0.07	4 ^A ±0.14	8.23 ^C ±0.13
C (drought)	2.10 ^E ±0.12	5.50 ^{CD} ±0.11	0.12 ^E ±0.07	0.03 ^D ±0.01	2.3 ^B ±0.14	1.66 ^G ±0.13
<i>L. agilis</i>	11.46 ^A ±0.12	6.83 ^{BC} ±0.11	0.50 ^A ±0.07	0.15 ^A ±0.03	4 ^A ±0.14	11.1 ^A ±0.13
<i>L. agilis</i> (drought)	8.50 ^B ±0.12	9.83 ^A ±0.11	0.24 ^D ±0.07	0.09 ^E ±0.04	2 ^B ±0.14	6.12 ^D ±0.13
<i>L. acido</i>	6.13 ^A ±0.12	4.63 ^D ±0.11	0.15 ^E ±0.07	0.09 ^E ±0.04	3 ^{AB} ±0.14	5.5 ^E ±0.13
<i>L. acido</i> (drought)	4.06 ^D ±0.12	5.8 ^{CD} ±0.11	0.09 ^F ±0.07	0.02 ^D ±0.01	2 ^B ±0.14	3.73 ^F ±0.13
<i>L. pla</i>	10.96 ^A ±0.12	6.76 ^C ±0.11	0.42 ^B ±0.07	0.12 ^B ±0.01	2.6 ^{AB} ±0.14	10.3 ^B ±0.13

<i>L. pla</i> (drought)	7.33 ^C ±0.12	7.56 ^B ±0.11	0.2 ^D ±0.07	0.06 ^E ±0.0 3	2.3 ^B ±0.14	4.86 ^E ±0.1 3
----------------------------	-------------------------	-------------------------	------------------------	-----------------------------	------------------------	-----------------------------

Alpha 0.05. Between the means, there is significant statistical difference




Figure 1: Shows Petri plate assays of stress induce wheat plant and *L. agilis* and *L. plantarum*

C= Control, D= drought

3.4 Pot experiment

All treatments including primed wheat seeds containing strains of *L. agilis* and *L. plantarum* demonstrated a significant increase in the germination rate when compared to unprimed seeds. Drought and heat stress showed significant decrease in RL, SL, LA, FW, DW in comparison with the unstressed controlled plants. In stressed plants, *L. plantarum* treatment revealed a significant growth in RL, SL, LA, FW, DW ranges from 7 to 20%. Whereas, in case of *L. agilis* treated plants the values ranged from 14 to 25%. A combined treatment of *L. agilis* and *L. plantarum* exhibited a maximum rise range from 15 to 70% (table 4).

Table 4: Pot experiment after the treatment of *Lactobacillus agilis* and *Lactobacillus plantarum* under stress and non-stressed conditions

Treatments	RL (cm)	SL (cm)	LA (cm ²)	F.W (g)	D.W (g)	GP
Control	13 ^{BC} ±0.1	18.6 ^{FG} ±0.2	12.6 ^{CD} ±0.1	0.37 ^A ±0.2	0.25 ^B ±0.1	100 ^A ±0.3
Heat	12 ^C ±0.1	21.6 ^{BCD} ±0.2	15.6 ^{BC} ±0.1	0.18 ^C ±0.2	0.07 ^D ±0.1	96.6 ^B ±0.3
Drought	11 ^C ±0.1	16 ^G ±0.2	10 ^D ±0.13	0.28 ^{AB} ±0.2	0.183 ^C ±0.11	100 ^A ±0.3
D+H	10 ^C ±0.1	16.6 ^{FG} ±0.2	10 ^D ±0.13	0.15 ^{DE} ±0.2	0.11 ^C ±0.1	100 ^A ±0.3
<i>L. agilis</i> +D	16 ^A ±0.1	24.6 ^{AB} ±0.2	18. ^A 3±0.1	0.31 ^{AB} ±0.2	0.22 ^B ±0.1	100 ^A ±0.3
<i>L. agilis</i> +H	13 ^{BC} ±0.1	20 ^{CDE} ±0.2	12.6 ^{CD} ±0.1	0.16 ^{DE} ±0.2	0.12 ^C ±0.1	100 ^A ±0.3
<i>L. agilis</i> +D+H	17 ^A ±0.1	26.3 ^A ±0.2	16.6 ^{AB} ±0.1	0.07 ^{EF} ±0.2	0.043 ^D ±0.11	100 ^A ±0.3
<i>L. pla</i> +H	12 ^C ±0.1	19 ^G ±0.2	13 ^{CD} ±0.1	0.02 ^F ±0.2	0.01 ^{CD} ±0.11	96.6 ^B ±0.3
<i>L. pla</i> +D	11 ^C ±0.1	17 ^{FG} ±0.2	10.6 ^D ±0.1	0.16 ^{CD} ±0.2	0.123 ^C ±0.11	96.6 ^B ±0.3
<i>L. pla</i> +D+H	11 ^C ±0.1	19.6 ^{DEF} ±0.2	12 ^{CD} ±0.1	0.073 ^{DEF} ±0.2	0.043 ^D ±0.11	96.6 ^B ±0.3

<i>L. agilis</i> + <i>L. pla</i> +H	15 ^A ±0.1	22.3 ^{BC} ±0.2	15.6 ^{ABC} ± 0.13	0.27 ^B ±0.2	0.20 ^{BD} ±0.	100 ^A ±0.
<i>L. agilis</i> + <i>L. pla</i> +D	15 ^{AB} ±0. 1	22 ^{BC} ±0.2	15 ^{BC} ±0.1 3	0.16 ^C ±0.2	0.08 ^D ±0.1 1	100 ^A ±0. 3

Alpha 0.05. Between the means, there is significant statistical difference

3.5 *L. agilis* and *L. plantarum* treatments impact on photosynthetic pigments in stressed plants

Chlorophyll *a*, *b* and carotenoid concentrations in wheat plants were decreased after heat and drought stress. Compared with the non-stressed control, drought stress decreased carotenoid to 6%, chlorophyll *a* to 35%, and chlorophyll *b* to 14%. In contrast to the non-stressed control, heat stress reduced the levels of carotenoid by 33%, chlorophyll *a* by 63%, and chlorophyll *b* by 81% (figure 2). All of the treatments notably increased the amounts of carotenoid, chlorophyll *a*, and chlorophyll *b* in the unstressed plants. When plants under drought stress were treated with *L. agilis*, levels of carotenoids increased by 37%, and levels of chlorophyll *a* and chlorophyll *b* also increased by 55% and 10%, respectively, in comparison to plants not under drought stress. After being treated with *L. agilis*, heat-stressed plants displayed increased levels of carotenoids by 55% and chlorophyll *a* and chlorophyll *b* by 70% and 82%, respectively, in comparison to the control group. Plants exposed to drought stress exhibited increased levels of carotenoids (46%) and chlorophyll *a* and chlorophyll *b* (54 and 19%) respectively, following treatment with *L. plantarum*, as compared to the control group under drought stress. Figure 3 illustrates the increase in carotenoids (59%), chlorophyll *a* (73%), and chlorophyll *b* (85%), in heat-stressed plants treated with *L. plantarum* as opposed to drought-stressed control. Plants under drought and heat stress exhibited increased levels of carotenoids (58%), as well as chlorophyll *a* (68%) and chlorophyll *b* (52%), following treatment with *L. agilis*. Drought and heat-stressed plants, when treated with *L. plantarum*, showed elevations in the carotenoids by 31% and chlorophyll *a* and chlorophyll *b* levels by 22% and 28%, respectively when compared with drought- and heat-stressed control. Drought-stressed plants, after combined treatment with *L. agilis* and *L. plantarum*, showed 37% elevation in the carotenoids and 44% chlorophyll *a* levels when compared to the drought-stressed control group. Heat-stressed plants, after being treated together with *L. agilis* and *L. plantarum*, showed 60% elevations in the carotenoids concentration and 62%, 72% increase in chlorophyll *a* and chlorophyll *b* levels, respectively, when compared with the heat-stress

control group. In drought- and heat-stressed plants, after combined treatment with *L. agilis* and *L. plantarum*, the concentration of carotenoids increased to 71% and chlorophyll *a* and *b* levels to raised to 78% and 69% respectively, after being compared with control of drought and heat-stress (figure 2).




Figure 2: Shows contents of carotenoid, chlorophyll *a* and *b* after stress, *L. agilis* and *L. plantarum* treatments

Alpha 0.05. Between the means, there is significant statistical difference

3.6 Effects of *L. agilis* and *L. plantarum* treatments on antioxidant enzyme activities in stressed plants

After the wheat plants exposed to drought and heat stress exhibited 21% and 29% elevation in the POD, respectively, in comparison to non-stressed plants (figure 3). In plants with drought stress, a further rise in the CAT concentration was noted after the treatments of *L. agilis* and *L. plantarum* with 33% and 11%, respectively, as compared with the drought-stressed control group. In plants with heat stress, a notable increase in the CAT concentration was observed after the treatments of *L. agilis* and *L. plantarum* with 19% and 20%,

respectively, as compared with the heat-stressed control group. In plants with drought and heat stress, the level of POD in combined treatments increased by 39% and 41%, respectively. After comparison to non-stressed plants, wheat plants exposed to heat and drought conditions showed elevations in the CAT of 26% and 21%, respectively (figure 3). In plants subjected to drought stress, an increase in the CAT level was noted after treatments of *L. agilis* and *L. plantarum* with 40% and 41%, respectively, as compared to drought-stressed control group. Similarly, plants under heat stress showed a significant raise in the CAT level after treatments with *L. agilis* at 29% and *L. plantarum* at 5% as compared with the heat-stressed control group. Exposed wheat plants to drought and heat stress exhibited 93% and 91% elevation in the SOD, respectively, compared with the non-stressed plants (figure 3). When *L. agilis* was applied to drought-stressed plants, there was an additional 5% rise in SOD content as compared to the control group. After comparison to the heat-stressed control group, there was a notable rise (42%) in the SOD level in heat-stressed plants following the application with *L. agilis*. Exposed wheat plants with drought and heat stress exhibited 91% and 93% elevation in the APX, respectively, compared with the non-stressed plants. After *L. agilis* treatment, there was a further 25% increase in the APX level in drought-stressed plants after being compared with the drought-stressed control group. After comparison to the heat-stressed control, there was a notable rise (24%) in the APX level in plants that had received *L. agilis* treatment (figure 3).




Figure 3: Activities of peroxidase, catalase, super oxide dismutase and ascorbate peroxidase in stress, *L. agilis*, *L. plantarum* treatments

Alpha 0.05. Between the means, there is significant statistical difference

4. Discussion

The selected *Ficus carica*, *Azadirachta indica*, and *Ocimum tenuiflorum* plants have been used for medicinal purposes a long time ago, and most of the researchers have concluded the role of extracellular metabolites in the rhizosphere region of the plants to promote plant growth and other potent characteristics. Technologies based on microbe are becoming common, producing nutrients and plant-growth promoting constituents in the rhizospheric plants region (Naamala, 2020). As revealed from the current study, inoculation of seeds with Rhizospheric bacterial isolates significantly enhanced the growth and yield of the wheat plant when compared to non-treatment controls. Results of the current study are correlated to a study that analysed PGPRs assistance in maintaining plant survival under stress conditions by producing various secondary metabolites, phytohormones and enzymes (Mishra, 2018). The

study found that *L. agilis* and *L. plantarum* significantly affected wheat growth and yield, with direct exposure to drought and heat shocks increasing antioxidant (POD, SOD, CAT, APX) levels.

In the current study, drought and heat exposure significantly reduced the germination, shoot length (SL), root length (RL), leaf area (LA), fresh weight (FW), and dry weight (DW) in untreated plants, which is related to the study of Saadaoui *et al.* (2023), in which it is reported that abiotic stresses severely affected the germination and biomass of wheat. Current findings illustrated the momentous rise in the contents of chlorophyll after PGPR treatments, that also correlated to the study of Neshat *et al.* (2022), which depicted the linkage between the PGPR and chlorophyll content, enhanced antioxidant levels, and better plant growth under the stress condition. In this study, stresses of drought and heat to the treatments, reduced the photosynthetic pigments significantly in leaves, as revealed by the lower values of chlorophyll *a*, *b*, and carotenoids, which related to a study in which chlorophyll synthesis was directly affected by drought and heat stresses through defective electron transport chain mechanisms (Abbas *et al.*, 2018). All the *L. agilis* and *L. plantarum* inoculation significantly enhanced the photosynthetic pigments in both stressed and non-stressed plants, which suggested that plant and bacterial enzymes are responsible for the synthesis of chlorophyll pigments along with essential plant nutrients. The current finding correlated with the research led by Vejan *et al.* (2016), who found that chlorophyll *a* and *b* contents increased after the application of *B. subtilis* S4 in stressed plants. In this study, the treatments of *L. agilis* and *L. plantarum* showed significant enhancements in the chlorophyll *a*, *b*, carotenoid contents, and antioxidant enzymes in wheat plants.

5. Conclusion

Current findings suggest that *L. agilis*, *L. plantarum*, and *L. acidophilus* enhanced the wheat plant's vigor and biomass. However, *L. agilis* and *L. plantarum* showed significant impact individually and in consortium to mitigate the adversative effects of drought and heat stresses on wheat plants. Moreover, *L. agilis* and *L. plantarum*-mediated stress alleviation was attributed to the modulation of the morphological, physiological, and biochemical responses of stressed wheat. These strains can be tested in field conditions to validate their potential as biofertilizers.

6. References

Aaboud, M., Aad, G., Abbott, B., Abeloos, B., Abidi, S. H., AbouZeid, O. S., Abraham, N. L., Abramowicz, H., Abreu, H., Abreu, R., & others. (2018). Search for dark matter and other new phenomena in events with an energetic jet and large missing transverse momentum using the ATLAS detector. *Journal of High Energy Physics*, 2018(1), 1–53.

Abbas, S. K., Hassan, H. A., Asif, J., Ahmed, B., Haider, S. S., & others. (2018). Integration of TTF, UTAUT, and ITM for mobile Banking Adoption. *International Journal of Advanced Engineering, Management and Science (IJAEMS) Vol-4, Issue-5*.

Ahmed, N., Brawley, V., Hegde, M., Bielamowicz, K., Kalra, M., Landi, D., Robertson, C., Gray, T. L., Diouf, O., Wakefield, A., & others. (2017). HER2-specific chimeric antigen receptor--modified virus-specific T cells for progressive glioblastoma: a phase 1 dose-escalation trial. *JAMA Oncology*, 3(8), 1094–1101.

Bulgarelli, D., Schlaeppli, K., Spaepen, S., Van Themaat, E. V. L., & Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology*, 64, 807–838.

Chappelle, E. W., Kim, M. S., & McMurtrey III, J. E. (1992). Ratio analysis of reflectance spectra (RARS): an algorithm for the remote estimation of the concentrations of chlorophyll a, chlorophyll b, and carotenoids in soybean leaves. *Remote Sensing of Environment*, 39(3), 239–247.

Conrath, U., Beckers, G. J. M., Langenbach, C. J. G., Jaskiewicz, M. R., & others. (2015). Priming for enhanced defense. *Annu. Rev. Phytopathol*, 53(1), 97–119.

Cossart Y. (1987). Bergey's Manual of Systematic Bacteriology Volume 2. *Pathology*, 19:324. doi: 10.1016/S0031-3025(16)36764-2.

Daryanto, S., Wang, L., & Jacinthe, P.-A. (2016). Global synthesis of drought effects on maize and wheat production. *PloS One*, 11(5), e0156362.

De Man, J. C., Rogosa, D., & Sharpe, M. E. (1960). A medium for the cultivation of lactobacilli. *Journal of applied Bacteriology*, 23(1), 130-135.

El-Shabrawi, H., Kumar, B., Kaul, T., Reddy, M. K., Singla-Pareek, S. L., & Sopory, S. K. (2010). Redox homeostasis, antioxidant defense, and methylglyoxal detoxification as markers for salt tolerance in Pokkali rice. *Protoplasma*, 245(1), 85-96.

Holt J.G., Krieg N.R., Sneath P.H.A. (1994). *Bergey's Manual of Determinative Bacteriology*. The Williams and Wilkins Co.; Baltimore, MD, USA.

Ighodaro, O. M., & Akinloye, O. A. (2018). First line defence antioxidants-superoxide dismutase (SOD), catalase (CAT) and glutathione peroxidase (GPX): Their fundamental role in the entire antioxidant defence grid. *Alexandria Journal of Medicine*, 54(4), 287–293.

Khalid, A., Arshad, M., & Zahir, Z. A. (2004). Screening plant growth-promoting rhizobacteria for improving growth and yield of wheat. *Journal of Applied Microbiology*, 96(3), 473–480.

Khan, A. N., Yasmin, H., Ghazanfar, S., Hassan, M. N., Keyani, R., Khan, I., ... & Ahmad, A. (2021). Antagonistic, anti-oxidant, anti-inflammatory and anti-diabetic probiotic potential of *Lactobacillus agilis* isolated from the rhizosphere of the medicinal plants. *Saudi J. Biol. Sci.*, 28(11), 6069-6076.

Kodydková, J., Vávrová, L., Kocik, M., & Zak, A. (2014). Human catalase, its polymorphisms, regulation and changes of its activity in different diseases. *Folia Biologica*, 60(4), 153.

Mishra, J., Fatima, T., & Arora, N. K. (2018). Role of secondary metabolites from plant growth-promoting rhizobacteria in combating salinity stress. In *Plant microbiome: stress response* (pp. 127–163). Springer.

Neshat, M., Abbasi, A., Hosseinzadeh, A., Sarikhani, M. R., Dadashi Chavan, D., & Rasoulnia, A. (2022). Plant growth promoting bacteria (PGPR) induce antioxidant tolerance against salinity stress through biochemical and physiological mechanisms. *Physiology and Molecular Biology of Plants*, 28(2), 347-361.

Naamala, J., & Smith, D. L. (2020). Relevance of plant growth promoting microorganisms and their derived compounds, in the face of climate change. *Agronomy*, 10(8), 1179.

Onsa, G. H., bin Saari, N., Selamat, J., & Bakar, J. (2004). Purification and characterization of membrane-bound peroxidases from Metroxylon sagu. *Food Chemistry*, 85(3), 365–376.

Pii, Y., Mimmo, T., Tomasi, N., Terzano, R., Cesco, S., & Crecchio, C. (2015). Microbial interactions in the rhizosphere: beneficial influences of plant growth-promoting rhizobacteria on nutrient acquisition process. A review. *Biology and Fertility of Soils*, 51(4), 403–415.

Rao, G. N. (1996). New diet (NTP-2000) for rats in the National Toxicology Program toxicity and carcinogenicity studies. *Fundamental and Applied Toxicology*, 32(1), 102–108.

Saadaoui, W., Tarchoun, N., Msetra, I., Pavli, O., Falleh, H., Ayed, C., ... & Petropoulos, S. A. (2023). Effects of drought stress induced by D-Mannitol on the germination and early seedling growth traits, physiological parameters and phytochemicals content of Tunisian squash (*Cucurbita maxima* Duch.) landraces. *Frontiers in Plant Science*, 14, 1215394.

Tian, J., Chow, J. C., Cao, J., Han, Y., Ni, H., Chen, L.-W. A., Wang, X., Huang, R., Moosmu, H., Watson, J. G., & others. (2015). A biomass combustion chamber: Design, evaluation, and a case study of wheat straw combustion emission tests. *Aerosol and Air Quality Research*, 15(5), 2104–2114.

Vejan, P., Abdullah, R., Khadiran, T., Ismail, S., & Nasrulhaq Boyce, A. (2016). Role of plant growth promoting rhizobacteria in agricultural sustainability—a review. *Molecules*, 21(5), 573.

Wintermans, P. C. A., Bakker, P. A. H. M., & Pieterse, C. M. J. (2016). Natural genetic variation in *Arabidopsis* for responsiveness to plant growth-promoting rhizobacteria. *Plant Molecular Biology*, 90(6), 623–634.

Xiong, L., Wang, R.-G., Mao, G., & Koczan, J. M. (2006). Identification of drought tolerance determinants by genetic analysis of root response to drought stress and abscisic acid. *Plant Physiology*, 142(3), 1065–1074.

Zhang, G. S., Zhang, F. X., & Li, X. T. (2019). Effects of polyester microfibers on soil physical properties: Perception from a field and a pot experiment. *Science of the Total Environment*, 670, 1–7.

Zhao, S., Ding, G., Gao, Y., & Han, J. (2017). Approximating discrete probability distribution of image emotions by multi-modal features fusion. *Transfer*, 1000(1), 4669–4675.

17%

SIMILARITY INDEX

11%

INTERNET SOURCES

15%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Yashfa Tanveer, Humaira Yasmin, Asia Nosheen, Sajad Ali, Ajaz Ahmad. "Ameliorative effects of plant growth promoting bacteria, zinc oxide nanoparticles and oxalic acid on *Luffa acutangula* grown on arsenic enriched soil", *Environmental Pollution*, 2022
Publication 1%
- 2 Submitted to Higher Education Commission Pakistan
Student Paper 1%
- 3 Yashfa Tanveer, Saman Jahangir, Zafar Abbas Shah, Humaira Yasmin et al. "Zinc oxide nanoparticles mediated biostimulant impact on cadmium detoxification and in silico analysis of zinc oxide-cadmium networks in *Zea mays L. regulome*", *Environmental Pollution*, 2023
Publication 1%
- 4 link.springer.com
Internet Source 1%
- 5 peerj.com

1 %

6

Sidra Shoukat, Ayesha Tassarwar, Rumana Keyani, Mohsin Zafar et al. "Exogenous application of sodium hydrosulfide and salicylic acid mitigate salinity stress in maize by regulating ionic balance, biochemical attributes, photosynthetic pigments and some key antioxidants", South African Journal of Botany, 2023

Publication

1 %

7

Mirza Hasanuzzaman, Masayuki Fujita. "Selenium Pretreatment Upregulates the Antioxidant Defense and Methylglyoxal Detoxification System and Confers Enhanced Tolerance to Drought Stress in Rapeseed Seedlings", Biological Trace Element Research, 2011

Publication

1 %

8

www.researchgate.net

Internet Source

1 %

9

www.frontiersin.org

Internet Source

<1 %

10

ebin.pub

Internet Source

<1 %

11

"Microbial BioTechnology for Sustainable Agriculture Volume 1", Springer Science and

<1 %

Business Media LLC, 2022

Publication

12

Nahar, Kamrun, Mirza Hasanuzzaman, Md. Mahabub Alam, and Masayuki Fujita.

"Glutathione-induced drought stress tolerance in mung bean: Coordinated roles of the antioxidant defense and methylglyoxal detoxification systems", *AoB Plants*, 2015.

Publication

13

Sana Abid, Arshad Farid, Rameesha Abid, Mujeeb Rehman et al. "Identification, Biochemical Characterization, and Safety Attributes of Locally Isolated *Lactobacillus fermentum* from *Bubalus bubalis* (buffalo) Milk as a Probiotic", *Microorganisms*, 2022

Publication

14

"Wheat Production in Changing Environments", *Springer Science and Business Media LLC*, 2019

Publication

15

Ali S. Alqahtani, Fahd A. Nasr, Mohammad Z. Ahmed, Muath Y. Bin Mansour et al. "protective and anti-inflammatory effects of and its flavonoids profile ", *Open Chemistry*, 2023

Publication

16

Jun Nie, Shixiang Fan, Xu Luo. "First Account of the Breeding Biology of Indian Blue Robin

<1 %

<1 %

<1 %

<1 %

<1 %

(*Larvivora brunnea*) in Southwest China",
Animals, 2023

Publication

17

www.scielo.br

Internet Source

<1 %

18

PRASHANT SINGH, Menka Tiwari. "Bolstering Wheat's Immunity: BABA-Mediated Defense Priming Against *Bipolaris sorokiniana* Amid Competition", Cold Spring Harbor Laboratory, 2024

Publication

<1 %

19

prp.hec.gov.pk

Internet Source

<1 %

20

www2.mdpi.com

Internet Source

<1 %

21

aem.asm.org

Internet Source

<1 %

22

www.researchsquare.com

Internet Source

<1 %

23

Xiaoxia Yang, Rui Liu, Miao Jing, Ningning Zhang, Cuiying Liu, Jiakun Yan. "Variation of Root Soluble Sugar and Starch Response to Drought Stress in Foxtail Millet", Agronomy, 2023

Publication

<1 %

24

bmcchem.biomedcentral.com

Internet Source

<1 %

25

journals.plos.org

Internet Source

<1 %

26

repository.brieflands.com

Internet Source

<1 %

27

www.ncbi.nlm.nih.gov

Internet Source

<1 %

28

Fatima Bibi, Noshin Ilyas, Muhammad Arshad, Azeem Khalid, Maimoona Saeed, Sabah Ansar, Jacqueline Batley. "Formulation and efficacy testing of bioorganic fertilizer produced through solid-state fermentation of agro-waste by *Burkholderia cenocepacia*", *Chemosphere*, 2021

Publication

<1 %

29

Umakanta Sarker, Shinya Oba. "Catalase, superoxide dismutase and ascorbate-glutathione cycle enzymes confer drought tolerance of *Amaranthus tricolor*", *Scientific Reports*, 2018

Publication

<1 %

30

real.mtak.hu

Internet Source

<1 %

31

www.e3s-conferences.org

Internet Source

<1 %

32	www.nisshinbo.co.jp Internet Source	<1 %
33	Min-Ji Kim, Ramalingam Radhakrishnan, Sang-Mo Kang, Young-Hyun You, Eun-Ju Jeong, Jong-Guk Kim, In-Jung Lee. "Plant growth promoting effect of <i>Bacillus amyloliquefaciens</i> H-2-5 on crop plants and influence on physiological changes in soybean under soil salinity", <i>Physiology and Molecular Biology of Plants</i> , 2017 Publication	<1 %
34	edubirdie.com Internet Source	<1 %
35	krishikosh.egranth.ac.in Internet Source	<1 %
36	law.resource.org Internet Source	<1 %
37	patents.justia.com Internet Source	<1 %
38	publications.aston.ac.uk Internet Source	<1 %
39	www.coursehero.com Internet Source	<1 %
40	www.publish.csiro.au Internet Source	<1 %

41

www.scirp.org

Internet Source

<1 %

42

"Microorganisms in Saline Environments: Strategies and Functions", Springer Science and Business Media LLC, 2019

Publication

<1 %

43

Benchao Hou, Xiaomin Wang, Zhijian He, Haiyun Liu. "Integrative approach using network pharmacology, bioinformatics, and experimental methods to explore the mechanism of cantharidin in treating colorectal cancer", Naunyn-Schmiedeberg's Archives of Pharmacology, 2024

Publication

<1 %

44

Jaya Nepal, Xiaoping Xin, Gabriel Maltais-Landry, Wiqar Ahmad et al. "Carbon nanomaterials are a superior soil amendment for sandy soils than biochar based on impacts on lettuce growth, physiology and soil biochemical quality", NanoImpact, 2023

Publication

<1 %

45

K Adachi, I Suzuki. "Purification and properties of glutamate synthase from *Thiobacillus thioparus*", Journal of Bacteriology, 1977

Publication

<1 %

46

M. Farooq. "Plant drought stress: effects, mechanisms and management", *Agronomy for Sustainable Development*, 01/2009

Publication

<1 %

47

Patrícia F. Ávila, Allan H.F. de Mélo, Rosana Goldbeck. "Cello-oligosaccharides production from multi-stage enzymatic hydrolysis by lignocellulosic biomass and evaluation of prebiotic potential", *Innovative Food Science & Emerging Technologies*, 2023

Publication

<1 %

48

Rafaela Mendonça Nozari, Luisa Machado Ramos, Laura Alcântara da Luz, Rafael Nolibos Almeida et al. "Halotolerant *Streptomyces* spp. induce salt tolerance in maize through systemic induction of the antioxidant system and accumulation of proline", *Rhizosphere*, 2022

Publication

<1 %

49

Simona Gabrielli, Lucia Fontanelli Sulekova, Federica Furzi, Elisa Biliotti et al. "Molecular characterization of *Blastocystis* subtypes in HIV-positive patients and evaluation of risk factors for colonization", *Research Square Platform LLC*, 2019

Publication

<1 %

50

jm.um.ac.ir

Internet Source

<1 %

51

ntp.niehs.nih.gov

Internet Source

<1 %

52

test.techscience.com

Internet Source

<1 %

53

worldwidescience.org

Internet Source

<1 %

54

www.iamm.fr

Internet Source

<1 %

55

www.teses.usp.br

Internet Source

<1 %

56

Noshin Ilyas, Komal Mumtaz, Nosheen Akhtar, Humaira Yasmin et al. "Exopolysaccharides Producing Bacteria for the Amelioration of Drought Stress in Wheat", Sustainability, 2020

Publication

<1 %

57

Silva, E.N.. "Photosynthetic changes and protective mechanisms against oxidative damage subjected to isolated and combined drought and heat stresses in Jatropha curcas plants", Journal of Plant Physiology, 20100915

Publication

<1 %

58

B. Hameeda, O. P. Rupela, Gopal Reddy, K. Satyavani. "Application of plant growth-promoting bacteria associated with composts and macrofauna for growth promotion of

<1 %

Pearl millet (*Pennisetum glaucum* L.)", Biology and Fertility of Soils, 2006

Publication

59

Ingunn Stensland, John R. Pluske. "Use of probiotics and prebiotics in pig nutrition in the post-weaning period", Burleigh Dodds Science Publishing Limited, 2017

Publication

<1 %

60

Raju Bheemanahalli, Ranadheer Reddy Vennam, Purushothaman Ramamoorthy, K. Raja Reddy. "Effects of post-flowering heat and drought stresses on physiology, yield, and quality in maize (*Zea mays* L.)", Plant Stress, 2022

Publication

<1 %

61

journalcra.com
Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On